



# Comprehensive Guide for First Aid & CPR



# Table of Contents

<b>The Fundamental Principles of the International Red Cross And Red Crescent Movement</b>	<b>01</b>		
<b>01 THE RED CROSS</b>	<b>02</b>		
Our History.....	03		
The Fundamental Principles.....	03		
Red Cross Symbols.....	04		
Who We Are.....	05		
How We Help.....	05		
<b>02 RESPONDING TO EMERGENCIES</b>	<b>07</b>		
Preparing to Respond.....	08		
Preparing for Emergencies at Home.....	09		
Recognizing that an Emergency Exists.....	11		
Willingness to Act.....	11		
Legal Issues Around First Aid.....	12		
Good Samaritan Laws.....	12		
Workplace First Aiders.....	12		
Getting Permission to Help.....	12		
Duty to Report Child Abuse or Neglect.....	13		
Responding to Disclosures of Violence.....	13		
The Emergency Medical Services System.....	14		
Chain of Survival Behaviours.....	14		
Understanding Your Role as a First Aider.....	14		
Understanding Your Role in the EMS System.....	15		
Activating EMS.....	15		
When to Call EMS/9-1-1.....	16		
Poison Control Centres.....	16		
Calling EMS/9-1-1 for a Child in Your Care.....	17		
After an Emergency.....	17		
Lowering the Risk of Infection.....	18		
How to Prevent Diseases from Spreading.....	19		
Personal Precautions.....	19		
Equipment Precautions.....	20		
Environmental Precautions.....	22		
<b>03 CHECK, CALL, CARE</b>	<b>24</b>		
Check.....	25		
Call.....	28		
Care.....	30		
Secondary Assessment.....	33		
Shock.....	36		
Providing First Aid for Someone with a Disability	37		
<b>04 CHOKING</b>	<b>40</b>		
Partial Choking.....	43		
Complete Choking.....	44		
Responsive Adult or Child over 12 Months.....	44		
If You Are By Yourself and Choking.....	44		
Responsive Baby (Less Than 12 Months).....	47		
Unresponsive Choking.....	48		
<b>05 CIRCULATION EMERGENCIES</b>	<b>49</b>		
Cardiovascular Disease.....	50		
Heart Attack.....	51		
Angina.....	53		
Stroke.....	53		
Transient Ischemic Attack (TIA).....	55		
Life-Threatening Bleeding.....	55		
Life-Threatening External Bleeding.....	55		
Using a Tourniquet.....	57		
Using Hemostatic Dressings.....	58		
Life-Threatening Internal Bleeding.....	58		
<b>06 CPR &amp; AED</b>	<b>59</b>		
Cardiac Arrest.....	60		
Cardiac Arrest, Adult (After the Onset of Puberty)....	61		
Cardiac Arrest, Child (1 Year to the Onset of Puberty)....	62		
Cardiac Arrest, Baby (Less Than 12 Months).....	63		
Continue CPR Until.....	64		
What to Do if the Rescue Breaths Don't Go In.....	65		
Compression-Only CPR.....	65		
CPR Compression Depth.....	66		
Chest Compression Rates.....	66		
Chest Recoil.....	66		
Decreasing Compression Fraction (Off-Chest Time)....	67		
Special Considerations.....	67		
CPR for a Pregnant Woman.....	67		
Air in the Stomach.....	67		
Vomiting.....	67		
Mouth-to-Nose Breathing.....	69		
Mouth-to-Stoma Breathing.....	69		
Automated External Defibrillation.....	69		
Using an AED.....	70		
Medical Direction and AEDs.....	73		
AED Maintenance.....	74		
Working as a Team.....	74		
<b>07 BREATHING EMERGENCIES</b>	<b>75</b>		
Respiratory Distress.....	76		
Hyperventilation.....	77		
Asthma.....	78		
Using an Inhaler.....	80		
Using an Inhaler with a Spacer.....	80		
Allergic Reactions.....	80		
Anaphylaxis.....	82		

# Table of Contents

<b>08 WOUND CARE</b>	<b>85</b>	<b>10 BONE, MUSCLE, AND JOINT INJURIES</b>	<b>117</b>	<b>13 POISONS</b>	<b>154</b>
Bandaging Guidelines.....	86	Splinting Guidelines.....	120	Identifying Hazardous Materials.....	156
Triangular Bandages.....	86	Osteoporosis.....	124	Swallowed Poisons.....	158
Cuts and Scrapes.....	87			Inhaled Poisons.....	159
Puncture Wounds.....	88			Carbon Monoxide Poisoning.....	159
Scalp Injuries.....	88	<b>11 SUDDEN MEDICAL EMERGENCIES</b>	<b>125</b>	Absorbed Poisons.....	160
Infection.....	89	Fainting.....	126	Poison Ivy, Sumac, and Oak.....	161
Burns.....	90	Diabetic Emergencies.....	127	Giant Hogweed and Wild Parsnip.....	162
Thermal Burns.....	91	Using a Glucagon Kit.....	129	<b>Injected Poisons.....</b>	<b>163</b>
Chemical Burns.....	93	Seizures.....	129	Poisoning Caused by Alcohol or Drugs.....	164
Electrical Burns.....	93	Febrile Seizures.....	132	Drug Overdose.....	164
Radiation Burns.....	94	Mental Health Crisis.....	132	Opioid Overdose.....	165
Special Considerations When Caring for Burns.....	94	Self-Inflicted Injuries.....	133	Alcohol Poisoning.....	165
Bruises.....	94	Childbirth.....	133	<b>Bites and Stings.....</b>	<b>166</b>
Splinters.....	95	Miscarriage.....	135	Animal Bites.....	166
Nosebleeds.....	96			Rabies.....	167
Foreign Objects in the Nose.....	97	<b>12 ENVIRONMENTAL ILLNESS</b>	<b>136</b>	Insect Stings.....	167
Knocked-Out Teeth.....	97	How Body Temperature is Controlled.....	137	Tick Bites.....	168
Eye Injuries.....	98	Heat-Related Illnesses.....	139	Lyme Disease.....	169
Ear Injuries.....	100	Heat Cramps.....	140	Snakebites.....	170
Impaled Objects.....	101	Heat Exhaustion.....	140	Spider Bites.....	172
Amputations.....	102	Heat Stroke.....	142	<i>Black Widow Spiders</i> .....	172
Crush Injuries.....	103	<b>Cold-Related Illnesses.....</b>	<b>145</b>	<i>Brown Recluse Spiders</i> .....	172
Chest Injuries.....	103	Frostbite.....	145	Stings from Marine Life.....	173
Penetrating Chest Injuries.....	104	Hypothermia.....	147		
Blunt Chest Injuries.....	105	Freezing of Skin to Metal Objects.....	150		
Abdominal Wounds.....	105	Snow Blindness.....	150		
Blast Injuries.....	106	Cold-Water Immersion.....	151		
<b>09 HEAD, NECK, AND SPINAL INJURIES</b>	<b>107</b>	Drowning.....	152		
The Neck and Spine.....	108	Responsive Drowning Person.....	152		
The Brain.....	109	Unresponsive Drowning Person.....	153		
Concussion.....	112	Seasickness.....	153		
Shaken Baby Syndrome.....	115				

# Table of Contents

<b>14 CHILDHOOD ILLNESSES</b>	<b>175</b>	
Protecting Children from Infection and Disease	176	
Questions to Ask Yourself if You Think a Child May Be Ill.....	177	
When to Call the Parent of a Child in Your Care	178	
Common Childhood Conditions: Guidelines for Care.....	178	
Comparison Chart for Epiglottitis and Croup.....	187	
Childhood Fevers.....	188	
Taking a Child's Temperature.....	188	
Complications Related to Fever.....	189	
Giving Medications.....	189	
When to Give Medications.....	189	
General Rules for Medication.....	190	
<i>Medication in the Eye</i> .....	190	
<i>Medication in the Nose</i> .....	190	
<b>15 CARING FOR CHILDREN</b>	<b>191</b>	
Toy and Equipment Safety.....	192	
Equipment for Babies.....	192	
Safe Use of Cribs.....	192	
Safe Use of Playpens.....	192	
Safe Use of High Chairs.....	192	
Safe Use of Baby Strollers.....	193	
Safe Use of Pacifiers.....	193	
Equipment for Children.....	193	
Safe Use of Child Bike Trailers.....	193	
Safe Use of Safety Gates.....	193	
Toys for Children and Babies.....	194	
Playground Safety.....	194	
Car Safety.....	195	
Safe Use of Car Seats.....	195	
Passenger Rules for Children: Safety Education.....	195	
<b>APPENDIX A – KIT CONTENTS</b>	<b>203</b>	
Example of an Emergency Supplies Kit.....	203	
Example of an Emergency Car Kit.....	203	
<b>APPENDIX B – MOVING AN ILL OR INJURED PERSON</b>	<b>204</b>	
Body Mechanics.....	205	
How to Move an Ill or Injured Person.....	205	
Stretchers and Lifting Devices.....	208	
<b>APPENDIX C – ANATOMY AND PHYSIOLOGY</b>	<b>211</b>	
Body Systems.....	211	
Interrelationships of Body Systems.....	211	
Respiratory System.....	215	
Circulatory System.....	216	
Nervous System.....	217	
Musculoskeletal System.....	218	
<i>Muscles</i> .....	218	
<i>Skeleton</i> .....	219	
Integumentary System.....	220	
Digestive System.....	221	
Urinary System.....	222	
Reproductive System.....	223	
<b>GLOSSARY</b>	<b>224</b>	
<b>ACKNOWLEDGEMENTS</b>	<b>228</b>	

# The Fundamental Principles of the International Red Cross and Red Crescent Movement

## Humanity

The International Red Cross and Red Crescent Movement, born of a desire to bring assistance without discrimination to the wounded on the battlefield, endeavours in its international and national capacity to prevent and alleviate human suffering wherever it may be found. Its purpose is to protect life and health and ensure respect for the human being. It promotes mutual understanding, friendship, co-operation, and lasting peace amongst all peoples.

## Impartiality

It makes no discrimination as to nationality, race, religious beliefs, class, or political opinions. It endeavours to relieve the suffering of individuals, being guided solely by their needs, and to give priority to the most urgent cases of distress.

## Neutrality

In order to continue to enjoy the confidence of all, the Movement may not take sides in hostilities or engage at any time in controversies of a political, racial, religious, or ideological nature.

## Independence

The Movement is independent. The National Societies, while auxiliaries in the humanitarian services of their governments and subject to the laws of their respective countries, must always maintain their autonomy so that they may be able at all times to act in accordance with the principles of the Movement.

## Voluntary Service

It is a voluntary relief movement not prompted in any manner by desire for gain.

## Unity

There can be only one Red Cross or one Red Crescent Society in any one country. It must be open to all. It must carry on its humanitarian work throughout its territory.

## Universality

The International Red Cross and Red Crescent Movement, in which all Societies have equal status and share equal responsibilities and duties in helping each other, is worldwide.

The Fundamental Principles were proclaimed by the XXth International Conference of the Red Cross, Vienna, 1965. This is the revised text contained in the Statutes of the International Red Cross and the Red Crescent Movement, adopted by the XXVth International Conference of the Red Cross, Geneva, 1986.

In keeping with the Fundamental Principles of the Red Cross, the Society is committed to Social Justice in the elimination of Society structures and actions that oppress, exclude, limit, or discriminate on the basis of race, gender, ethnicity, financial ability, sexual orientation, religion, disability, or age.



# 1 The Red Cross

## Our History



In June of 1859, Henry Dunant saw an unforgettable scene: 40,000 dead and wounded soldiers left on the field after the Battle of Solferino in Italy. Dunant organized local villagers into first aid teams to help as many of the wounded as possible, saving thousands of lives.

To prevent this horror from happening again, Dunant decided to create a neutral organization to care for wounded soldiers and prisoners—an organization that would be respected and protected by both sides in any conflict. The result was the Red Cross. Dunant spent the rest of his life trying to reduce the suffering caused by war. He lobbied governments, organized Red Cross Societies in different countries, and spoke to the public.

In the spring of 1885, during Louis Riel's Northwest Rebellion, Dr. George Sterling Ryerson stitched a Red Cross made of cotton onto white material and used this as a flag to distinguish the horse drawn wagon being used to transport the wounded. This was one of the first Red Cross flags ever flown in Canada.

In 1896, Dr. Ryerson founded the first overseas branch of the British Red Cross, which later became the Canadian Red Cross.

In 1901, Dunant won the first Nobel Peace Prize. By founding what is now the International Red Cross and Red Crescent Movement, he has saved the lives of millions of people.



On May 19, 1909, *The Canadian Red Cross Society Act* was approved by Parliament, and decreed that the Canadian Red Cross would serve as an auxiliary to the Government of Canada for Canadians, which was a measure that was in accordance with the Geneva Conventions. (Prior to 1909, the Canadian Red Cross operated as a branch of the British Red Cross.) As a result, for more than a century, Canadians have had their own national Red Cross Society dedicated to improving the situations of the most vulnerable people in Canada and around the world.

Today, there are Red Cross or Red Crescent Societies in more than 190 countries, all dedicated to the same ideals and fundamental principles.

## The Fundamental Principles

In every country, our programs and activities are guided by seven Fundamental Principles. The Tanzanian Red Cross has created a short, simple version of these principles:

**Humanity:** We serve people, but not systems.

**Impartiality:** We care for the victims and the aggressors alike.

**Neutrality:** We take initiatives, but never take sides.

**Independence:** We bow to needs, but not rulers.

**Voluntary Service:** We work around the clock, but never for personal gain.

**Unity:** We have many talents, but a single idea.

**Universality:** We respect nations, but our work knows no bounds.

Essentially, we provide help to people in need, whatever their race, gender, culture, ethnicity, financial ability, sexual orientation, religion, disability, or age.

### Red Cross Symbols

The International Red Cross and Red Crescent Movement has three official symbols: the Red Cross, Red Crescent, and Red Crystal.

All three symbols are equivalent and share the same status under international law. Each national society of the International Red Cross and Red Crescent Movement chooses one of these three symbols to use in their country.

The three emblems are used to identify military medical services as well as the people, programs, and objects connected with the activities of the International Red Cross and Red Crescent Movement.

Recognized as a symbol of humanity and neutrality, the emblem is the only protection Red Cross workers have when they bring relief and protection to victims of armed conflicts. The emblem identifies and protects the sick, the wounded and those who come to their aid, and medical equipment, buildings, and vehicles.



#### The Red Cross


The Red Cross symbol—five red, equal-sized squares forming a cross resting on a white background—was adopted in 1863 at the first International Conference of what would one day be the Red Cross and Red Crescent Society. Today it is one of the most recognized emblems in the world, and continues to be an internationally respected symbol of protection and neutrality.

The Red Cross is the emblem that is used by the Canadian National Society. No organization—except the Canadian Red Cross and the medical services of the armed forces—has the right to use the Red Cross emblem in Canada. The Red Cross emblem must be readily recognized and respected around the world as a trusted symbol of protection and humanitarian aid. Its use is legislated by the Geneva Conventions Act, the Trade Marks Act, and the Canadian Red Cross Society Act.



#### The Red Crescent

Although the Red Cross emblem has no religious, political, or cultural associations, it was sometimes believed to be related to the Christian cross, and this association was problematic in certain regions and conflicts. In 1929, the International Federation of the Red Cross agreed to the adoption of an additional, equivalent emblem: The Red Crescent. This emblem could be used by any national society in place of the Red Cross.



#### The Red Crystal

In December of 2005, the International Red Cross and Red Crescent Movement welcomed the decision to create an additional emblem alongside the Red Cross and Red Crescent: the Red Crystal.

## Who We Are

### Our Mission

The mission of the Canadian Red Cross is to improve the lives of vulnerable people by mobilizing the power of humanity in Canada and around the world.

### Our Vision

The Canadian Red Cross is the leading humanitarian organization through which people voluntarily demonstrate their caring for others in need.

### Our Values

Our actions and decisions are based on:

- Humanitarian values, as expressed in our Fundamental Principles;
- Respect, dignity, and care for one another within and outside the Canadian Red Cross; and
- Integrity, accountability, effectiveness, and transparency.

### Our Volunteers

The Canadian Red Cross relies on more than 25,000 volunteers across the country to fulfill its mission. Red Cross volunteer opportunities are dynamic and varied, with engagement levels varying by geographic location and program. For more information on volunteering, visit [redcross.ca](http://redcross.ca) or contact your local Red Cross office.

## How We Help

The Canadian Red Cross provides help in many ways:



### Disaster Management

While many associate the Canadian Red Cross with its disaster relief efforts around the world, each year thousands of Canadians receive aid following emergencies such as house fires, flooding, ice storms, and even chemical

spills. Following a disaster, Canadian Red Cross volunteers ensure those affected have access to shelter, food, clothing, hygiene items, and emotional support. As part of a global network of 190 national Societies, the work and expertise of the Canadian Red Cross often extends beyond Canada's borders. Trained Canadian personnel and resources, including an Emergency Response Unit specializing in health, are poised for deployment within hours of a disaster.



### International Operations

The Canadian Red Cross works around the world to protect and promote the rights of all people affected by armed conflict. The Geneva Conventions, laws that protect the wounded, the sick, prisoners of war, and civilians, serve as a mandate for the Canadian Red Cross

in its mission to provide assistance to those in need. The Canadian Red Cross educates Canadians on international issues including humanitarian law and development efforts. The Society recruits and trains qualified Canadian delegates for international field operations in countries affected by conflict or disaster or for long-term development missions.



## First Aid Programs

As the largest provider of first aid training in Canada, the Canadian Red Cross has been offering first aid and CPR training for more than 50 years. With courses for individuals, groups, workplaces, and first responders, participants learn how to prevent injuries and to think, react, and respond in emergency situations.



## Swimming and Water Safety Program

Since 1946, the Canadian Red Cross has been teaching Canadians how to stay safe around the water through our Swimming and Water Safety programs. Water safety activities are offered at community pools, beaches, schools and summer camps to help people of all ages and abilities gain knowledge and skills in every aspect of water safety.



## Respect Education

The Canadian Red Cross aims to prevent abuse, neglect, harassment, and interpersonal violence to ensure everyone's right to live, play, and learn in a safe environment. Nationally recognized, award-winning programs teach children, youth, and adults about healthy relationships,

the signs of abuse, dating violence, personal safety, bullying prevention, and how to get help. Prevention educators with extensive training in abuse prevention deliver workshops to school and community groups, sports teams, coaches, educators, and parents or guardians. The aim is to stop the hurt before it starts.



## Community Health and Wellness

Canadian Red Cross staff and volunteers work together to provide a variety of services that help people remain living independently in their homes with comfort and dignity, despite injury, illness, or other circumstances. Nutrition-based programs, such as Meals on

Wheels, provide balanced meals that eliminate the guesswork for clients and caregivers in meeting daily dietary requirements; transportation services ensure people in the community get to and from medical appointments, the grocery store, and more; and health equipment loans provide short-term loans of clean, safe medical equipment to people who are recovering from injury, illness, or surgery. Our programs are designed to enhance quality of life, support independence, and provide peace of mind. The services offered in each area vary according to the needs of the community.



## 2 Responding to Emergencies

## Preparing to Respond

Make sure you have easy access to items that will help you respond to an emergency. Keep a first aid kit in your home and vehicle, and ensure that you know the location of the first aid kit and automated external defibrillator (AED) in your workplace. Download the Canadian Red Cross First Aid App to your mobile device so that you always have a first aid reference readily available.



You can purchase first aid kits and supplies from the Canadian Red Cross online store ([products.redcross.ca](http://products.redcross.ca)), a drug store, or a medical supplies store. Whether you buy a first aid kit or assemble one yourself, make sure it has all of the items you may need. Perform regular maintenance and safety checks on all first aid kits and equipment. Remove or replace any items that are broken, expired, discoloured, or contaminated (e.g., a sterile tool in a torn package). You should also decontaminate any reusable equipment that has been used (e.g., a stretcher).



All workplace first aid kits must meet provincial/territorial legislation and occupational health and safety guidelines. Go to [redcross.ca/firstaidlegislation](http://redcross.ca/firstaidlegislation) for more details.

### First aid kits should include at least the following supplies:

- Emergency telephone numbers for EMS/9-1-1, your local Poison Control Centre, and your personal doctor(s)
- Home and office phone numbers for family members, friends, or neighbours who can help in an emergency
- Sterile gauze pads (dressings) in small and large squares to place over wounds
- Adhesive tape
- Roller and triangular bandages to hold dressings in place or make slings
- Adhesive bandages in assorted sizes
- Scissors
- Tweezers
- Safety pins
- Cold packs or instant ice packs
- Disposable non-latex gloves, such as surgical or examination gloves
- Flashlight, with extra batteries in a separate bag
- Antiseptic wipes or soap
- Pencil and pad
- Emergency blanket
- Eye patches
- Thermometer
- Barrier devices, such as a pocket mask or face shield
- Canadian Red Cross first aid manual

## Preparing For Emergencies at Home

To be ready for an emergency at home:

- Keep important information about yourself and your family in a handy place. Include your address, everyone's date of birth, health card numbers, medical conditions, allergies, and prescriptions and dosages. List the names and phone numbers of your doctors.
- Post the numbers for the police, fire department, emergency medical services (EMS), and Poison Control Centre near every phone in your home and save these numbers in your mobile phone's contacts list.
- Teach children how to call for help.
- Install smoke and carbon monoxide detectors. Test them regularly.
- Keep first aid kits handy in your home and car. Make sure everyone in the family knows where they are and what is in them.
- Review and practice the first aid skills that you learn so that you will have the confidence and knowledge to respond in an emergency.

Most communities are served by the emergency telephone number 9-1-1. If your community does not operate on a 9-1-1 system, search online or in your local phone directory for the numbers of the police department, fire department, and EMS system. Also include the number for your provincial or territorial Poison Control Centre's hotline on your list. Teach everyone in your home how and when to use these numbers.


Take steps to make it easier for EMS personnel and others to help you, should an emergency occur:

- Make sure your house or apartment number is large, easy to read, and well-lit at all times. Numerals (e.g., "87") are easier to see and read than words (e.g., "eighty-seven").
- Keep relevant medical information, such as a list of the medications that each family member takes, in an accessible place at home and when you go out (e.g., on the refrigerator door and in your wallet or mobile phone).

Mobile phones and other electronic devices can display emergency contacts. The letters "ICE" (for "In Case of Emergency") may appear in the phone's contacts list next to the name of a doctor, spouse, or other important contact. Assigning "ICE" to a contact allows emergency personnel to reach someone who knows important facts about the ill or injured person. If the mobile phone has been programmed to include the ICE contact with a "0" in front, it will appear as the first contact in the person's list.

If you have a chronic medical condition such as diabetes, epilepsy, or allergies, consider wearing a medical identification product to alert EMS personnel of your condition in case you are unable to do so. There are also applications you can use to create digital identification tags in your mobile phone. These tags can be viewed even when your phone is locked, and so can provide important information in an emergency (such as your medical conditions, blood type, and emergency contact details).

In a life-threatening emergency, every second counts. By preparing for emergencies, you can help ensure that care begins as soon as possible—for yourself, a family member, a co-worker, or a member of your community.



# Signs of an Emergency

## UNUSUAL SOUNDS

- Screaming, moaning, shouting, or calls for help
- Sudden loud noises such as breaking glass, clashing metal, or screeching tires
- A change in the sound made by machinery or equipment
- An unusual silence

## UNUSUAL ODOURS

- A foul or unusually strong chemical odour
- The smell of smoke
- The smell of gas
- An unrecognizable odour
- An inappropriate odour (e.g., a sickly-sweet odour on a person's breath)

## UNUSUAL SIGHTS

- A stopped vehicle on the roadside or a car that has run off of the road
- Downed electrical wires
- Sparks, smoke, or fire
- A person who suddenly collapses or is lying motionless
- Signs or symptoms of illness or injury, such as profuse sweating for no apparent reason or an uncharacteristic skin colour

## UNUSUAL BEHAVIOURS

- Confusion in a person who is normally alert
- Unusual drowsiness
- Personality or mood changes (e.g., agitation in a person who is normally calm)



## Recognizing that an Emergency Exists

It will sometimes be obvious that an emergency exists. For example, a scream or cry for help, an unpleasant or unusual odour, or the sight of someone bleeding severely or lying motionless on the ground are all clear indications that immediate action is needed. But other times, the signs of an emergency may be more subtle, such as a slight change in a person's normal appearance or behaviour, or an unusual silence.

## Willingness to Act

Sometimes people don't want to get involved in an emergency. The four most common reasons are:

- 1. The Bystander Effect:** If there are other people at the scene, it is easy to think that they can take care of the emergency without your help. However, you should never assume that someone has taken action and is providing first aid just because you see a lot of people. Remember that there are many important jobs that you can do. You can help control the crowd, direct the actions of bystanders, call EMS/9-1-1, get supplies, or provide care to the ill or injured person. If you are unsure of what to do, ask others at the scene how you can help.
- 2. Unpleasant injuries or illnesses:** Some people may feel faint, upset, or nauseated when they see blood, vomit, or visible injuries. If this happens to you, close your eyes or turn away for a moment and take a few deep breaths to calm yourself before you deal with the situation. If you are still unable to give care, you can volunteer to help in other ways, such as by calling EMS/9-1-1 and bringing necessary equipment and supplies to the scene.

**3. Catching a disease:** You might be concerned that performing first aid will put you at risk of infection, but there are many ways to reduce this risk. If you take simple precautions to limit contact with the ill or injured person—such as wearing gloves and using a CPR breathing barrier—you can limit the possibility of catching a disease. Remember that EMS personnel provide care for ill and injured people every day without incident.

**4. Doing something wrong, or causing more harm:** You might be afraid that you will be sued if you make a mistake. As long as you act reasonably, you don't need to worry. All provinces and territories have laws to protect bystanders who give emergency help. Getting trained in first aid can give you the confidence, knowledge, and skills you need to respond appropriately to an emergency. Use your good judgment and stay within the realm of the skills in which you were trained. Once you start giving first aid, keep providing help until EMS personnel arrive. If you are unsure of what to do, call EMS/9-1-1 and follow the EMS dispatcher's instructions. The most harmful thing you can do is to do nothing at all.

Thinking about these things now and mentally preparing yourself for an emergency will help you overcome your fears.

## Legal Issues Around First Aid

### Good Samaritan Laws

Good Samaritan laws, which protect First Aiders from financial liability, were developed to encourage people to help others in emergency situations. These laws assume a First Aider will do his or her best to save a life or prevent further injury. Good Samaritan laws require the First Aider to use common sense and a reasonable level of skill, as well as to give only the type of emergency care that is within his or her training. Good Samaritan laws usually protect First Aiders who act in the same manner as a "reasonable and prudent person" would in the same situation. For example, a reasonable and prudent person would:

- Move a person only if the person's life was in danger.
- Ask a responsive person (or the parent, guardian, or caregiver of a child or baby) for permission to help (consent) before giving care.
- Check a person for life-threatening conditions before giving further care.
- Call EMS/9-1-1.
- Continue to give care until more highly trained personnel take over.

If a First Aider's actions are grossly negligent or reckless, or if the First Aider abandons the person after starting care, Good Samaritan laws may not protect the First Aider.



You may be legally obligated to provide first aid as needed to any child or baby in your care.

### Workplace First Aiders

First aid in the workplace can be governed by both national and provincial/territorial legislation. A workplace First Aider has additional responsibilities and may be legally required to provide first aid in the workplace if the need arises. He or she may also need to know other information, including:

- Where emergency equipment is located.
- How to properly complete documentation following a workplace incident.
- What the workplace-specific emergency procedures are.
- How to call for help in an emergency.

Refer to the legislation that governs your workplace for more information.



### Getting Permission to Help

Once you decide to act, you need to get the ill or injured person's permission (also known as consent) to assist.

To get permission:

1. Tell the person who you are.
2. Tell the person that you are here to help.
3. Ask the person if that is okay.

If a baby or a child is ill or injured:

- Ask the parent, guardian, or caregiver for permission.
- If the child or baby is alone, you can assume you have permission to give first aid.



## Special Situations

There are times when it is difficult to get permission to provide care for an ill or injured person or when the person denies the help of First Aiders or bystanders:

### **The Person Is Unresponsive Or Confused, Or Has a Mental Impairment**

Someone who is unresponsive or confused, or who has a mental impairment, may not be able to grant permission. In these cases, the law assumes the person would give permission if he or she were able to do so. This is called implied consent. Implied consent also applies when a child or baby needs emergency medical assistance and his or her parent, guardian, or caregiver is not present.

### **The Person Refuses Care**

An ill or injured person may refuse care, even if he or she desperately needs it. A parent or guardian also may refuse care for his or her child or baby. You must honour the person's wishes. Explain to the person why you believe care is necessary, but do not touch or give care to the person, and never attempt to give the person help by force. Remember, you should never put yourself in any danger. Stay nearby, if possible, in case the person later decides to accept your help or becomes unresponsive.

If you believe the person's condition is life-threatening, call EMS/9-1-1 and communicate that a person requires medical assistance but is refusing help. The EMS personnel who arrive will deal with the situation. If the person gives consent initially but then withdraws it, stop giving care and call EMS/9-1-1 if you have not already done so.

### **You and the Person Do Not Speak the Same Language**

If you do not speak the same language as the injured or ill person, obtaining consent may be challenging. Find out if someone else at the scene can serve as a translator. If a translator is not available, do your best to communicate with the person by using gestures and facial expressions. When you call EMS/9-1-1, explain that you are having difficulty communicating with the person, and tell the dispatcher which language you believe the person speaks. The dispatcher may have someone available who can help with communication.

### **Duty to Report Child Abuse or Neglect**

Every adult in Canada has a legal duty to report child abuse or neglect, even if it is not confirmed. Information around the specific how-to-report details can be found in your jurisdiction's Child Protection Act, but the duty to report is uniform in all acts. If you are responding to a first aid emergency and you think a child is being harmed, then there needs to be a report to child protection and/or police. Even if you are unsure, child protection needs to be informed and needs to guide the next steps. Also share your concerns with the EMS personnel who respond to the situation.

### **Responding to Disclosures of Violence**

A person revealing experiences of abuse—past or present—can be challenging and upsetting. This is even more true if the individual receiving the disclosure is close to the person and/or knows the perpetrator. Your response to the disclosure is critical. Studies show that the manner in which a disclosure of interpersonal violence is handled is a significant factor in determining the psychological impact on the victim. Indifference and blame are damaging responses that can have long-lasting consequences.


If you receive a complaint of abuse, neglect, harassment, or bullying, you must know what steps need to be taken, when they need to occur, and how they need to be carried out. The complaint cannot be diminished or ignored. There must always be a response. Both verbal and non-verbal disclosures need to be handled sensitively while following specific procedures. If the disclosure is from a young person, your jurisdiction's Child Protection Act may govern how you respond.

### The Emergency Medical Services System

The emergency medical services (EMS) system is a network of community resources and trained personnel, organized to give emergency care in cases of injury or sudden illness. The system begins when someone sees an emergency and decides to take action by calling EMS/9-1-1. This action allows the EMS dispatcher to take down information about the emergency and provide it to the trained EMS personnel who will respond to the scene. Many EMS dispatchers are also trained to provide first aid and CPR instructions over the phone to assist the caller until EMS arrives.

### Chain of Survival Behaviours

The Chain of Survival Behaviours is a series of steps that help ensure a positive outcome for an ill or injured person. As a First Aider, your role is to prepare, recognize emergencies, provide first aid and/or access help.



### Understanding Your Role as a First Aider

First aid is the immediate care that is given to an ill or injured person until more advanced care can be obtained.



Your role as a First Aider includes four basic steps:

1. Recognize the emergency.
2. Protect yourself, the ill or injured person, and bystanders.
3. Access help (e.g., call EMS/9-1-1).
4. Act according to your skills, knowledge, and comfort level.

One of the simplest and most important ways of providing first aid is to call for help (EMS/9-1-1). By making this call, you ensure that the ill or injured person receives care from highly trained medical professionals.

Responding to the scene of an emergency involves providing emotional support as well as caring for injuries. When providing care, you should:

- Speak in a calm and reassuring manner. Ask for the ill or injured person's name and use it often.
- Communicate both verbally and nonverbally to reassure the person.
- Position yourself at eye level when you talk to the ill or injured person. Avoid unnecessary physical contact and any body language that could appear threatening.
- Actively listen to the ill or injured person. This involves four behaviours:
  1. Making every effort to fully understand what the person is trying to say.
  2. Repeating back to the person, in your own words, what the person said.
  3. Avoiding criticism, anger, or rejection of the person's statements.
  4. Using open-ended questions (i.e., avoid questions that can be answered with "Yes" or "No").

### ***Understanding Your Role in the EMS System***

The emergency medical services (EMS) system is a network of professionals linked together to provide the best care for people in all types of emergencies. As a member of the community, you play a major role in helping the EMS system to work effectively. Your role in the EMS system includes four basic steps:

1. Recognizing that an emergency exists
2. Deciding to help/take action
3. Activating the EMS system
4. Giving care until EMS personnel take over



### ***Who Is Coming to Help***

Emergency medical services (EMS) personnel have advanced training that allows them to provide medical care outside of the hospital setting. Depending on their level of training, EMS personnel have different roles and responsibilities. There are different kinds of trained personnel who may respond when a call is placed to EMS/9-1-1.

**First Responders:** These include police, firefighters, and job-specific personnel such as athletic trainers and workplace first aid response teams. They are trained in time-sensitive lifesaving measures. Unlike lay responders (First Aiders), first responders have a legal obligation to act in an emergency when they are on duty.

**Paramedics:** These are highly specialized emergency personnel whose skills include both basic and advanced life support. Once on the scene, these professionals will take over the care of the person, including transportation to a hospital or other facility if necessary.



## Activating EMS

In a life-threatening emergency, it is critical that someone activate EMS by calling EMS/9-1-1. Activating EMS will send emergency medical help on its way as quickly as possible. The sooner EMS personnel arrive, the higher the chances of a positive outcome.

EMS varies from community to community. Most people in Canada call 9-1-1 for help in emergencies, but in some areas of Canada and in many workplaces you may need to dial a different designated emergency number instead. If you live or work in an area where 9-1-1 is *not* the number to call in an emergency, make sure you know the designated emergency number to call.

Phone carriers are required to connect any 9-1-1 call made from a mobile phone, even if the phone does not have an active service plan. In most areas, you cannot text 9-1-1. Unless you have confirmed that the 9-1-1 call centre in your area supports texting, you should always call.



## When to Call EMS/9-1-1

At times, you may be unsure whether EMS personnel are needed. To make the decision, use your best judgment based on the situation, your assessment of the ill or injured person, and information gained from this course and other training you may have received. Trust your instincts. When in doubt, make the call. If you think that an emergency exists, it probably does; you should call EMS/9-1-1 for professional help immediately.

Call EMS/9-1-1 if there is a danger to you or others, or if the ill or injured person has any of the following conditions:

- Unresponsiveness or an altered mental state (e.g., confusion)
- Difficulty breathing or no signs of breathing
- Persistent chest pain or pressure
- Life-threatening or uncontrollable bleeding
- Seizures
- Severe headaches
- Abnormal speech
- Injuries to the head, neck, or back
- Blood in the vomit or urine
- An apparent mental health crisis
- Imminent childbirth

You should also call EMS/9-1-1 if the ill or injured person is not easily accessible, or if the situation involves any of the following:

- Fire
- An explosion
- A motor vehicle collision
- Hazardous materials
- Downed electrical wires
- Swift-moving water

## Poison Control Centres

If none of the conditions above apply and you suspect that poisoning is the cause of the person's condition, call poison control instead of EMS/9-1-1. If any of the above conditions are true (for example, a person who may have swallowed poison is having difficulty breathing), or you are unsure of whether to call EMS/9-1-1 or poison control, call EMS/9-1-1.

## Calling EMS/9-1-1 for a Child in Your Care

If you are a professional caregiver, you may have additional considerations when calling EMS/9-1-1 for a child in your care.

Once you have activated EMS, call another caregiver to come and stay with any other children while you attend to the ill or injured child. If the child is being taken to the hospital, call his or her parent or guardian and ask the parent or guardian to meet you there. If you are travelling in an ambulance, tell them you will call again when you know the name and location of the hospital and that you may not know this information until after you arrive. If you cannot go with the child in the ambulance, be sure to give the paramedics the child's medical information and your contact information.

## After an Emergency

Being involved in an emergency and providing first aid can be stressful. After the emergency is resolved, you may have lingering feelings such as uneasiness, doubt, anxiety, and fear. After dealing with an emergency, it is often helpful to talk to somebody about the situation.

Everyone reacts to stressful situations differently. It is very difficult to predict who will or will not be affected by abnormal events. Some individuals may feel better in the few days after the event. Others may find that it takes longer for their stress to go away, and that it begins to impact their relationships.

Consider seeking professional help if you experience any of the following for more than two weeks after the emergency:

- Crying fits or uncontrollable anger
- Trouble eating
- Trouble sleeping
- Loss of engagement with former interests
- Unusual or unexplained pain (e.g., headaches or stomach aches)

- Fatigue
- Feelings of guilt, helplessness, or hopelessness
- Staying away from family and friends
- Ignoring daily tasks, such as going to work

For help dealing with ongoing stress symptoms, contact your local crisis intervention line. You may also speak to your family doctor or a mental health professional.

## Documentation

If an incident occurs in a workplace (including a marine environment), it must be documented properly. The documentation may be paper and/or electronic. The reports you create are official legal documents: paper versions should be filled out in pen (not pencil), with any changes or corrections crossed out and initialed.

Good documentation is:

- Complete and accurate
- Legible
- Objective
- Completed as soon as possible (while your memory of the incident is fresh)

## Lowering the Risk of Infection

Giving first aid care is a hands-on activity that can put you in close contact with another person's bodily fluids (such as saliva, mucus, vomit, and blood), which may contain harmful pathogens (microorganisms that can cause disease or infection). Some pathogens pose a particular risk because of their long-term effects on the health of the infected person.

Pathogens spread from person to person through four modes of disease transmission. Some pathogens spread through only one of these routes, while others spread through several.

MODE OF TRANSMISSION		
	Definition	Example
<b>Direct Contact</b>	 Occurs when bodily fluids from an infected person enter another person's body	A cut on your hand comes into contact with an infected person's blood. Examples: HIV/AIDS, Hepatitis, Herpes
<b>Indirect Contact</b>	 Occurs when germs from an infected object or surface enter another person's body	You are not wearing gloves and you pick up a tissue that has been used by an infected person, and then you rub your eye. Examples: Meningitis, Influenza
<b>Airborne Transmission</b>	 Occurs when a person breathes in germs from the air	You are not wearing a face mask while providing first aid to an infected person. The infected person sneezes, sending germs into the air, and then you breathe them in. Examples: Measles, Tuberculosis
<b>Vector-Borne Transmission</b>	 Occurs when germs are introduced directly into the body	An infected mosquito bites you, injecting pathogens into your blood. Examples: Lyme Disease, Malaria

## How to Prevent Diseases from Spreading

### Personal Precautions

Personal precautions are the actions that individuals can take to reduce the risk of disease transmission. Examples include washing your hands frequently, treating all blood and other bodily fluids as infectious materials, covering your mouth and nose when sneezing, eating well, and getting enough exercise and sleep.

One way to significantly reduce your risk of infection is to inspect your hands regularly and cover even small cuts with adhesive bandages. This reduces the number of points where germs can enter your body.

### Handwashing

Proper handwashing is an important precaution for preventing the spread of germs that cause many infectious diseases. Frequent handwashing is important as part of a daily routine, for example, before eating or drinking and after using the washroom.

Wearing gloves helps to protect you from infection, but it does not eliminate the need for proper hygiene. First Aiders should always wash their hands thoroughly when they have finished giving care, even if they wore gloves the entire time.

Use the following guidelines when washing your hands:

1. Always use warm running water and a mild soap.



2. Create a lather by rubbing your hands together with the soap—friction with the soap suds is what actually removes the germs from hands. Rub your hands together vigorously for at least 30 seconds.



3. Wash all parts of each hand. Be sure to scrub your palms and wrists, between your fingers, under your fingernails, and around the backs your hands.



4. Dry your hands with a clean paper towel.



5. Turn the faucet off using the towel as a barrier between your hands and the faucet handle. Use the towel as a barrier for the door handle as well.



## When to Wash Your Hands

Wash your hands after touching anything that is likely to carry germs (e.g., after wiping your nose, changing a diaper, or touching a pet), and before touching anything that could carry germs into the body (e.g., before handling food or touching an open cut). It is also a good habit to wash your hands frequently, for example when returning home from a public place.

In a first aid emergency, remember to wash your hands:

- Before and after contact with an ill or injured person.
- After handling dirty articles, instruments, or dressings.
- Before or after treating wounds.
- After removing gloves.

## Hand Sanitizers

Hand sanitizers are not a substitute for hand washing, and are not suitable for cleaning visibly soiled hands. They should be used only as a temporary measure in situations where you are unable to wash your hands immediately. If you can see dirt on your hands, you should wash them with soap and water before using hand sanitizer.

When using an alcohol-based hand sanitizer to help decontaminate your hands, use the amount of product recommended by the manufacturer. Rub it thoroughly over all the surfaces of your hands, including your nails and in between your fingers, until the product dries.

## Immunization

Most Canadians have been immunized against common diseases such as measles and tetanus. Some vaccine-preventable diseases are serious, and can lead to disability or death. Immunizations protect not only you, but also your family, friends, and co-workers.

Immunization introduces a substance into the body that builds up its resistance to germs that cause a specific disease. Some vaccinations may wear away with time. As a result, it's possible that your immunity to a particular disease could decline. It is important to keep up with your immunization needs since protection against diseases through immunization is a lifelong process.

Some vaccines require more than one injection and a certain period of time between injections. If you are planning a trip outside the country, find out well before you leave which immunizations are recommended and required in order to safely go to the countries you will be visiting.

Because the risk of disease varies from place to place, this information cannot cover all the hazards that you might face. For specific guidelines on your particular situation, talk to your doctor or your community public health centre.

## Equipment Precautions

Personal protective equipment (PPE) is the general term for items that protect you from contact with pathogens and contaminated objects. Examples include barrier devices such as safety glasses, goggles, face masks, CPR breathing barriers, and gloves. You should always use some type of barrier device between yourself and any material that could pose the risk of infection.

Any disposable equipment that you have used (e.g. gloves, breathing barriers) should be immediately disposed of in a labelled leak-proof container with a biohazard symbol.



## Gloves

Wearing gloves is an important way to protect yourself while giving first aid. Most gloves found in first aid kits are disposable and latex-free, as some people are allergic to latex. Disposable gloves are meant to be worn once and then discarded; never wash or reuse disposable gloves. Gloves should fit properly. You should throw out any gloves that are discoloured, torn, or punctured.

Wear gloves when:

- You are providing care, especially if there is a possibility that you will come into contact with another person's blood or other potentially infectious material.
- You must handle items or surfaces soiled with blood or other potentially infectious materials.

When you are wearing gloves, try to limit how much you touch other surfaces. Avoid touching anything while wearing soiled gloves as pathogens can easily transfer from that surface to another person. If possible, remove soiled gloves and replace them with a clean pair before touching other surfaces or equipment in your first aid kit.

Consider asking people to begin helping themselves while you retrieve and/or put on your gloves. For example, a bleeding person can use his or her own hand to apply pressure to the wound. This can also be effective in situations where gloves are not available.

When you are finished providing care, remove your gloves using the proper technique to reduce the risk of contaminating your skin. Dispose of the gloves properly and wash your hands. When multiple people are in need of care, remove your gloves, wash your hands, and put on a clean pair before assisting the next person.

## Proper Technique for Removing Gloves

To remove soiled gloves without contaminating your skin:

1. With one hand, pinch the glove at the wrist of your second hand, being careful to touch only the glove's outer surface.
2. Pull the glove down and off your hand.
3. Form the glove into a ball and hold it in the palm of your gloved hand.



4. Insert your gloveless thumb under the glove of your other hand at the inner wrist.
5. Pull the glove downward off the hand, turning it inside out and trapping the other balled glove inside.



6. Discard gloves appropriately.
7. Wash your hands thoroughly with soap and water for at least 30 seconds, using the proper handwashing technique.





## CPR Breathing Barriers

CPR breathing barriers are used to reduce the risk of infection when giving rescue breaths by eliminating the need for mouth-to-mouth contact, protecting you from bodily fluids such as saliva and blood.

The most basic and portable type of breathing barrier is a face shield—a flat piece of thin plastic that is placed over a person's face, with an opening or valve that allows you to breathe into the person's mouth.

A pocket mask is a transparent, flexible device that creates a tight seal over the person's nose and mouth. The opening contains a filter or a valve that protects you from coming into contact with the person's bodily fluids and exhaled air.

Breathing barriers sized specifically for children and babies are available. If possible, select equipment that is appropriately sized for the injured or ill person.

## Environmental Precautions

Environmental precautions refer to the systems that are implemented in an area to reduce exposure to germs. Examples include making sure there is proper ventilation in the workplace, that people don't use the same sink for hygiene and food preparation, and that any contaminated materials are immediately disposed of in a dedicated biohazard garbage.

Designate a container for sharps (potentially infected sharp objects such as used needles and lancets). It should be puncture-resistant, leak-proof, and labelled with warning signs such as the biohazard symbol. To avoid infection, put all needles and similar sharp objects into this container immediately after they are used. It should not be used for other purposes.



Many environmental precautions are the responsibility of the employer, but you can make suggestions if you notice something that could be improved.




## Cleaning and Disinfecting Surfaces and Equipment

Reusable equipment and surfaces that have been contaminated by blood or other potentially infectious materials need to be properly cleaned and disinfected before the equipment is put back into service or the area is reopened. Clean and disinfect surfaces and equipment as soon as possible after the incident occurs. Remember to wear appropriate personal protective equipment (PPE).

If blood or other potentially infectious materials have spilled onto the floor or another surface, take steps to prevent unnecessary people from accessing the area. If the spill contains sharp objects (e.g., shards of broken glass), do not pick them up with your hands. Instead, use tongs, a disposable scoop and scraper, or two pieces of cardboard to remove and dispose of them. Carefully wipe up the spill using absorbent towels, and/or use a solidifier (a fluid-absorbing powder).

After cleaning up the spill, flood the area with a freshly mixed disinfectant solution of approximately 1 part bleach to 9 parts water (a 10% solution). Always ensure that there is good ventilation, and wear gloves and eye protection when using bleach. Let the solution stand on the surface for at least 10 minutes, then use clean absorbent materials (such as paper towels) to wipe up the liquid and dry the area. Dispose of all materials used to clean up the spill in a labelled biohazard container. If a biohazard container is not available, place the soiled materials in a sealable plastic bag or a plastic container with a lid, seal the container, and dispose of it properly.

Soft materials such as clothes and linens that have been contaminated should be washed with soap and the hottest water available, at least 70°C (158°F). Hard-surfaced materials, such as dustpans, should be cleaned with a disinfectant solution. There is no need to dispose of soft materials or instruments as long as they are washed properly.



### **Handling an Exposure Incident**

An exposure incident occurs when one person's bodily fluids risk infecting another person through contact with the eyes, mucous membranes (e.g., in the mouth or nose), or an opening in the skin (e.g., a cut). Being pricked by a used syringe is also an example of an exposure incident.

In the event of an exposure incident, follow these steps immediately:

- Decontaminate the exposed area, if possible:
  - If your skin was exposed, wash the contaminated area with soap and water.
  - For infectious splashes into your mouth or nose, flush the area with water.
  - For infectious splashes into your eyes, irrigate the eyes with water, saline, or a sterile irrigant for 15 to 20 minutes.
- Report the exposure incident to EMS personnel or your healthcare provider.

If the exposure incident occurred in a workplace setting, notify your supervisor and follow your company's exposure control plan for reporting the incident and receiving post-exposure follow-up care. Some workplaces have additional requirements dictated by regulations and legislation: ensure that you are aware of any local, provincial/territorial, or organizational laws and regulations that apply to you.



# 3 Check, Call, Care

If you encounter an ill or injured person, there are three simple steps that you must take:



## Check

Once you recognize an emergency you must first check the scene, then check the person.

### Check the Scene

Before rushing to help an ill or injured person, stop and take a good look at the scene. This will form an initial impression. Try to answer these questions:

#### Is the scene safe for you, the ill or injured person, and any bystanders?

Enter a scene only if it is safe to do so. Look for hazards such as spilled chemicals, hostile bystanders, or oncoming traffic. Listen for things such as ringing alarms or leaking gas. Smell for things such as gas or smoke. Avoid entering confined areas with poor ventilation and places where there is a risk of explosion (e.g., from leaking propane or natural gas). Do not enter the scene if the person is hostile or threatening suicide.

If hazards are present, stay at a safe distance and call EMS/9-1-1. Leave dangerous situations for EMS personnel. Once EMS personnel arrive and make the scene safe, you can offer your assistance as appropriate. Remember that conditions at the scene may change, so a safe area could quickly become dangerous. It is important to continuously monitor your surroundings even if your initial assessment revealed a low level of risk.

#### What happened? How did it happen?

Take note of anything that might reveal the cause of the emergency. If the person is unresponsive and there are no witnesses, your check of the scene may offer the only clues as to what happened. Use your senses to detect anything out of the ordinary, such as broken glass, a spilled bottle of medication, or an unusual smell or sound. Keep in mind that

the ill or injured person may not be exactly where he or she was when the emergency occurred—someone may have moved the person, or the person may have moved in an attempt to get help.

### How many ill or injured people are there?

Look carefully for more than one ill or injured person. A person who is moving or making noise, or who has very visible injuries, will likely attract your attention right away, but there may be a person who is silent and not moving or a person obscured by debris or wreckage who you do not notice at first. It is also easy to overlook a small child or a baby.

### Is there someone to help?

Take note of bystanders who can be of assistance. A bystander who was there when the emergency occurred or who knows the person may be able to provide valuable information. Bystanders can also assist in other ways, such as by calling EMS/9-1-1, waiting for EMS personnel and leading them to the site of the emergency, getting needed items (such as an AED and first aid kit), controlling crowds, and reassuring the injured or ill person.

### What is your initial impression of the ill or injured person?

Before you reach the person, try to form an initial impression about the person's condition and what is wrong. For example, does the person seem alert, confused, or sleepy? Look at the person's skin—does it appear to be its normal colour, or does it seem pale, ashen (grey), or flushed? Is the person moving or motionless? Does the person have any immediately identifiable injuries? Look for signs of a life-threatening illness or injury, such as unresponsiveness, trouble breathing, or life-threatening bleeding.



## Check the Person (Primary Assessment)

If it is safe to do so, quickly check the person to determine his or her initial condition and whether there are any life-threatening concerns. This is called the primary assessment.

1. Check the person to see if he or she is responsive:
  - Does the person respond when you talk to him or her (e.g., does the person open his or her eyes, move, moan, or talk to you)?
  - Does the person respond when tapped on the shoulder?A person who does not respond at all is considered to be unresponsive.
2. Check the person's ABCs: Airway, Breathing, and Circulation




Checking the ABCs is a rapid assessment and should be done as quickly as possible.

## A = Check the Airway

Your first job is to make sure the person has an open airway. The airway is the pathway that connects the mouth and nose to the lungs. If it is closed or blocked, air cannot get in. The airway is most commonly blocked by the tongue. If the person is speaking, moaning, or crying, the person's airway is open.

If the person is unresponsive, perform a head-tilt/chin-lift. The head-tilt moves the back of the tongue away from the airway, and the chin-lift opens the epiglottis.



### Performing the Head-Tilt/Chin-Lift

To perform a head-tilt/chin-lift, place one hand on the person's forehead and 2 or 3 fingers under the chin. Gently tilt the head back until the chin is pointing upward.

## B = Check Breathing

Next, check for breathing. Someone who can speak or cry is breathing. Check for normal breathing for 5 to 10 seconds. To assess breathing, put your cheek close to the person's face so that you can hear and feel air coming out of the person's nose and mouth while you watch the chest rising and falling. A person is breathing normally if air is moving in and out of the lungs and the chest is rising and falling in a normal, regular pattern.

### Agonal Respiration

A person who is not breathing normally may be occasionally gasping for air: this is a reflex action called "agonal respiration" and unlike normal breathing, it is irregular and sporadic. Because agonal respiration does not oxygenate the blood, it is not considered normal breathing. A person who is experiencing agonal respiration is in cardiac arrest and requires immediate assistance.



Agonal respirations originate from lower brainstem neurons as higher centres become increasingly hypoxic (oxygen-deprived) during cardiac arrest. With agonal respiration, the diaphragm is still receiving intermittent residual impulses from the brain, resulting in sporadic gasping breaths. Agonal respiration is sometimes referred to as "air hunger" and it can appear as snorting, gurgling, moaning or gasping, a gaping mouth, or laboured breathing. The duration differs from person to person, lasting from a few minutes to hours.

## C = Check Circulation

If the person is breathing normally (more than an occasional gasp), his or her heart is beating. Checking circulation means quickly looking at the person from head to toe for signs of life-threatening external bleeding. Life-threatening bleeding must be controlled as soon as possible.



Unresponsiveness, trouble breathing, and life-threatening bleeding are all signs of a life-threatening emergency. If your initial check of the person reveals any of these conditions, you must provide care for that condition immediately. Make sure that someone calls EMS/9-1-1 right away, and obtain an AED and first aid kit if these items are available.

## Obvious Signs of Death

Obvious signs of death include:

- Torso transection (torso cut into two pieces)
- Decapitation (detached head)
- Decomposition

If you encounter a person with one or more of these signs, call EMS/9-1-1 and follow the dispatcher's instructions. Leave the body and scene exactly as they were found, as the area could be considered a crime scene.

## Rolling a Person from a Face-Down to a Face-Up Position

A person can collapse in a variety of positions. If a person is lying face down, is unresponsive, and not breathing, or if the face-down position makes it impossible to check the person's ABCs, then the person must be rolled into a face-up position without delay.

Move an ill or injured person only if:

- The person's position stops you from giving care for a life-threatening injury or illness.
- The person is blocking access to someone with a more serious injury or illness.
- The scene is becoming unsafe.

To roll a person from a face-down to a face-up position:

1. Support the head while rolling the person.
2. Try to roll the person as one unit (head, back, and legs at one time).
3. Open the airway with a head-tilt/chin-lift once the person is facing upward.
4. Check the person's ABCs.



## Call

If you identify an unresponsive individual or an individual with a life-threatening condition in your initial check, you must always activate EMS.

Whenever possible, use a mobile phone or ask a bystander to call EMS/9-1-1. Having someone else call is better than doing it yourself; this way, you can stay with the ill or injured person and continue to give first aid. If you are alone with the person and you do not have a mobile phone, call out loudly for help. If no one comes, get to a phone as quickly as you can and call EMS/9-1-1. As soon as you hang up, return to the person.

If someone else has to leave to make the call for you, ask that person to come back and tell you what the EMS/9-1-1 dispatcher said. If using a mobile phone, the person can stay with you while placing the call and speaking to the dispatcher.



Any time you activate EMS, you should also obtain a first aid kit and an AED if they are available.

## If You Are Alone: Call First or Care First?

If you are alone without a phone and there is no one to send to call EMS/9-1-1, you may need to decide whether to call first or give care first. Unless the situation specifically requires you to provide care before calling EMS/9-1-1, you should always activate EMS first so that help will arrive as soon as possible.

You should provide care first in the following situations:

**The person is choking or is experiencing anaphylaxis and carrying epinephrine.**


In these cases, the person could die before EMS arrives, so providing immediate care is your highest priority. Moreover, these situations can be corrected by First Aiders, so immediate care may actually improve the person's condition. If a person experiencing anaphylaxis is *not* carrying epinephrine, there is little that you can do as a First Aider, so you should call EMS/9-1-1 first before providing care.

**The person has life-threatening bleeding**

Life-threatening bleeding quickly reduces the body's blood volume, which can be fatal. Apply direct pressure immediately, then call EMS/9-1-1 as soon as possible. In many cases, the person will be able to apply pressure to his or her own injury while you (or a bystander) call EMS/9-1-1.

**The person is an unresponsive child or baby who is not breathing.**

Children's organs require high levels of oxygen in the blood to function efficiently and can rapidly shut down when oxygen levels are lowered. For this reason, cardiac arrest in children is more likely to be caused by low oxygen levels than by a problem with the heart itself. If you find a non-breathing child or baby who is unresponsive, you must introduce oxygen into his or her system as soon as possible. Immediately do 2 minutes of CPR (5 cycles of 30 compressions and 2 breaths) to increase the oxygen level, then quickly call EMS/9-1-1 before returning to provide more care.



**When You Call EMS/9-1-1**

When you call, the EMS dispatcher who answers will likely ask:

- Where is the emergency (e.g., the address, nearby intersections, or landmarks)?
- What is the nature of the emergency (i.e., is police, fire, or medical assistance needed)?
- What telephone number are you calling from?
- What is your name?
- What has happened?
- How many people are involved and what is their condition?

Don't hang up until the dispatcher tells you to. The dispatcher may need more information. Many dispatchers are also trained to give first aid and CPR instructions over the phone, which can be helpful if you are unsure of what to do or need to be reminded of the proper care steps.

**Radio Medical Advice in a Marine Environment**

In a marine environment, it may not be possible for EMS personnel to come to your assistance in an emergency. You may need to contact a doctor and request advice on providing care, even if it is outside of your training. Document the doctor's orders carefully.

Medical advice can be accessed by:

- Mobile phone.
- Radiotelephone contact with a shore radio station.
- Radiotelephone contact with a nearby port or ship with a doctor on board.

You should prepare the following information and present it to the doctor:

- Routine information about the ship
- Routine information about the person
- Details of the illness/injury
- Findings from your secondary assessment
- Care rendered and the person's response to that care

Transport Canada provides a reference regarding radiotelephone procedures (TP 9878E: Safety and Distress Radiotelephone Procedures), which should be displayed next to your vessel's radiotelephone. The procedures in this reference advise that safety and distress radio messages should be made on Channel 16 (156.8 MHz) or on frequency 2,182 kHz, MF.



### Care

Care for any life-threatening conditions first. Give the care that is needed, within the scope of your knowledge and training, and follow these general guidelines:

- Monitor the person's breathing and level of responsiveness.
- Help the person rest in the most comfortable position. If necessary, roll the person into the recovery position.
- Keep the person from getting chilled or overheated.
- Reassure the person by repeating that you are there to help and that EMS personnel have been called (if this is true).
- Continue to watch for changes in the person's condition.

### Recovery Position

A person who is unresponsive or has an altered level of responsiveness should not be left in a face-up position, as the airway can become blocked by the person's tongue or the person's saliva or other bodily fluids. You can help protect the airway by rolling the person onto his or her side and into the recovery position.



It is usually safest for any ill or injured person to be in the recovery position, so long as this doesn't interfere with providing care. However, the recovery position is unnecessary if the person is fully responsive and able to protect his or her own airway.

You should avoid rolling a person if doing so could worsen his or her condition (e.g., if the person's leg appears to be broken). As always, you should move the person only if it is safe to do so.

To roll a person safely into the recovery position:

1. Raise the person's arm that is closest to you.
2. Place the arm farther from you across the person's chest with the palm against the cheek.
3. Raise the knee of the leg farther from you.
4. Roll the person toward you as one unit by pulling the raised knee and supporting the head and neck with your other hand
5. Position the person on his or her side, and slide the bent knee into a position that prevents the person from rolling onto his or her face.
6. Move the person's other arm into a position of comfort in front of the body.
7. Reassess the person's ABCs.

The steps above are recommended, but there are other effective methods of moving a person into the recovery position. Regardless of how you do it, the important things to remember are as follows:

- Support and protect the head while rolling the person.
- Try to roll the person as one unit (head, back, and legs at the same time).
- Roll the person into a position where the body will stay safely on its side
- Position the head so that it keeps the airway open
- Check the ABCs after you complete the roll.

## ***When to Stop Giving Care***

Once you begin providing care to an injured or ill person, you must continue to give the appropriate care until:

- Another trained First Aider or EMS personnel takes over.
- You are too exhausted to continue.
- The scene becomes unsafe.

## ***When More Than One Person Is Ill or Injured***

If you are in a situation where there are several ill or injured people, the general principle is to provide care to the people who need it the most urgently. This involves deciding whose illness or injuries pose the greatest risks, as well as deciding who you are most able to help given your level of training. The process of sorting and providing care for multiple ill or injured people according to the severity of their conditions is called triage.

As an example, if one person has minor bleeding and another person has life-threatening bleeding, you should help the person with life-threatening bleeding first. EMS personnel will arrive before the minor bleeding becomes serious.

## ***Guidelines for Helping an Ill or Injured Person with Medication***

You should only help a person take his or her medication if:

- It is safe to do so.
- The person is responsive and has in some way expressed a need for help with finding, preparing, and/or taking the medication.

When helping someone with his or her medication, check the label to ensure that it is the correct medication and, if it is a prescription medication, that it was prescribed for the person who will be taking it. You should also carefully read over any instructions or warnings (e.g.,

how much the person should take and when it should be taken). You should read the label at least twice: once when you find the medication, and again before you give it to the person.

If there is no name on the label, confirm with the person that it is the medication he or she wants to take. You must also ask if the person has any allergies or is taking anything that could interfere with or react negatively with the medication. Quickly review these details with the ill or injured person and obtain his or her permission before helping.



Write down what medication was taken, how much of it was taken, and when it was taken. This information will be useful to EMS personnel.

## Finding the Medication

When helping a responsive ill or injured person find his or her medication, listen to instructions from the person on where to find it. If the person cannot speak, look in places where people typically keep their medication (e.g., backpack, purse, or pocket), or look for physical cues from the person (e.g., the person pointing at a bag). If the person cannot speak and you are helping with prescription medication, ask the person to confirm the name on the label by using physical signals (shaking or nodding the head, blinking once or twice, etc.). If the person is unable to do this, you may be able to confirm the person's identity by asking bystanders, looking for a piece of photo ID, or checking for a medical identification product.

## Preparing the Medication

Preparing the person's medication varies depending on the type of the medication and its instructions. For example, preparing oral medication may simply involve handing the person the correct number of tablets. In

the case of an inhaler or auto-injector, you may have to remove the cover and follow the preparation instructions provided on the package before handing it to the person for self-administration.

## Guiding the Person in Taking the Medication

Guiding the person in taking his or her medication also varies depending on the type of medication. For example, with tablets, this may involve telling the person to chew them or place them under the tongue. With an inhaler, this may involve instructing the person to shake the inhaler and exhale before dispensing the medication, or helping the person hold the inhaler to his or her mouth. When simply guiding the person, you do not press on the inhaler's plunger yourself. With an auto-injector, you may guide the person by showing where on the outer thigh the epinephrine should be injected.

## Giving Lifesaving Medication

Giving lifesaving medication involves helping an ill or injured person who has in some way expressed a need for help in taking his or her salbutamol (Ventolin) inhaler or using an epinephrine auto-injector. With an inhaler, you would dispense the correct number of sprays (from a metered-dose inhaler) either into a spacer or directly into the person's mouth as he or she inhales. With an auto-injector, you would inject the epinephrine directly into the person's outer thigh.

## Giving Ingested Medications

Helping with ingested medications should be limited to finding the medication, preparing the medication, and guiding the person in taking the medication. In cases where the person is physically unable to get the medication into his or her mouth (e.g., the person has injured his or her hands), First Aiders may assist by placing the correct number of pills into the mouth or holding a liquid dispenser up to the person's mouth, if the person has explicitly asked the First Aider to do so. Otherwise, you should never put anything into a person's mouth, as this can be a choking hazard.



## Secondary Assessment

After you have identified and cared for any life-threatening conditions found in the primary assessment, you must check the person for other injuries and conditions that may require care. This is called the secondary assessment.

If the person's ABCs seem normal, do a secondary assessment to look for injuries or conditions that were not identified in your primary assessment. The secondary assessment consists of three steps:

1. Asking questions
2. Checking the quality of the person's vital signs
3. Doing an injury check



If possible, record the findings from the secondary assessment or have someone else record them to help you remember. When EMS personnel arrive, tell them what you found.

## Ask Questions

Interview the ill or injured person (if he or she is responsive) and any bystanders at the scene to get more information. Asking the following SAMPLE questions can provide useful information about the person's situation:

- S** **SIGNS AND SYMPTOMS** — Do you have any cuts or bruises? How do you feel? Do you feel any pain? Does anything feel different?
- A** **ALLERGIES** — Are you allergic to anything?
- M** **MEDICATIONS** — Do you take any medicine? What is it for?
- P** **PAST MEDICAL HISTORY** — Do you have any medical conditions such as heart disease or other illnesses? Has this happened before?
- L** **LAST ORAL INTAKE** — When did you last eat or drink? What did you have?
- E** **EVENTS LEADING UP TO THE EMERGENCY** — What happened?

## Check the Vital Signs

Check the quality of the person's vital signs by evaluating his or her level of responsiveness, breathing, and skin.

**Level of Responsiveness:** Is the person alert or sleepy? Does the person seem confused? Is the person's awareness increasing, decreasing, or staying the same?

**Breathing:** Listen for sounds. Is the breathing fast or slow? Effective or adequate? Shallow or deep? Is breathing painful for the person?

**Skin:** Is it dry or wet? Is it an unusual colour or temperature?

A person's respiratory rate directly impacts the amount of oxygen that enters his or her bloodstream. Slower breathing brings less air into the lungs in a given period, and therefore decreases the amount of oxygen that crosses into the bloodstream. Breathing that is either too fast or too slow can also throw off the balance of oxygen and carbon dioxide in a person's blood. In a first aid emergency, a person's breathing rate can be affected by injury, blood loss, shock, pain, and anxiety. Providing continual care can help restore the person's breathing rate to normal.

An ill or injured person's skin may lose its underlying red tones, becoming more grey or white than usual. In people with darker skin, these changes may be harder to identify: skin may appear ashen-grey, yellow-brown or grayish-green. Changes may be most easily visible on the inside of the lips, the nail beds, or the skin around the mouth, which may be paler or have a dark blue tone due to the blueish tint of deoxygenated blood. The important thing is to note any changes from the person's normal skin tone, if possible.

### Injury Check

The goal of an injury check is to look carefully for injuries that were not identified during the primary assessment. An injury check may involve a focused examination or a hands-on check, depending on the comfort level of the person and whether the person is responsive.

### Focused Examination

If the person is responsive and able to answer questions, do a focused examination. Keep watching the person's vital signs. If the person's condition deteriorates, stop the examination and give first aid immediately.

To do a focused examination:

1. Explain that the purpose of the examination is to identify injuries.
2. Ask the person if anything hurts or feels uncomfortable.

3. If the person indicates an area of pain, discomfort, or concern, look at the area for signs of injury including discolouration (bruises) or deformities (odd shapes). For privacy reasons, do not remove any of the person's clothing unless it makes providing first aid difficult.
4. If there are no signs of any injury, ask focused questions about how the person feels. The symptoms the person describes might help to determine whether EMS/9-1-1 needs to be called and what kinds of care should be provided.
5. If you find a medical identification product, be sure to read it carefully. It may indicate what is wrong, who to call for help, and what care to give.
6. If the person doesn't complain of any pain or tenderness, and there are no signs of injuries, ask the person to rest for a few minutes in a comfortable position. Check the quality of the person's vital signs and ensure the ABCs are still unaffected. If there is no visible problem, help the person to stand up slowly when he or she is ready.
7. Based on your findings, decide whether you need to call EMS/9-1-1, and provide first aid care as needed.





### *Hands-On Check*

If a person is breathing but unresponsive, or is otherwise unable to communicate what is wrong, you may need to do a hands-on check to assess whether further first aid is required.

Conducting a hands-on check involves systematically checking the person from head to toe for signs of injury. Look and feel for any abnormalities such as bumps, soft spots, deformity, bruising, and bleeding. You should also look at the ground around the person for signs of blood or other body fluids. Continue to watch the person's ABCs and vital signs as you perform this check. If the person's condition deteriorates, stop the hands-on check and give first aid immediately.

! If the person is responsive, you will usually be able to get the information you need from a focused exam and sample questions alone, but there may be situations where touching the person is necessary to assess an injury. If so, you may need to do a portion of the hands-on check, focusing on the injured area.



Begin by checking the head, then work downward, focusing on the chest, abdomen, and legs before checking the arms. This prioritizes the areas that are more prone to life-threatening bleeding and organ damage.

When checking the chest, feel the ribs for signs of deformity. If the person is unresponsive, watch the rib cage as the person breathes. Both sides should expand at the same time. If the person is responsive, ask him or her to take a deep breath (if it doesn't cause any pain) to make it easier to check the expansion of the rib cage.

When checking the abdomen, press on it gently. It should be soft to the touch. If it feels hard or gentle pressure is painful, check carefully for bruising (some of the person's clothes may need to be moved or removed at this point).

Pushing on a person's injured pelvis can cause serious injury or worsen an existing condition, so this area should not be touched during the hands-on check.

! Be careful not to reach underneath someone during a hands-on check because there could be glass or other objects that could hurt you.

## Continual Care

A person's condition can change while you are waiting for EMS personnel to arrive. Do the following to provide continual care:

1. Have the person rest.
2. Help the person maintain a normal body temperature.
3. Monitor the person's condition and ABCs.
4. Move the person into the recovery position, if necessary.
5. Give comfort and reassurance to the person.

Helping the person stay calm has physiological benefits, such as decreasing the heart rate, reducing pain, slowing and deepening breathing, stabilizing blood pressure, and relaxing the body's muscles.

To help a person calm down and control his or her breathing:

1. Encourage the person to maintain eye contact with you.
2. Have the person breathe in through his or her nose while you raise your arm and count to five. Breathe with the person.
3. Have the person breathe out through his or her mouth, making an audible sound, while you slowly lower your arm and count to ten.
4. Repeat 3 to 5 times, providing encouragement in a calm voice.

## Should You Transport an Ill or Injured Person to the Hospital?

You may decide to take the ill or injured person to a hospital, urgent care clinic, doctor's office, or community first aid station if:

- The person is a friend or family member.
- The person's injuries are minor and non-life-threatening.
- There is no risk of the person's condition getting worse.
- There is no risk of the person experiencing an altered level of responsiveness.

If you decide it is safe to transport the person yourself, take someone else with you, if possible, to help keep the ill or injured person comfortable and to watch for any changes in his or her condition. This will let you focus on driving safely. Remember to obey traffic laws. No one will benefit if you are involved in a motor vehicle collision or get a speeding ticket on your way to the medical facility.

Never let an ill or injured person drive a motor vehicle for any reason. Driving is a high-risk activity that requires concentration, so getting behind the wheel while ill or injured is extremely dangerous. Even a person whose condition is mild could get worse while he or she is driving, and his or her ability to safely operate the vehicle could easily be compromised. Remember that it is not only the health and safety of the driver that is at stake, but that of everyone else on and near the road.



## Shock

A person who is ill or injured may go into shock. Shock happens when the vital organs do not get enough oxygen-rich blood. Shock is a life-threatening condition.

## Common Causes

Be on the lookout for shock when providing care for any injury or sudden illness, or when someone has been involved in a serious incident (even if he or she is not badly injured). Shock is often caused by significant fluid loss, for example, diarrhea and vomiting. This is especially true in children, who can become dehydrated more easily.

Other causes of shock include the following:

- Significant blood loss
- Heart damage
- Extensive burns
- Infection
- Anaphylaxis
- Severe infection (e.g., septicemia)

### **What to Look For**

The following are signs and symptoms of shock:

- Anxiety
- Cool, clammy skin
- Skin that is paler than normal
- Weakness
- Confusion
- Excessive thirst
- Rapid breathing
- Drowsiness or loss of responsiveness
- Nausea and vomiting

### **What to Do**

The best thing you can do when a person is in shock is to call EMS/9-1-1. While you are waiting for EMS personnel to arrive, provide care by:

1. Caring for the cause of the shock.
2. Having the person rest.
3. Keeping the person warm.
4. Monitoring the person's ABCs.
5. Providing comfort and reassurance.

### **Providing First Aid for Someone with a Disability**

It is not uncommon to feel intimidated when providing first aid to a person who has different abilities, because there is often uncertainty about how to care for the person. Remember that people are more similar than they are different, both physiologically and psychologically, and most aspects of first aid care do not vary from person to person.

Remember that a person's particular needs may not be visible, and that the person (or the person's direct caregiver) is the best source of information about his or her unique situation. If you are unsure, ask what you can do to help. The person is likely to appreciate your consideration.

### **General Tips for Providing First Aid to a Person with a Disability**

Remember the following tips when providing care for a person with any type of disability:

- As with all first aid, you need permission to help and should explain what you intend to do. These basic principles do not change because the person has different abilities.
- Avoid stereotypes and make no assumptions about what abilities the person has or does not have. Remember that not all disabilities are visible.
- Be confident and reassuring.
- If you cannot understand what the person is saying, politely ask him or her to repeat it.
- Don't touch or speak to service animals. They are working and have to pay attention at all times.
- Get permission before touching assistive devices, including wheelchairs.

## *Providing First Aid for Someone with Vision Loss*

Examples of vision loss include tunnel vision (where a person cannot see objects to the side), lack of central vision, and total blindness, in which the person receives no visual information from the world around them. Most people with vision loss still have some degree of sight.

Keep the following in mind when providing first aid to an individual with vision loss:

- If you need to leave the person, let him or her know that you are leaving and will be back.
- Narrate your actions so that the person can follow what you are doing.
- If you're unsure about how to provide directions, simply ask the person how he or she would like you to do so.

## *Providing First Aid for Someone with Hearing Loss*

Keep the following points in mind when providing first aid to an individual with any type of hearing loss:

- Get the person's attention before speaking. The best way to do this is usually by lightly touching the person's shoulder or gently waving your hand.
- Ask the person how you can help. Don't shout.
- If communication is difficult, ask if the person prefers to communicate in another way, such as using a pen and paper.
- Be patient if you are using a pen and paper to communicate. American Sign Language may be the person's first language and it has its own grammatical rules and sentence structure.
- If the person uses a hearing aid and you are having trouble communicating, minimize background noise or move to a quieter area, if possible.

## *Providing First Aid for Someone Who Is Deafblind*

"Deafblind" describes people who have some degree of both vision and hearing loss. Many people who are deafblind are accompanied by an intervenor (a professional who helps with communication). If an intervenor is accompanying the person, tell him or her who you are, but then speak directly to the ill or injured person. Don't touch the person abruptly and don't touch the person without permission.

## *Providing First Aid for Someone with a Physical Disability*

There are many different types of physical disability, and not all of them are visible. The important thing is not to diagnose the person: what matters is determining whether the conditions you encounter are pre-existing or whether they are signs of the injury or illness you are providing care for.

Keep the following tips in mind when providing first aid to an individual with a physical disability:

- Speak naturally to the person (physical and mental disabilities are not the same thing).
- Adjust your position if necessary so that you can make eye contact with the person while providing care.

### *Providing First Aid for Someone with an Intellectual or a Developmental Disability*

It may not be apparent that a person has an intellectual or developmental disability unless this fact is communicated to you.

Keep the following points in mind when providing first aid to an individual with an intellectual or a developmental disability:

- As much as possible, interact with the person as you would with anyone else in the same situation.
- Use straightforward language when communicating.
- Give one piece of information at a time.

### *Providing First Aid for Someone with a Speech or Language Impairment*


Speech and language impairments may affect a person's ability to communicate verbally, with written language or both. Assistive devices such as communication boards are often used by individuals with severe speech or language impairments.

Keep the following tips in mind when providing first aid to an individual with a speech or language impairment:

- If possible, ask questions that can be answered with "Yes" or "No."
- Give the person time to communicate and answer your questions.
- Wait for the person to finish speaking and do not try to finish his or her sentences.

# 4 Choking





The airway is the passage that connects the nose and mouth with the lungs. Choking occurs when the airway becomes partially or completely blocked by a foreign object (e.g., a piece of food or a small toy), by swelling in the mouth or throat, or by fluids, such as vomit or blood. If the airway is blocked by the person's tongue or by swelling, this is called an anatomical obstruction. If it is blocked by a physical object, this is called a mechanical obstruction.

Children younger than 5 years old have a particularly high risk of choking because their airways are about the size of their little fingers, but a person of any age can choke.

## Anatomy and Physiology

The respiratory system consists of the airway and the lungs. When breathing in, air moves from the outside world into the lungs through the airway. The respiratory system constantly supplies the body with the oxygen it needs and removes carbon dioxide. The process is largely involuntary and is controlled by the brain.

## Partial Choking

Partial choking happens when the airway is partially blocked. Coughing is the body's way of clearing the airway, and so it may indicate a partial airway obstruction. A person who is coughing is still able to breathe. Since forceful coughing usually eliminates the obstruction, encourage the person to keep coughing. Stand by and monitor the person in case further help becomes necessary. Note that the care for complete choking will be ineffective for partial choking, because it depends on creating

pressure behind the blockage (which is impossible unless the blockage is complete).

If the person is or becomes too weak to cough, his or her condition will quickly deteriorate into complete choking. If a choking person is unable to cough forcefully for any reason, call EMS/9-1-1 immediately and monitor the person's condition closely.

## Complete Choking

Complete choking happens when the airway is completely blocked. When a person is experiencing complete choking, he or she is unable to breathe and is in a life-threatening situation. Immediate first aid (and possibly medical intervention) is required to remove whatever is blocking the airway.

## Common Causes

Any medical condition that affects a person's ability to chew and/or swallow increases his or her risk of choking. So can dental problems or poorly fitting dentures that affect a person's ability to chew food properly.

Common causes of choking include the following:

- Trying to swallow large pieces of food
- Eating while talking, laughing, walking, or running
- Eating too quickly

# Choking Hazards

## FOODS

- Nuts and seeds
- Hot dogs and sausages
- Chunks of meat or cheese
- Chunks of fruit (e.g., apples)
- Small fruits (e.g., whole grapes and cherries)
- Hard raw vegetables (e.g., carrots and celery)
- Popcorn
- Peanut butter
- Hard, gooey, or sticky candy (e.g., peppermints, marshmallows, and gummy bears)
- Foods that break easily into small pieces (e.g., teething biscuits and cookies)

## HOUSEHOLD OBJECTS AND TOYS

- Plastic bags, broken or uninflated balloons (the thin material can block the airway)
- Coins
- Buttons
- Small "button" batteries (e.g., those found inside watches, car key fobs, and hearing aids)
- Magnets
- Marbles
- Small rocks, beads, or decorative stones
- Pen and marker caps
- Jewellery
- Pills and vitamins
- Items from the garbage or recycling (e.g., corners of milk bags and pull tabs from cans)
- Toys meant for older children, which may be small or have small parts

## Prevention

The following may reduce the risk of choking in adults:

- Chewing food well before swallowing
- Eating slowly and calmly
- Avoiding talking, laughing, walking, and other kinds of physical activity while chewing

The following may reduce the risk of choking in children and babies:

- When babies start eating solid food, begin with purées as opposed to solid pieces.
- Always supervise children and babies when they are eating.
- Teach children to eat calmly, chew properly, and not to speak with a mouthful of food.
- Encourage children to sit while eating.
- Make sure that babies and young children cannot reach objects small enough for them to swallow.
- Remember that children can choke on soft plastic (e.g., the cut-off corners of milk bags) and rubber (e.g., burst balloons) as well as hard objects, so keep these items out of reach.



Toys smaller than 4 cm (1.5 in.) in diameter are choking hazards for babies and toddlers. Use the following guideline: if an object can fit through a toilet paper tube, it is not safe for children younger than 4 years old.



## Partial Choking

### What To Look For



Typical signs of partial choking include the following:

- A look of panic, with wide eyes
- Forceful or weak coughing
- One or both hands clutching the throat

### What To Do

#### Call

It is not usually necessary to call for help for partial choking, though you may need to do so if the obstruction does not clear or the person's condition worsens.

#### Care

1. If the person is coughing or can speak, encourage him or her to cough forcefully, and do not interfere. Forceful coughing may be enough to clear the obstruction on its own.
2. Encourage or assist the person to sit upright, if possible, as this will make coughing easier and more effective.
3. Continue to monitor the person's condition until either the obstruction clears itself or the person's condition worsens.

## Complete Choking

### What To Look For



The signs and symptoms of complete choking include the following:

- An inability to speak, cough, or breathe
- High-pitched squeaking noises
- Flushed (red) skin that becomes bluish or paler than normal
- A look of panic with wide eyes
- One or both hands clutching the throat

### What To Do

#### Responsive Adult or Child Over 12 Months



##### Call

- If possible, send someone to call EMS/9-1-1 and get an AED while you care for the person.
- If you are alone, immediately begin providing care for complete choking. Shout for help to try to attract the attention of a bystander, but do not delay care by calling EMS/9-1-1 yourself.



##### Care

1. Try to dislodge the object by combining any two of the following three options: back blows, abdominal thrusts, and chest thrusts. Continue alternating between the two methods until the object comes out, the person begins to breathe, or the person becomes unresponsive.

You may need to adapt which combination of the methods you use depending on the specific circumstances. For example, if the person is pregnant, or you cannot fully reach around the person, you will not be able to provide abdominal thrusts and will therefore have to do a combination of back blows and chest thrusts.


### Back Blows

1. Stand behind the person (or kneel, if you are significantly taller) and wrap one arm diagonally across the person's chest.
2. Bend the person forward at the waist until the upper airway is at least parallel to the ground.
3. With the heel of your free hand, deliver up to 5 firm back blows between the shoulder blades.



### Abdominal Thrusts

1. Stand behind the person and ensure that your stance is stable. If the person is much shorter than you are, it may be more effective to kneel.
2. Wrap your arms around the person's waist, make a fist, and place it just above the belly button.
3. Cover your fist with your other hand and give up to 5 quick, inward and upward thrusts into the person's abdomen.



### Chest Thrusts

1. Stand behind the person (or kneel, if you are significantly taller) and wrap both of your arms around the person's chest, just under the armpits.
2. Make a fist and place it in the middle of the person's chest, with your thumb facing inward, and place your other hand over your fist.
3. Give up to 5 chest thrusts by pulling straight back toward you. If you find your thrusts are not effective, pull more sharply and deeply.



## Special Choking Circumstances

### Larger or Pregnant Responsive, Choking Adult



1. If you cannot reach far enough around a person to perform abdominal thrusts, or if you are aiding a woman who is obviously pregnant, alternate between 5 firm back blows and 5 chest thrusts.
2. Continue until the object comes out, the person begins to breathe or cough, or the person becomes unresponsive.

### If You Are By Yourself and Choking



1. Dial EMS/9-1-1 and leave the phone off the hook. This will tell the dispatcher to send help.
2. If there are people nearby, move to a place where you can be noticed.
3. Attempt to dislodge the object by performing an abdominal thrust against a safe object with no sharp edges or corners, such as the back of a chair or the edge of a table.

### Responsive, Choking Toddler

1. If the child is short enough, kneel on the floor with one knee raised.
2. Perform 5 back blows with the child leaning across your raised leg, alternating with either 5 chest thrusts or 5 abdominal thrusts.
3. Continue until the object comes out, the child begins to breathe or cough, or the child becomes unresponsive.



Calling EMS/9-1-1 does not always tell the dispatcher your location. Enhanced 9-1-1 services may be able to triangulate your position from a mobile phone call, but it is accurate only to within several hundred metres. If you are using a VOIP system, the location of the call is dependent upon how the system is set up. Additionally, if you call from a landline phone, your exact position within the building will not be pinpointed. While the phone call is important, it is equally important that you move to a public area or a place where you are likely to be seen.

## Choking Baby (Less Than 12 Months)



### Call

- Shout for help.
- If the baby is making high-pitched noises, is wheezing, can no longer make a sound, or becomes too weak to cough, have someone call EMS/9-1-1 and get an AED while you care for the baby.
- If you are alone, immediately begin care for complete choking. Shout for help to try to attract the attention of a bystander.



### Care

1. Sit or kneel holding the baby.
2. Position the baby face down along your forearm, holding his or her jaw in your hand.
3. Rest your forearm on your leg so that the baby's head is lower than his or her body.
4. Deliver 5 firm back blows between the shoulder blades with the heel of your free hand.



5. If the object has not been dislodged, turn the baby face-up, ensuring you support the head.



6. Place 2 fingers on the middle of the chest just below the nipple line and quickly deliver 5 firm chest compressions, pushing down 1/3 of the chest's depth.
7. Repeat the 5 firm back blows and 5 chest compressions until the object is coughed up, the baby starts to cry, breathe, or cough, or the baby becomes unresponsive.



## After the Object Comes Out

An object lodged in the airway can cause damage to the sensitive tissues in the trachea and the back of the throat. This can cause bleeding and swelling.

Any person who has suffered complete choking should always seek medical attention, even if the object comes out. If you have done back blows, abdominal thrusts, or chest thrusts, those procedures may have caused damage (in addition to the damage caused by the choking itself), so any person given this care should be assessed in a hospital.

A person who has experienced partial choking should seek medical attention if he or she experiences any kind of respiratory distress after the incident. The person should be monitored for several hours after the incident to ensure that there is no damage: if there is no sign of injury in the first few hours, the person is probably fine.


## ***Unresponsive Choking***

If the choking person becomes unresponsive, lower him or her to the ground as safely as possible and immediately begin CPR, starting with chest compressions.



# 5 Circulation Emergencies

Circulation emergencies are sudden illnesses or injuries involving the heart or the blood vessels. Because every minute counts when a person is experiencing a circulation emergency, the person's survival often depends on First Aiders acting quickly and giving appropriate care until EMS personnel arrive and take over.



## Cardiovascular Disease

Cardiovascular disease is an abnormal condition that affects the heart and blood vessels. It is one of the leading causes of death in Canadians, and it is the underlying cause of many circulation emergencies.

Although a circulation emergency such as angina, a heart attack, or a stroke can occur suddenly, the underlying cardiovascular disease often develops over a long period of time. In fact, it can begin as early as the teenage years.

Cardiovascular disease occurs when the arteries become hardened and narrowed—a process called atherosclerosis. This damage occurs gradually, as cholesterol and fatty deposits called plaque build up on the inner artery walls. As this buildup worsens, the arteries become narrower, reducing the amount of blood that can flow through them. The narrowing of arteries increases the risk of artery blockages, which can cause angina, heart attacks, transient ischemic attacks (TIAs), and strokes.

## Prevention

Cardiovascular disease has both controllable and uncontrollable factors. Uncontrollable factors include age, sex, family history, and ethnicity.

Fortunately, many risk factors of cardiovascular disease are the result of lifestyle choices and are therefore within a person's control.

### Smoking

Smoking is a major risk factor for developing cardiovascular disease, but it is also one of the most controllable factors. Smokers have a risk of sudden death that is 2 to 4 times greater than that of non-smokers. Second-hand and smokeless tobacco also increase the risk of cardiovascular disease.

Fortunately, the risk starts to drop as soon as a person stops smoking. Compared to a smoker, the risk of cardiovascular disease decreases by half within 1 year of quitting smoking. Within 15 years, the risk is close to that of a non-smoker. Many effective community programs are available for individuals who wish to quit smoking permanently.

### Cholesterol

Cholesterol is produced naturally in the body, and is necessary for cellular function. However, not all cholesterol is good for you: low density lipoproteins (LDL, or "bad cholesterol") can build up in the bloodstream and stick to the blood vessel walls, narrowing the blood vessels and reducing blood flow.

It is easy to measure cholesterol with a blood test. Cholesterol can be controlled with a proper diet and exercise, along with medication if necessary.

## Blood Pressure

High blood pressure can stress and weaken blood vessels throughout the body, and it can contribute to the narrowing of blood vessels. High blood pressure can be hard to detect, so testing for it should be a routine examination. High blood pressure can be controlled with exercise, diet control, stress reduction, and medication.


## Weight

Obesity directly contributes to heart disease. Middle-aged people who are significantly overweight have 3 times the risk of a fatal heart attack when compared to people who are at a healthy weight. Obesity also increases the risk of high blood pressure and blood cholesterol levels. Maintaining a healthy weight requires exercise and a balanced diet.

## Stress

Stress is a normal part of life. Learning to cope with it well helps you to stay healthy and reduces your risk of developing cardiovascular disease, among other illnesses.

Stress can be reduced by developing rewarding hobbies and exercising regularly. It can also be helpful to set realistic goals, practice relaxation exercises, and avoid caffeine, which can reduce your ability to handle stress.



## Heart Attack

A heart attack occurs when the heart muscle cannot get enough oxygen because of a blockage in one of the arteries that feeds it.

## What to Look For



The signs and symptoms of a heart attack vary from person to person, and can be different in women and men. Even a person who has had a heart attack before may not experience the same signs and symptoms if he or she has a second heart attack.

A person who is having a heart attack may experience chest pain, which can range from mild to unbearable. The person may complain of pressure, squeezing, tightness, aching, or heaviness in the chest. The pain or discomfort is persistent, lasting longer than 3 to 5 minutes. It is not relieved by resting, changing position, or taking medication. It may be difficult to distinguish the pain of a heart attack from the pain of indigestion, heartburn, or a muscle spasm.

Other signs and symptoms of a heart attack include the following:

- Discomfort or pain that spreads to one or both arms, the jaw, the shoulder, the neck, or (more commonly in women) the back or the upper part of the stomach
- Problems breathing (e.g., noisy breathing, shortness of breath, or breathing that is faster than normal)
- Cold, sweaty skin
- Skin, lips, and fingers that are bluish, ashen (grey), or paler than normal
- Feelings of anxiety, denial, or impending doom

## "Soft Signs" of a Heart Attack

During a heart attack, many females, elderly people, and people with diabetes tend to experience "soft signs," which are milder or more generalized than the "classic" signs and symptoms of a heart attack. These "soft signs" may be experienced for hours, days, or even weeks prior to the heart attack and are often dismissed as nothing out of the ordinary.

"Soft signs" include:

- Mild, unfocused chest discomfort that:
  - Comes and goes
  - Does not necessarily feel painful
  - Gets better with rest and worse with activity, or gets progressively worse
- Extreme fatigue (tiredness)
- Gastric discomfort, nausea, or vomiting
- Flu-like symptoms
- Dizziness or light-headedness



Not everyone experiences chest pain during a heart attack.

## What to Do



### Call

If you suspect a person is having a heart attack, call EMS/9-1-1 and get an AED immediately.



### Care

1. Have the person rest comfortably.
2. If acetylsalicylic acid (ASA) is available, encourage the person to chew 160 to 325 mg (either 1 adult-dose or 2 low-dose ASA tablets), unless the person has an allergy or a contraindication to ASA, such as a bleeding disorder. Ensure that the person thoroughly chews and swallows the ASA. Do not repeat the dosage.
3. If the person takes a prescribed medication to relieve chest pain (e.g., nitroglycerin), offer to locate the medication and help the person to take it.
  - If the person has nitroglycerin in a spray or pill form, ensure that the person sprays or places the nitroglycerin under his or her tongue.
4. Reassure the person. Anxiety may increase the person's discomfort.





Other painkilling medications such as acetaminophen (e.g., Tylenol®) or ibuprofen (e.g., Advil®) do not have the same effect as ASA in reducing damage due to heart attacks. Do not substitute ASA with acetaminophen or ibuprofen.

If given in conjunction with nitroglycerin, erectile dysfunction drugs can cause blood pressure to lower to a life-threatening level. If the person has recently taken an erectile dysfunction drug (in the previous 24 to 48 hours), advise the person not to take nitroglycerin.

## Angina

Angina occurs when the heart requires more oxygen than it is receiving. This occurs when two factors are combined: 1) the arteries are narrowed by cardiovascular disease, reducing blood flow; and 2) the heart's oxygen needs are elevated (e.g., during times of physical activity or emotional stress). Angina causes a painful squeezing, suffocating, or burning feeling in the chest.

Uncontrolled high blood pressure, anemia, and certain heart disorders can all contribute to a person's risk of angina.


The signs and symptoms of angina are similar to the signs and symptoms of a heart attack, but the pain is usually triggered by physical activity or stress, goes away if the person rests, and is often relieved by medication (e.g., nitroglycerin). Because it is not possible for First Aiders to distinguish between angina and a heart attack, you should treat a person with angina as you would treat a person having a heart attack.

## Myth-Information

**Myth:** You should delay calling EMS/9-1-1 for a person if you suspect that he or she is experiencing angina and not a full heart attack.

You should always call EMS/9-1-1 if you have any reason to suspect that a person might be having a heart attack. It is better to have EMS personnel arrive at a scene and not be needed than to have a person who is having a heart attack not receive medical attention.

Many people who are having a heart attack delay seeking care because they hope they are experiencing signs and symptoms of a more minor condition that will go away with time, such as indigestion, heartburn, a muscle strain, or the flu. People often worry about calling EMS/9-1-1 for a "false alarm." Without immediate medical intervention, heart attacks are often fatal, so the risk of not seeking care outweighs any inconvenience.



## Stroke

A stroke happens when the blood flow to part of the brain is interrupted, resulting in the death of brain cells. Strokes can cause permanent brain damage, but with quick action, the damage can sometimes be stopped or reversed. Although strokes are most common in older adults, a person of any age, even a child, can have a stroke.

## Causes of Stroke

### Ruptured Aneurysm

A hemorrhagic stroke occurs when a brain aneurysm ruptures. A brain aneurysm is a swollen, fragile section in the wall of an artery that carries blood to the brain. When it ruptures, it causes bleeding in the surrounding brain tissues, killing the nearby brain cells and causing a stroke.

As an intact brain aneurysm usually has no signs or symptoms, it can go undetected. However, an unruptured brain aneurysm may press against areas in the brain and, depending on the severity of the aneurysm and its location, it can cause signs and symptoms such as neck pain, acute headaches, blurred vision, and changes in speech.

### Blood Clot

An ischemic stroke occurs when a clot becomes lodged in an artery of the brain, usually because the artery has been narrowed by cardiovascular disease. The blockage prevents blood from reaching the brain cells it usually supplies.

### Brain Tumour

A stroke can also occur when a tumour in the brain presses on an artery. The tumour has a similar effect to a blood clot, resulting in decreased blood flow or a complete blockage of blood flow to part of the brain.

### Prevention

Because the risk factors for stroke are similar to those for heart disease, the risk of a stroke can be reduced by following the same lifestyle changes discussed for preventing cardiovascular disease.

## What to Look For



The signs and symptoms of strokes can vary from person to person. A person who is having a stroke may suddenly develop one or more of the following signs and symptoms:

- A sudden, severe headache
- Dizziness or confusion
- Unresponsiveness or temporary loss of responsiveness
- Sudden loss of bladder or bowel control

### FAST

When trying to determine if a person is having a stroke, remember the acronym FAST:

- F** **FACE** — Facial numbness or weakness, especially on one side.
- A** **ARM** — Arm numbness or weakness, especially on one side.
- S** **SPEECH** — Abnormal speech, difficulty speaking or understanding others, or a loss of speech.
- T** **TIME** — Time is important; call EMS/9-1-1 immediately.

## What to Do



### Call

Call EMS/9-1-1 and get an AED.



### Care

1. Have the person rest in a comfortable position. If the person prefers to lie down, or is unresponsive, drooling, or having trouble swallowing, place the person in the recovery position.



2. Note when the signs and symptoms first started (or, if you do not know when the signs and symptoms started, note the last time the person was known to be well).
3. Monitor the person's condition and provide reassurance until EMS personnel arrive; a stroke can be an extremely frightening experience.



Some of the medications used to treat strokes can only be administered within a short window of time after the onset of signs and symptoms. It is crucial to note when the person first began to experience the signs and symptoms, or the last time the person was known to be well.

## Transient Ischemic Attack (TIA)

A transient ischemic attack (TIA) is also referred to as a "mini-stroke" and is caused by a temporary drop in blood flow to part of the brain. A TIA may be caused by a tumour or a clot in an artery in the brain. The signs and symptoms of a TIA are the same as the signs and symptoms of a stroke, but they disappear within a few minutes or hours.

A person who has had a TIA is at a very high risk of having a stroke in the near future. In fact, more than 10% of people who have a TIA will have a stroke within 3 months, with half of these strokes happening within 48 hours of the TIA.

### Myth-Information

**Myth:** A person with a suspected stroke does not need medical attention if his or her signs and symptoms dissipate.

TIA signs and symptoms will disappear within a few hours, but the person still needs to be checked by a medical professional. TIAs are warning signs and are often quickly followed by strokes. Always call EMS/9-1-1 if the person has demonstrated signs and symptoms of a TIA or stroke, even if the signs and symptoms have lessened in severity or disappeared completely.

## Life-Threatening Bleeding

Life-threatening bleeding refers to a situation where large amounts of blood loss occur either externally (outside the body) or internally (within the body). Life-threatening bleeding can occur whenever one or more of the body's blood vessels are broken. It must be controlled immediately.

Bleeding from arteries (arterial bleeding) is often rapid and profuse, and it is always life-threatening. This is because arterial blood is under direct pressure from the heart, so it usually spurts from the wound, making it difficult for clots to form. As a result, arterial bleeding is harder to control than bleeding from veins and capillaries.

Veins are damaged more often than arteries because they are closer to the skin's surface. Bleeding from veins (venous bleeding) is easier to control than arterial bleeding. Venous blood is under less pressure than arterial blood and flows from the wound at a steady rate without spurting. Only damage to veins deep in the body, such as those in the torso or thigh, produces profuse bleeding that is hard to control.

## Prevention

The following precautions may reduce the likelihood of an injury that causes life-threatening bleeding:

- Be familiar with your surroundings.
- Keep all sharp objects, such as knives, in a safe place.
- Get proper training on machinery in the workplace.
- Always follow the outlined safety procedures when operating equipment.
- Wear and use appropriate safety equipment at work and at home.
- Only use equipment for its intended purpose.
- Stay alert when operating equipment.

## Life-Threatening External Bleeding

### What to Look For



The signs and symptoms of life-threatening external bleeding include:

- Blood spurting or flowing freely from a wound
- Blood that fails to clot after you have taken all measures to control bleeding
- Large amounts of blood loss

## What to Do

### Call

Call EMS/9-1-1 and get an AED.

### Care

1. Expose the wound.
2. Apply firm and direct pressure to the wound.



3. While maintaining direct pressure, apply a dressing and bandage, and then secure them in place.



4. Reassess the wound to see if the pressure has stopped the bleeding. If not, consider using a tourniquet if the wound is on a limb.





## Using a Tourniquet

A tourniquet is a tight band placed around an extremity (e.g., an arm or leg). It constricts blood vessels to stop life-threatening bleeding in circumstances where normal bleeding control is impossible or ineffective.

Tourniquets are commercially made, but they can also be improvised with any wide strip of material (e.g., a triangular bandage). If you find yourself in a situation where you need to apply a tourniquet, a commercially manufactured tourniquet is preferred over an improvised device. Follow the manufacturer's instructions for applying the tourniquet. Although tourniquets may have slightly different designs, all are applied in generally the same way. Once a tourniquet is in place, you should not remove it for any reason.

## When to Apply a Tourniquet

Using a tourniquet involves a level of risk, which must be balanced against the potential benefits. You should only apply a tourniquet in the following situations:

- The person has life-threatening external bleeding that cannot be controlled using direct pressure.
- The person is in a physical location that makes it impossible to apply direct pressure (e.g., the person or the person's injured limb is trapped in a confined space).
- You must move the person and are unable to maintain direct pressure while doing so.

## Applying a Tourniquet

1. Apply the tourniquet 5 to 10 cm (2 to 4 in.) above the injury. If there is a joint in this range, apply the tourniquet above it, at least 2.5 cm (1 in.) away from the joint.



2. Tighten the tourniquet until the bleeding stops.



3. Secure the tourniquet in place.



4. Document the time the tourniquet was tightened.





## Using Hemostatic Dressings

A hemostatic dressing is a dressing treated with a substance that speeds clot formation. As is the case with tourniquets, hemostatic dressings are used in cases of life-threatening bleeding where standard first aid procedures fail or are not practical. Typically, hemostatic dressings are used on parts of the body where a tourniquet cannot be applied, such as the neck or torso. A hemostatic dressing can also be used to control bleeding from an open wound on an arm or a leg if a tourniquet is ineffective. The hemostatic dressing is applied at the site of the bleeding (possibly inside the wound) and is used along with direct pressure.

## Life-Threatening Internal Bleeding

Internal bleeding is the escape of blood from arteries, veins, or capillaries into spaces in the body. Life-threatening internal bleeding usually occurs in injuries caused by a violent blunt force, such as when someone falls from a height. You should suspect internal bleeding after any injury that involved a forceful blow to the body. Internal bleeding may also occur when a sharp object, such as a knife, penetrates the skin and damages internal structures.

## What to Look For



Signs and symptoms of life-threatening internal bleeding include the following:

- Bruising in the injured area
- Soft tissues (e.g., the abdomen) that are tender, swollen, or hard
- Blood in saliva or vomit
- Pain
- Severe thirst, nausea, and vomiting
- Anxiety

## What to Do



### Call

Call EMS/9-1-1 and get an AED. You cannot provide first aid for life-threatening internal bleeding as a First Aider. Life-threatening internal bleeding, especially if it results from damage to an organ, requires in-hospital medical intervention. Help the injured person rest in the most comfortable position and provide continual care while waiting for EMS personnel.



A person with life-threatening internal bleeding may be very thirsty, but giving anything by mouth (even water) can cause serious complications.

# 6 CPR & AED






CPR, or cardiopulmonary resuscitation, is a skill that is used when a person is in cardiac arrest. CPR involves giving sets of 30 chest compressions followed by sets of 2 rescue breaths; it keeps oxygenated blood moving to the brain and other vital organs until advanced medical help arrives. Because we breathe in 21% oxygen and breathe out 16%, even exhaled breath has more than enough to sustain life.

### Cardiac Arrest

Cardiac arrest occurs when the heart stops beating or beats too ineffectively to circulate blood to the brain and other vital organs. Under normal circumstances, a network of special cells in the heart muscle conducts electrical impulses that coordinate contractions, causing the heart to beat rhythmically. In cardiac arrest, the electrical impulses become abnormal and chaotic. This causes the heart to lose the ability to beat rhythmically, or to stop beating altogether.



The respiratory and circulatory systems are very closely linked. After breathing stops, it is not long before the heart stops. This is especially true in children: pediatric cardiac arrests are often the result of airway or breathing emergencies.



### Sudden Cardiac Arrest

Cardiac arrest can happen suddenly and without any warning signs; this is called sudden cardiac arrest. People who have a history of cardiovascular disease or a congenital heart disorder are at higher risk for sudden cardiac arrest. However, sudden cardiac arrest can happen in people who appear healthy and have no known heart disease or other risk factors for the condition. A person who experiences sudden cardiac arrest is unlikely to survive without immediate care.



### Clinical and Biological Death

Clinical death occurs when breathing stops and the heart stops beating. A person without a heartbeat is clinically dead. Because permanent brain damage does not begin for 4 to 6 minutes, clinical death is reversible with immediate care and intervention.

Biological death occurs when the brain becomes irreversibly damaged, which occurs after about 8 to 10 minutes without oxygen. This is also called "brain death."

## Common Causes

Cardiovascular disease is the most common cause of cardiac arrest. Other common causes include the following:

- Drowning
- Severe blood loss
- Suffocation or complete choking
- Electrocution
- Drug overdose
- Severe chest injuries
- Commotio cordis
- Other heart diseases and abnormalities

Cardiovascular disease and certain congenital heart conditions (i.e., conditions that a person is born with) can increase the risk of cardiac arrest. Breathing emergencies can also lead to cardiac arrest. For example, choking or drowning can interrupt the body's supply of oxygen, causing the heart to stop beating. Every organ in the body needs a steady supply of oxygen in order to work properly, and the heart is no exception. Severe trauma, electrical shocks, and drug overdoses are other potential causes of cardiac arrest. Although cardiac arrest is more common in adults, it does occur in young people as well. The most common causes of cardiac arrest in children and babies are breathing emergencies, congenital heart disorders, and trauma.

Cardiac arrest is a life-threatening condition, so starting CPR is the top priority. A person in cardiac arrest may have other conditions as well, such as a possible head, neck, or spinal injury, but caring for other injuries is lower priority than starting CPR and getting an AED.

## What to Look For

When a person experiences cardiac arrest, signs and symptoms may include the following:

- The person suddenly collapses.
- The person is unresponsive and not breathing, or is unresponsive with agonal breaths.



For each minute that CPR and use of an AED are delayed, the person's chance of survival is reduced by about 10%.

## Cardiac Arrest, Adult (After the Onset of Puberty)

### What to Do



#### Call

Have someone call EMS/9-1-1 and get an AED. If you are alone with an adult, call EMS/9-1-1 yourself, get an AED, and then return to care for the person.



## Care

1. Place both of your hands on the centre of the person's chest.
2. Do 30 compressions:
  - Push down at least 5 cm (about 2 in.), pushing deeply and steadily.
  - Allow the chest to fully recoil after each compression (do not lean on the chest).
  - The chest compression rate should be between 100 and 120 beats per minute (30 compressions in 15 to 18 seconds).
3. Give 2 rescue breaths:
  - Open the airway by doing a head-tilt/chin-lift.
  - Place your barrier device over the person's mouth and nose.
  - If using a plastic face shield, seal your lips tightly over the person's mouth and pinch the nose. If using a face mask, ensure it is fitted tightly over the mouth and nose.



- Give 2 rescue breaths. Each breath should last 1 second, with just enough volume to make the chest start to rise.
4. If both breaths go in, repeat the cycle of 30 compressions and 2 breaths.



If there are two First Aiders present, they should alternate every 5 cycles (about every 2 minutes).

## Cardiac Arrest, Child (1 Year to the Onset of Puberty) What to Do



### Call

Have someone call EMS/9-1-1 and get an AED. If you are alone with the child and are not in close proximity to a phone, do 5 cycles (2 minutes) of CPR before calling EMS/9-1-1. Take the child with you to call EMS/9-1-1 and get an AED, and then continue to provide care.



## Care

### 1. Do 30 compressions:

- Put 2 hands on the centre of the child's chest.
- Push down at least 1/3 of the chest's depth, pushing deeply and steadily.
- The chest compression rate should be between 100 and 120 beats per minute (30 compressions in 15 to 18 seconds).
- Allow the chest to fully recoil after each compression (do not lean on the chest).

### 2. Give 2 breaths:

- Open the airway by doing a head-tilt/chin-lift.
- Place your barrier device over the child's mouth and nose.
- If using a plastic face shield, seal your lips tightly over the child's mouth and pinch the nose. If using a face mask, ensure it is fitted tightly over the mouth and nose.
- Give 2 breaths. Each breath should last 1 second, with just enough volume to make the chest start to rise.



3. If both breaths go in, repeat the cycle of 30 compressions and 2 breaths.

## Cardiac Arrest, Baby (Less Than 12 Months)

### What to Do



### Call

Have someone call EMS/9-1-1 and get an AED. If you are alone with the baby and are not in close proximity to a phone, do 5 cycles (2 minutes) of CPR before calling EMS/9-1-1. Take the baby with you to call EMS/9-1-1 and get an AED, and then continue to provide care.

 Care


## 1. Do 30 compressions:

- Put 2 fingers on the middle of the baby's chest, just below the nipple line.
- Push down at least 1/3 of the chest's depth, pushing deep and pushing steady.
- The chest compression rate should be between 100 and 120 beats per minute (30 compressions in 15 to 18 seconds).
- Allow the chest to fully recoil after each compression (do not lean on the chest).



## 2. Give 2 breaths:

- Open the airway by doing a head-tilt/chin-lift.
- Place your barrier device over the baby's mouth and nose.



- If using a plastic face shield, seal your lips tightly over the baby's mouth and nose. If using a face mask, ensure it is fitted tightly over the mouth and nose.
- Give 2 breaths. Each breath should last 1 second, with just enough volume to make the chest start to rise.

## 3. If both breaths go in, repeat the cycle of 30 compressions and 2 breaths.

**Continue CPR Until...**

Once you begin CPR, continue giving sets of 30 chest compressions and 2 rescue breaths until:

- EMS personnel take over.
- An AED is available and there is no one else who can apply the pads to the person's chest.
- Another trained First Aider is available and can take over compressions.
- You have performed approximately 2 minutes of CPR for a child or baby (5 sets of 30 chest compressions and 2 rescue breaths) and you need to call EMS/9-1-1.
- You are alone and too tired to continue.
- The scene becomes unsafe.
- You notice an obvious sign of life, such as movement. If the person shows an obvious sign of life, stop CPR and check the person's ABCs. If the person is breathing, place him or her in the recovery position, and continue to monitor the person's condition until EMS personnel take over.

## What to Do If the Rescue Breaths Don't Go In

If the chest does not rise after the first breath:

1. Reposition the head to adjust the airway, and attempt to give another breath.
2. If the second breath does not go in, begin the CPR sequence again (30 chest compressions).

3. After the 30 chest compressions, look in the person's mouth. If you see an object, carefully remove it:
  - With one hand, grasp the lower jaw and tongue (if possible) while opening the mouth.
  - Place one finger from your other hand into the side of the person's mouth, then sweep behind the object to lift it out.



4. After looking in the mouth (even if nothing is visible) open the airway and attempt to give a breath:
  - If the breath goes in, give a second breath.
  - If the breath does not go in, resume the CPR sequence (starting with 30 chest compressions).

5. Continue to give CPR. After each set of compressions and before attempting to give rescue breaths, look for an object in the person's mouth and remove it if possible.



## Compression-Only CPR

Compression-only CPR uses chest compressions (without rescue breaths) to pump the heart and circulate oxygen that is already in the person's body. If you are unwilling or unable to give rescue breaths for any reason (e.g., you do not have a barrier device or the ill or injured person has facial trauma), compression-only CPR is acceptable. As a minimum, you should always perform compression-only CPR for any person in cardiac arrest. You are encouraged to perform traditional CPR (compressions with rescue breaths) whenever possible.

Compression-only CPR is always better than no care, and can be effective. However, if you are caring for a child, a baby, or any person who entered cardiac arrest because of a respiratory problem (e.g., choking, drowning, anaphylaxis, asphyxiation), the person may not have much oxygen remaining in his or her blood, and circulating deoxygenated blood is not very useful. In these circumstances, traditional CPR with rescue breaths is the recommended method of care.


## CPR Compression Depth for an Adult

When providing CPR to an adult, you should perform compressions in the middle of the chest on the lower half of the sternum. Push deeply and push steadily, ensuring a compression depth of at least 5 cm (about 2 in.), while avoiding excessive depths greater than 6 cm (about 2.5 in.). While there is a risk of injury with compression depths greater than 6 cm, these injuries tend to be non-life-threatening. It is more important to push deeply and push steadily than to worry about pushing too deeply.

## CPR Compression Depth for a Child or a Baby

Because of the variation in size among children or babies, compression depth for this age group is measured as a fraction of the total chest depth, not in centimetres. When providing CPR to a child or baby (i.e., a person who has not yet reached puberty), you should perform chest compressions on the lower half of the sternum and depress the chest to at least 1/3 of its depth. Ideally, chest compressions should be at the upper limit of 1/2 of the front-to-back (anteroposterior) depth.

CPR compression depths may be impacted by items such as heavy clothing (e.g., a winter coat or motorcycle jacket) or sports equipment. You should not delay starting CPR to remove them, but you may have to push harder to ensure that the minimum chest compression depth is reached.




## Chest Compression Rates

First Aiders should perform chest compressions at a rate of 100 to 120 beats per minute. Counting out loud as you give compressions can help you to keep a steady, even rhythm that allows time for adequate chest recoil. For compressions one through 12, say "one and two and three and four and five and..." up to 12. When you get to 13, say the numbers without saying "and" in between: "13, 14, 15, 16..." and so on. Push down as you say the number and come up as you say "and" or the second syllable of the number.

## Chest Recoil

You should avoid leaning on the person's chest wall in between compressions. Instead, ensure that the chest recoils fully. It is during recoil that the coronary arteries of the heart are filled with blood and the heart muscle gets oxygen. If you are leaning on the person's chest between compressions, the coronary arteries will not get sufficient blood flow and heart death can occur. Remember: Compression is for the brain, recoil is for the heart.



## ***Myth-Information: Upper-Body Strength Required for CPR***

**Myth:** Performing CPR compressions requires significant upper-body strength.

If you are performing compressions properly, the power comes from the weight of your upper body, not from your arm muscles. Incorrect technique or body position can cause your arms and shoulders to tire quickly when you are giving compressions. Avoid rocking back and forth, because this makes your compressions less effective and wastes your energy.

### ***Decreasing Compression Fraction (Off-Chest Time)***

During CPR, you should minimize interruptions to chest compressions. Any interruption decreases the total number of compressions given, and therefore the amount of blood that is circulated. Compressions also build up pressure in the person's arteries: when compressions are interrupted, the arterial pressure drops, which means that the first compressions you give after the interruption are rebuilding the pressure and not yet circulating blood effectively.

Interruptions should be limited to critical interventions, such as applying AED pads to the person, allowing the AED to analyze or administer a shock, and clearing the airway. Pause to reassess the person's ABCs only if there has been an obvious change in the person's condition. Lengthy ventilation steps (opening airway, excessive breathing) should be avoided. There is no need for you to reposition your hands after every compression—the hand is placed on the middle of chest.



## ***Special Considerations***

### **CPR for a Pregnant Woman**

When performing CPR for a pregnant woman, raising her right hip 7.5 to 10 cm (3 to 4 in.) will help blood return to the heart. If a bystander is available (or an appropriate object is within easy reach), ask the person to find a soft object that you can place under the woman's hip, but do not interrupt or delay starting CPR to find an object yourself.

### **Air in the Stomach**

When giving rescue breaths, it is important to keep the person's head tilted back and breathe with just enough force to cause the chest to rise. Failing to tilt the head back, giving too much air in one breath, or blowing too forcefully can push air into the person's stomach instead of into his or her lungs.

Air in the stomach can make someone vomit. When an unresponsive person vomits, the stomach contents may get into the lungs—this is called aspiration. Aspiration makes giving rescue breaths more difficult and reduces the person's chance of successful resuscitation. To prevent aspiration, remember to give breaths only until the chest starts to rise.

### **Vomiting**

In some situations a person may vomit while you are giving CPR. If this happens:

1. Turn the person onto his or her side, facing you.
2. Quickly wipe the person's mouth clean.
3. Reposition the person on his or her back and continue with CPR.

**CPR SUMMARY**

	<b>Hand Position</b>	<b>Compress</b>	<b>Breathe</b>	<b>Cycle</b>	<b>Compression Rate</b>
<b>Adult</b>	Two hands on the middle of the chest	At least 5 cm (about 2 in.)	Just enough volume to make the chest start to rise (1 second per breath)	30 compressions and 2 breaths	Rate of 100–120 per minute: 30 compressions in about 15–18 seconds.
<b>Child</b>	Two hands on the middle of the chest	At least 1/3 of the chest's depth	Just enough volume to make the chest start to rise (1 second per breath)	30 compressions and 2 breaths	Rate of 100–120 per minute: 30 compressions in about 15–18 seconds.
<b>Baby</b>	Two fingers on the middle of the chest (just below the nipple line)	At least 1/3 of the chest's depth	Gently, with just enough volume to make the chest start to rise (1 second per breath)	30 compressions and 2 breaths	Rate of 100–120 per minute: 30 compressions in about 15–18 seconds.



## Mouth-to-Nose Breathing

Sometimes you cannot seal your mouth over the person's mouth to give rescue breaths because:

- The person's jaw or mouth is injured.
- The person's jaw or mouth is shut too tightly to open.
- Your mouth is too small to cover the person's mouth.
- The person has blood coming out of his or her mouth.

If this happens, tilt the person's head as usual using a head-tilt/chin-lift and breathe into the person's nose. Block or close the mouth with your hand to stop air from escaping.

## Mouth-to-Stoma Breathing



Some people have had an operation to remove part of the trachea. They breathe through an opening called a stoma in the front of the neck. Because air passes directly into the trachea through the stoma instead of through the mouth and nose, you should give rescue breaths into the stoma. Block the person's mouth and nose with your hand to stop air from escaping.

## Automated External Defibrillation

While CPR can help prevent brain damage and death by keeping oxygenated blood moving throughout the body, an automated external defibrillator (AED) can correct the underlying problem for some people who go into sudden cardiac arrest.

Two abnormal heart rhythms in particular, ventricular fibrillation (V-fib) and ventricular tachycardia (V-tach), can

lead to sudden cardiac arrest. In V-fib, the heart muscle quivers weakly ("fibrillates") instead of contracting strongly. In V-tach, the heart muscle contracts too rapidly ("tachy-" means "fast"). Both of these rhythms impair the heart's ability to pump and circulate blood throughout the body and both are life-threatening.



In many cases, an electrical shock delivered by an AED can correct V-fib and V-tach rhythms. The shock disrupts the heart's electrical activity long enough for it to spontaneously develop an effective rhythm on its own. Using an AED as soon as possible (along with starting CPR immediately) gives the person the best chance of surviving cardiac arrest.

Different types of AEDs are available, but all are similar to operate and guide you using visual displays, voice prompts, or both. If your place of employment has an AED on site, know where it is located, how to operate it, and what the procedures around it are (e.g., for reporting its use or replacing its batteries). Also take note of the location of AEDs in public places that you frequent, such as shopping centres, airports, recreation centres, and sports arenas.

In more than 80% of all sudden cardiac deaths, the person's heart rhythm is "shockable" (ventricular tachycardia or ventricular fibrillation). In other words, defibrillation could have helped. If the person's heart has no electrical activity (asystole or flatline), defibrillation won't help. For each minute that the person has to wait for defibrillation, the chance of survival drops between 7 and 10%.

## Using an AED

1. Open and turn on the AED.
2. Prepare to apply the AED pads:
  - Remove or cut away clothing and undergarments to expose the person's chest, and move or remove any objects (including jewellery) that could come into contact with the pads.
  - If the person's chest is wet, dry it using a towel or gauze pad. Dry skin helps the AED pads to stick properly.
  - Small amounts of chest hair will not interfere with pad adhesion. If the person has thick chest hair, shave the areas where the pads will be placed using the razor included with the AED prep kit.



3. Check if the person has an implanted pacemaker:

- Look on the chest for a small scar and a lump about the size of a matchbox.
- Apply the AED pads approximately 2.5 cm (1 in.) away from the pacemaker.



4. Apply the AED pads:

- Use the appropriate size of pads—adult, child, or baby.
- A child is considered to be between 1 and 8 years old. If child AED pads are unavailable, use adult pads.



- A baby is considered to be 1 year old or less. If baby AED pads are unavailable, use child or adult pads.
- Follow the diagrams on the pads to position them correctly on the person.
- Pads must be placed at least 2.5 cm (1 in.) apart. If there is not enough space on the chest to leave 2.5 cm (1 in.) between the pads, place one pad on the person's chest (anterior) and one on the person's back (posterior).



5. Follow the AED's prompts to "Analyze Heart Rhythm:"
  - Most AEDs will begin to analyze the heart rhythm automatically, but some may require you to push an "analyze" button to start this process.
  - No one should touch the person while the AED is analyzing the heart rhythm because this could result in a faulty reading.



6. If the AED prompts you to do so, deliver the shock:
  - Stand clear and say "I'm clear, you're clear, everybody's clear."
  - Ensure that no one is touching the person (as they can be shocked too), then press the "shock" button.



7. After a shock is delivered (or if the AED determines that no shock is necessary), immediately resume CPR, starting with compressions.
8. Continue to follow the AED's automated prompts.



The AED will continue to check the heart rhythm every 2 minutes. Listen for prompts from the AED and continue giving CPR and using the AED until you notice an obvious sign of life or EMS personnel arrive. If you notice an obvious sign of life, stop CPR, but leave the AED turned on and the pads in place on the person's chest. Continue to follow the AED's prompts.

### ***Considerations for Safe and Effective AED Use***

#### **Environmental Considerations**

**Flammable or combustible materials:** Do not use an AED around flammable or combustible materials, such as gasoline or free-flowing oxygen.

**Metal surfaces:** It is safe to use an AED when the person is lying on a metal surface, as long as the AED pads do not touch the metal.

**Water:** If the person is in water, remove him or her from the water before using the AED. Remember that many liquids conduct electricity: if the person is lying in a puddle when the shock is delivered, anyone who is touching the same puddle is at risk of being shocked as well. It is safe to use an AED on ice or snow.

**Inclement weather:** It is safe to use AEDs in all weather conditions, including rain and snow, but their effectiveness can be reduced if the pads are wet. Provide a dry environment if possible (for example, shelter the person with umbrellas), but do not delay defibrillation to do so. An AED is designed to function in a range of weather conditions, but it is still a battery-operated electronic device, and excessive cold, heat, moisture, etc., can result in maintenance errors or prevent the pads from adhering properly. If this happens, move the person into a different environment, if possible, before attempting defibrillation again.

### Person-Specific Considerations

**Pregnancy:** It is safe to use an AED on a woman who is pregnant.

**Pacemakers and implantable cardioverter-defibrillators (ICDs):** A person who has a known arrhythmia (irregular heartbeat) may have a pacemaker or an ICD. These small devices are surgically implanted under the skin to automatically prevent or correct an irregular heartbeat. You may be able to see or feel the outline of the pacemaker or ICD in the area below the person's collarbone, or the person may wear a medical identification product indicating that he or she has a pacemaker or ICD. If the person has a pacemaker or ICD, adjust the pad placement to avoid placing the AED pads directly over the device, as this can interfere with the delivery of the shock. If you are not sure whether the person has an implanted device, place the pads as you normally would.

**Transdermal medication patches:** Some types of medications, including nitroglycerin and smoking cessation medications, are delivered through patches applied to the skin. Remove any medication patches that you see before applying AED pads. Wear gloves to prevent absorption of the drug through your own skin.

**Chest hair:** Time is critical in a cardiac arrest situation and sparse chest hair rarely interferes with pad adhesion. In most cases, you should proceed as you normally would—attach the AED pads, pressing firmly

to ensure adhesion. However, if the person has a great deal of thick chest hair and it seems like it could interfere with pad-to-skin contact, quickly shave the areas where the pads will be placed and then attach the pads.

**Jewellery and body piercings:** You do not need to remove the person's jewellery or body piercings before using an AED. However, if possible, avoid placing the AED pads directly over any metallic jewellery or piercings. If making a small adjustment to pad placement allows you to avoid the jewellery then do so, but if you must move the pads significantly it is better to place the pads directly over the jewellery.

### Using an AED on a Child or Baby

The procedure for using an AED on a child or baby is the same as the procedure for using an AED on an adult. Some AEDs come with pediatric AED pads that are smaller and designed to deliver a less powerful shock. These pads should be used on children up to 8 years of age or weighing less than 55 pounds. Other AEDs have a key or switch that configures the AED for use on a child. If pediatric AED pads are not available or the AED does not have a pediatric setting, it is safe to use the adult pads and setting on a child or baby.

### Medical Direction and AEDs

In some professions, defibrillation is designated as a controlled medical act. If you are providing care under the authority of a medical director, he or she may choose to adopt different protocols than those of the Canadian Red Cross. The specific protocols will be written in a standing order. You should always follow the instructions of your medical director. The Canadian Red Cross accepts no responsibility for any modifications made to the AED protocol set out in this text.

## AED Maintenance

AEDs require minimal maintenance, but it is important to check them regularly according to the manufacturer's instructions or your employer's policy. This ensures that they remain in good working order and are ready for use whenever they are needed.

To ensure that your AED is ready to use:

- Familiarize yourself with the owner's manual and follow the manufacturer's instructions for maintenance.
- Familiarize yourself with the method the AED uses to indicate the status of the device. Many AEDs have a status indicator that displays a symbol or illuminates to indicate that the AED is in proper working order and ready to respond. The status indicator may also display symbols indicating that routine maintenance is needed (e.g., a battery change) or that a problem with the device has been detected. Some AEDs have a warning indicator that will illuminate or beep if the AED is not in proper working order or requires maintenance.
- Check regularly to make sure that the battery is properly installed and has not reached its expiration date.
- Make sure AED pads are adequately stocked, stored in a sealed package, and within their expiration date.
- After using the AED, make sure that all supplies are restocked and that the device is in proper working order.
- If at any time the AED fails to work properly or warning indicators appear, take the AED out of service and contact the manufacturer or the appropriate person at your place of employment (according to your employer's policy). You may need to return the AED to the manufacturer for service. If the AED stops working during an emergency, continue giving CPR until EMS personnel take over.



## Working as a Team

Remember: when you are giving CPR, you want to give high-quality compressions at the appropriate depth and rate and to minimize interruptions to compressions. If you are the only trained First Aider at the scene, you will begin to tire as you give CPR, and the quality of your compressions will diminish. You will also need to stop CPR to ready the AED for use when it arrives, which means that during that time there will be no oxygenated blood moving through the person's body.

Working as a team can increase the chance of survival for the person in cardiac arrest. First Aiders can share the responsibility for giving compressions, switching off every 2 minutes. This reduces fatigue and leads to higher-quality compressions.

Having two or more First Aiders at the scene also minimizes interruptions to chest compressions when the AED arrives. One First Aider should begin CPR while the second calls EMS/9-1-1, obtains the AED, and readies it for use by turning the device on, applying the pads to the person's chest, and plugging in the connector cable, if necessary. The First Aider performing CPR should not pause CPR until the AED is ready to analyze the person's heart rhythm and the second First Aider tells everyone to stand clear. While the AED is analyzing, the First Aiders should switch roles. The First Aider who is taking over compressions should hover with his or her hands positioned just above the person's chest so that he or she can immediately start compressions as soon as the AED prompts that a shock was delivered or that no shock was advised. The First Aiders then switch roles every time the AED analyzes the person's heart rhythm, which occurs every 2 minutes.

# 7

# Breathing Emergencies





A person who is having difficulty breathing is in respiratory distress. A person who is not breathing is in respiratory arrest. Both respiratory distress and respiratory arrest are breathing emergencies. Respiratory distress is especially dangerous in children, as it can quickly lead to cardiac arrest. Any child showing signs of respiratory distress is in a potentially life-threatening situation and must receive immediate care.

When a person is experiencing a breathing emergency, the oxygen supply to the person's body is either greatly reduced or cut off entirely, so it is important to act at once. If breathing stops or is restricted long enough, the person will become unresponsive, the heart will stop beating, and body systems will quickly fail. Brain cells begin to die after 4 to 6 minutes without oxygen.

A breathing problem can be identified by watching and listening to the person's breathing and by asking the person how he or she feels. If a person is having trouble breathing, do not wait to see if his or her condition improves without intervention, but begin providing care immediately.

## Respiratory Distress

### Common Causes

Respiratory distress may be caused by any of the following:

- Hyperventilation
- Asthma or chronic obstructive pulmonary disease (COPD)
- Pneumonia or bronchitis
- An allergic reaction
- Anaphylaxis
- A heart attack or heart failure
- Chest trauma
- Poisoning
- A drug overdose
- Electrocution
- Certain mental health conditions (e.g., panic disorders)

### What to Look For

The following are signs and symptoms of respiratory distress:

- Shortness of breath or gasping for breath
- Trouble speaking in complete sentences (due to difficulty breathing)
- Wheezing, gurgling, or high-pitched noises
- Breathing abnormally quickly or slowly
- Unusually deep or shallow breathing
- Cool, moist skin
- Bluish or ashen (grey) skin
- Flushed or pale skin
- Feelings of fear
- Dizziness or light-headedness

## What to Do



### Call

Call EMS/9-1-1.



### Care

1. If the person carries medication for respiratory distress, offer to help the person take his or her medication.
2. Encourage the person to sit down in a comfortable position (leaning forward may help make breathing easier).
3. Provide reassurance, as this can help reduce anxiety and help the person control his or her breathing.
4. Monitor the person's condition and provide continual care until EMS personnel arrive.

## Hyperventilation

Hyperventilation is a condition in which a person is breathing much more quickly than usual. This upsets the body's balance of oxygen and carbon dioxide.

## Common Causes

The following are common causes of hyperventilation:

- Strong emotions such as excitement, fear, or anxiety
- Asthma
- Injuries, especially injuries to the head
- Exercise
- Life-threatening bleeding

## Prevention

If you tend to hyperventilate due to anxiety, panic, or stress, relaxation techniques such as breathing exercises may help. If you hyperventilate due to a diagnosed medical condition, speaking to your doctor will help you learn how to treat or control it.

## What to Look For

The following are signs and symptoms of hyperventilation:

- Rapid, shallow breathing
- A feeling of suffocating or not getting enough air
- Fear, anxiety, or confusion
- A feeling of dizziness
- Numbness or tingling of the fingers and toes
- Muscle contractions, usually in the hands, feet, arms, and legs

## What to Do



### Call

It is not always necessary to call EMS/9-1-1 for a person who is hyperventilating. You should call EMS/9-1-1 and get an AED if the hyperventilation does not stop after a few minutes, the person becomes unresponsive, or you suspect that the person is hyperventilating because of an injury or illness.



### Care

1. Encourage the person to take controlled breaths by breathing in slowly, holding his or her breath for a few seconds, and then gradually exhaling.

## Myth-Information

**Myth:** If a person is hyperventilating, you should have him or her breathe into a paper bag.

This practice is unsafe and not recommended as a way to care for a hyperventilating person. The best way to treat hyperventilation is to encourage the person to take slow, controlled breaths.



## Asthma

Many people have asthma—a chronic illness in which certain substances or conditions (i.e., triggers) cause inflammation and swelling of the bronchioles (i.e., the small tubes at the base of the lungs), making it harder for air to move in and out of the lungs. Asthma is more common in children. People who have asthma usually know what can trigger an attack and take measures to avoid these triggers. Asthma is usually controlled with medication.

## Common Causes

Asthma can be triggered by:

- Air pollution or poor air quality (e.g., cigarette smoke)
- Allergies (e.g., to pollen, food, drugs, insect stings, or animals)
- Temperature fluctuations, extreme humidity, or extreme dryness
- Strong odours (e.g., perfume, cologne, scented cleaning products)
- Colds and flus
- Physical activity
- Respiratory infections
- Stress or anxiety

## Prevention

If you have asthma, the following precautions may help to prevent attacks:

- Know what triggers your attacks and avoid them if possible.
- Ensure that prescribed medication is always easily accessible in case of an attack.

If a child has asthma:

- Make sure that anyone who supervises the child knows about the asthma and how to help give medication if necessary.

## Asthma Medications

There are two main types of medication used by people with asthma: Long-term control medication and quick-relief (rescue) medication. A person with asthma may take one, both, or neither of them. Both long-term control medications and quick-relief (rescue) medications may be given through an inhaler, through a nebulizer, or orally.



Long-term control medications, sometimes called "preventers," are taken regularly, whether or not signs and symptoms of asthma are present. These medications help prevent asthma attacks by reducing inflammation and swelling and by making the bronchioles less sensitive to triggers.

Quick-relief (rescue) medications are taken when a person is experiencing an acute asthma attack. These medications are sometimes called "bronchodilators," because they work quickly to relax the muscles that tighten around the bronchioles, opening (dilating) them immediately so that the person can breathe more easily.

## What to Look For



Even when a person takes steps to manage his or her asthma by avoiding triggers and taking prescribed long-term control medications, he or she may still occasionally experience asthma attacks. The signs and symptoms of an asthma attack include the following:

- Wheezing or coughing, especially when exhaling
- Gasping for air
- Shortness of breath (feeling unable to get enough air into the lungs)
- Rapid, shallow breathing (or trouble breathing)
- Anxiety and fear
- Tightness in the chest
- Tingling in the hands and feet
- Sweating
- Inability to say more than a few words without pausing to breathe

## What to Do

### Call

Call EMS/9-1-1 and get an AED if the person is struggling to breathe, the person's breathing does not improve after taking his or her medication, or the person becomes unresponsive.

### Care

1. Eliminate any asthma triggers, if possible. If something in the environment is causing the attack, move the person away from the area if it is safe to do so.
2. Help the person get into a comfortable position and provide reassurance.

3. Help the person to take any prescribed quick-relief asthma medication that he or she has available.



## Providing Care for an Asthma Attack: Using an Asthma Action Plan

An individual who is at risk for ongoing asthma attacks may have a written plan that outlines the assessment and treatment steps, and clarifies when to call EMS/9-1-1. This is most common for children in child care situations. If you are a caregiver for a person who has an asthma action plan, familiarize yourself with it and be sure you understand what to do if the person has an attack.

### Using an Inhaler

1. Rapidly shake the inhaler 3 or 4 times.



3. Breathe out as much air as possible, away from the inhaler.
4. Bring the inhaler to the mouth.
5. Press the top of the inhaler while taking one slow, full breath.



2. Remove the cap from the inhaler.



6. Hold the breath for as long as is comfortable (up to 10 seconds), and then breathe out.

### Using an Inhaler with a Spacer

1. Rapidly shake the inhaler 3 or 4 times.
2. Remove the cap from the inhaler.
3. Remove the spacer's cap (if it has one), and put the inhaler into the spacer.
4. Bring the spacer to the mouth.
5. Press the top of the inhaler once to dispense the medication into the spacer.
6. Take slow, deep breaths, holding each breath for a several seconds (if possible).



### Allergic Reactions

Normally, our immune systems help to keep us healthy by fighting off harmful pathogens that can cause disease. In a person with an allergy, however, the immune system can react to normally harmless substances to produce allergic reactions.

An allergic reaction occurs in two stages. First, the immune system mistakes a normally harmless substance for a dangerous invader and produces antibodies that remain on alert for that particular allergen. When the body is exposed to the allergen again, these antibodies release a number of immune system chemicals, such as histamine, that cause allergy symptoms.

While certain substances are more common as allergens (e.g., peanuts and pollen), almost anything can produce an allergic reaction. The severity of an allergic reaction depends on the person. It can be as

minor as a slight irritation or as serious as a life-threatening emergency (anaphylaxis). The signs and symptoms of an allergic reaction depend on the trigger, but they typically involve the airways, skin, sinuses, nasal passages, cardiovascular system, and digestive systems.

### Preventing Allergic Reactions in a Child You Are Caring For

Discuss the allergy with the parent or guardian before you begin caring for the child, and ask at least the following questions:

- What is your child allergic to?
- What signs will tell us that your child is having an allergic reaction?
- What should we do if your child has a reaction?
- Has the doctor prescribed any allergy medication? For instance, should you give us your child's epinephrine auto-injector?

Make a list of the allergies of each child under your care, and keep it in a location where you can refer to it easily, especially in cooking and food service areas. Be sure to read all the ingredients on all food package labels when preparing and cooking food. Before serving a food that is a common allergen (e.g., eggs, shellfish, milk, nuts), ensure that the parent or guardian has given it to the child at home without causing a reaction.

### What to Look For



### What to Do



#### Call

Call EMS/9-1-1 and get an AED if the reaction is severe, the person is struggling to breathe, or the person loses responsiveness.



#### Care

1. Calm and reassure the person.
2. Try to identify the allergen and have the person avoid further contact with it.
3. Watch the person for signs of increasing distress.
4. If the person uses a medication to control allergic reactions (such as antihistamines), help him or her to take it.

Signs and symptoms of an allergic reaction can range from mild to very severe and include the following:

- Runny, itchy, or stuffy nose
- Sneezing
- Watery, itchy, red, or swollen eyes
- Nausea, vomiting, or diarrhea
- A rash or hives (raised, itchy areas of skin)
- Tingling of the mouth
- Swelling of the lips, tongue, face, or throat
- Coughing, chest tightness, wheezing, or shortness of breath
- Weakness, dizziness, or confusion

## Antihistamines

An antihistamine is a medication that counteracts the effects of histamine (a chemical released by the body during an allergic reaction). Antihistamines are supplied as pills, capsules, or liquids and are taken by mouth. The person should take the antihistamine according to the medication label and his or her healthcare provider's instructions.



## Anaphylaxis

Anaphylaxis is a severe allergic reaction. While mild allergic reactions are localized, anaphylaxis is a body-wide reaction and it can cause system-wide inflammation and swelling. In some cases, the air passages swell, making it difficult to breathe. It is important to act quickly when a person is experiencing anaphylaxis, as it can lead to death if it is not cared for immediately.

## Common Causes

Anything that causes other allergic reactions can also cause anaphylaxis, and the causes vary from one person to another. The most common allergens that trigger anaphylaxis include:

- Insect stings
- Food
- Medications

## Prevention

The following steps may help to prevent an anaphylactic incident:

- Avoid the substances, foods, or insects that cause reactions.
- Wear a medical identification product and carry the appropriate medication at all times.

## What to Look For



from two or more of these categories—especially after contact with an allergen—you should provide care for anaphylaxis:

- Skin (e.g., swelling of the lips, face, neck, ears, and/or hands, a raised, itchy, blotchy rash, flushing, or hives)
- Breathing (e.g., a feeling of tightness in the chest or throat, coughing, wheezing, or high-pitched noises)
- Alertness (e.g., weakness, dizziness, or unresponsiveness)
- Stomach (e.g., stomach cramps, nausea, vomiting, or diarrhea)

## What to Do



### Call

Call EMS/9-1-1 and get an AED.

The signs and symptoms of anaphylaxis may be similar to the signs and symptoms of a mild allergic reaction, but they are more pronounced. A person experiencing an anaphylactic emergency may develop one or more signs and symptoms within seconds or minutes of coming into contact with the allergen.

Anaphylaxis can affect a variety of body systems and can present in various ways. If a person exhibits signs and symptoms

from two or more of these categories—especially after contact with an allergen—you should provide care for anaphylaxis:

- Skin (e.g., swelling of the lips, face, neck, ears, and/or hands, a raised, itchy, blotchy rash, flushing, or hives)
- Breathing (e.g., a feeling of tightness in the chest or throat, coughing, wheezing, or high-pitched noises)
- Alertness (e.g., weakness, dizziness, or unresponsiveness)
- Stomach (e.g., stomach cramps, nausea, vomiting, or diarrhea)

## What to Do



### Call

Call EMS/9-1-1 and get an AED.



## Care

1. If the person has an epinephrine auto-injector, help him or her to use it.
2. Provide reassurance and encourage the person to breathe normally.
3. Help the person get into a comfortable position.



If the person's condition does not improve 5 minutes after the initial dose of epinephrine and EMS personnel have not yet arrived, help the person take a second dose, if available. The second dose should be given in the leg that you did not use for the first dose.

If responsive, the person may want to take additional medication such as an antihistamine.

### *Using an Epinephrine Auto-Injector*

1. Remove the safety cap.



2. Place the injection tip against the middle third of the outer thigh and push the epinephrine auto-injector firmly against the thigh with a quick motion. A click should be heard.
3. Hold in place as directed, usually for 5 to 10 seconds.

4. Remove the epinephrine auto-injector. Use caution if the needle tip is exposed.



5. Rub the injection site for 30 seconds to help with medication absorption.



6. Make sure the used epinephrine auto-injector is placed in a rigid container and goes with the person to the hospital.
7. Have the person rest quietly until EMS personnel arrive.



## Epinephrine

Epinephrine is a drug that slows or stops the effects of anaphylaxis. If a person has a risk of experiencing anaphylaxis, he or she may carry an epinephrine auto-injector (a syringe system that contains a single dose of epinephrine). Different brands of epinephrine auto-injectors may be available, but all work in a similar fashion. Some have audio prompts to guide the user.

Different auto-injectors contain different doses based on the weight of the person (0.15 mg for children weighing between up to 30 kg (66 lb.), and 0.3 mg for children and adults weighing more than 30 kg (66 lb.)).

Many healthcare providers advise that people with a known history of anaphylaxis carry at least two doses of epinephrine (two auto-injectors) with them at all times. This is because more than one dose may be needed to stop an anaphylactic reaction. The second dose should be administered if the person is still having signs and symptoms of anaphylaxis 5 minutes after the first dose.

# 8 Wound Care



## Stages of Wound Healing

Wounds go through several phases during the healing process. The first stage is the body's natural response to injury. It is at this stage that the characteristic signs of inflammation can be seen: heat, swelling, and pain. Infection is most prominent during the first stage of healing, but can occur at any time.

The second stage is where tissue is rebuilt by the body. The area will slowly heal with tissue that is sensitive and pink or red. This new skin tissue will slowly toughen and thicken during the final stage of healing.



Wounds such as bruises, scrapes, and small cuts are very common injuries, and are caused in a variety of ways.

### Prevention

The following general tips will help you to avoid both serious and minor wounds:


- Develop safe habits, such as never running with sharp objects.
- Use proper safety equipment when playing sports or engaging in recreational activities.
- Wear proper safety equipment in the workplace, and request training before using any potentially dangerous tools or equipment.
- Stay alert and focused when engaged in any potentially risky activity.

## Bandaging Guidelines

- Use clean, preferably sterile, dressings.
- Check circulation below the injury before and after applying a bandage. If circulation is reduced, loosen the bandage.
- If blood soaks through the bandage, place more dressings and another bandage on top. The dressing that is in contact with the wound should remain in place and not be removed.
- Dressings and bandages should be kept clean and dry to decrease the risk of infection. Once bleeding is under control and the wound has begun to heal, dressings that become wet should be replaced with dry dressings.



### Folding a Triangular Bandage Into a Broad or Narrow Bandage



1. Fold the tip of the triangle down to touch the base.
2. Fold the bandage in half lengthwise to make a broad bandage.
3. Fold it in half again to make a narrow bandage.

## Cuts and Scrapes

A cut is a wound where the skin has been split open. The edges of the cut can be jagged or smooth. Scrapes are wounds where the skin has been rubbed or scraped away. Signs and symptoms of a cut or scrape may include pain and bleeding.

### What to Do



#### Call

Call EMS/9-1-1 if you suspect that there may be more serious injuries.



#### Care

1. There is usually minimal bleeding with cuts and scrapes, but if the wound is bleeding significantly, apply direct pressure until it stops.
2. If possible, rinse the wound for 5 minutes with clean, running tap water.



3. If an antibiotic ointment or cream is available, ask the person if he or she has a sensitivity to any antibiotics, such as penicillin. If not, suggest the person apply it to the wound.



4. Cover the wound with a sterile non-stick dressing and/or bandage.
5. Ensure that the person knows to watch for signs of infection over the next few days.



## Puncture Wounds

A puncture wound occurs when a hole in the skin is created by a pointed object, such as nails, pieces of glass, and animal teeth.

### Common Causes

Puncture wounds may be caused by:

- Injuries from pointed objects such as nails or pieces of glass
- Animal bites

### Prevention

The following may help you to avoid injuries that cause puncture wounds:

- Stay away from unfamiliar animals.
- Wear shoes when walking outside.
- Clean up sharp objects, like glass, right away. Dispose of sharp objects safely (e.g., do not put broken glass directly into a garbage bag, and, if possible, remove nails from objects before putting the objects into a garbage bag).

### What to Look For

These signs may indicate an injury caused by a puncture wound:

- Bleeding
- Bruising
- A hole where the object went through the skin

### What to Do



#### Call

Call EMS/9-1-1 and get an AED if the wound is deep or large.



#### Care

1. If the wound is bleeding, apply direct pressure until the bleeding stops.
2. Once the bleeding is controlled, wash the wound thoroughly with water. If possible, rinse the wound for 5 minutes with clean, running tap water.
3. Cover the wound with a sterile dressing and/or bandage.
4. Ensure that the person knows to watch for signs of infection over the next few days.

## Scalp Injury

### What to Do



When performing a secondary assessment of a person with a suspected scalp injury, check the scalp thoroughly to ensure that you have uncovered the full extent of the injury.



Because the scalp contains many blood vessels, even small scalp wounds can have significant bleeding.

 **Call**

Call EMS/9-1-1 and get an AED if you are not sure how serious the scalp injury is or if you feel a dip, a soft area, or pieces of bone.

 **Care**

1. Put dressings on the wound and have the person hold them in place to control the bleeding.



2. Secure the dressings with a bandage.



3. If you feel a dip, a soft area, or pieces of bone:
  - Treat the injury as a head injury.
  - Put direct pressure on the wound only if the bleeding is life-threatening. Otherwise, try to control the bleeding with pressure on the area around the wound.
4. Provide continual care until EMS personnel arrive.

**Infection**

An infection occurs when dirt, foreign bodies, or other materials carrying germs get into a wound. Infections can be localized (e.g., in a cut) or systemic (throughout the body). Infections accompanied by nausea, fever, or general malaise are typically systemic, and can be life-threatening.

**Prevention**

The following general tips help to minimize the risk of infection:

- Always wash your hands before and after giving first aid.
- Wear gloves whenever possible if you will be coming into contact with someone's bodily fluids.
- Keep your immunizations up to date. If you have been wounded and do not know when your last tetanus shot was, seek medical attention.
- Keep any wound clean and wash it regularly. Change the dressing and bandages if they become dirty or wet.
- Use sterile dressings whenever possible. If sterile dressings are unavailable, use the cleanest option possible.

## What to Look For

Any of the following signs around an injury may indicate an infection:

- Redness
- Red streaks moving away from the wound
- Pus
- Heat or warmth
- Swelling
- Tenderness
- Excessive itchiness



## Chemical Burns

The following precautions will help prevent chemical burns:

- Store chemicals in their original containers.
- Wear protective gear when handling chemicals.
- Wash your hands after touching chemicals.
- Get trained in a hazardous materials training program, such as the Workplace Hazardous Materials Information System (WHMIS)/Globally Harmonized System (GHS) of Classification and Labelling of Chemicals.
- Read the label before using a product.
- Be aware of caustic plants in your area.

## Burns

Burns are soft-tissue injuries caused by chemicals, electricity, heat, or radiation.

### Prevention

There are different methods of prevention for different types of burns.

## Electrical Burns

The following tips will help prevent electrical burns:

- Keep electrical appliances away from water.
- If an electrical cord is frayed, fix it or dispose of it.
- If there are young children in the building, cover electrical outlets.
- Before approaching a person with a suspected electrical burn, make sure that trained personnel turn off the electrical current.

## Burns from Lightning Strikes

The following precautions will help prevent burns from lightning strikes:

- As soon as a storm is seen or heard, stop swimming or boating and get away from the water, because water conducts electricity.
- Stay indoors during thunderstorms. A picnic shelter or car (with the windows rolled up) will also provide some protection.
- If caught outside, stay away from telephone poles and tall trees. Stay off hilltops and try to crouch down in a ravine or valley if shelters are not available nearby.
- Stay away from things that conduct electricity, such as farm equipment, small metal vehicles (e.g., motorcycles, bicycles, and golf carts), wire fences, clotheslines, metal pipes, and railings.

## Thermal Burns

The following precautions will help prevent thermal burns:

- When cooking on the stove, turn the pot handles in and use only the back burners when possible.
- Keep the hot water tank temperature at or below 49°C (120°F).
- Keep children away from heat sources and appliances such as ovens, barbecues, space heaters, woodstoves, candles, and fireplaces.
- Keep hot drinks out of children's reach.

## Sunburns

The following precautions will help prevent sunburns:

- Limit exposure to the sun between 10:00 A.M. and 3:00 P.M., if possible.
- Wear light coloured clothing that covers as much of the body as possible.
- Use a broad-spectrum sunscreen with a sun protection factor (SPF) of at least 30 and apply it 15 to 30 minutes before going outdoors. Reapply sunscreen at least every 2 hours, as well as after being in the water and after sweating.

## What to Do



### Call

Call EMS/9-1-1 and get an AED immediately if:

- The burns make it difficult for the person to breathe.
- The person is in a great deal of pain or becomes unresponsive.
- The burns were caused by chemicals, explosions, or electricity.
- The burns involve a large amount of blistering or broken skin, or the burns cover the face, neck, hands, genitals, or a larger surface area.

If at least one of the above conditions is true and you are alone, call EMS/9-1-1 yourself, get an AED, and then return to care for the person. If it a superficial burn, you should not need to call EMS/9-1-1, unless the person is in a great deal of pain or becomes unresponsive. Always call EMS/9-1-1 for a full thickness burn, regardless of size.



### Care

1. While the care for all burns is similar, specific care steps can vary depending on the cause of the burn. Care should be taken to monitor for hypothermia when cooling large burns. This is particularly important in children. Cover the person with a blanket if the person complains of feeling cold.

## Thermal Burns

Thermal burns are caused by heat, hot liquid, steam, or open flames. The care that is required varies by the thickness of the burn, so you should determine the burn's seriousness before beginning to provide care.

### Superficial Burns

### What to Look For



- Redness
- Pain
- Possible swelling

## What to Do



### Care

1. Cool the affected area with clean running or standing water for at least 10 minutes. A clean cool or cold (but not freezing) compress can be used as a substitute.



## Partial Thickness Burns

### What to Look For



- Redness
- Pain
- Possible swelling
- Blisters

## What to Do



### Care

1. Cool the affected area with clean running or standing water for at least 10 minutes. A clean cool or cold (but not freezing) compress can be used as a substitute.
2. Remove jewellery and clothing from the burn site, but do not attempt to move anything that is stuck to the skin.
3. Once it is cool, cover the burn loosely with a dry, sterile dressing, preferably non-stick gauze.
4. Encourage the person to seek medical attention even if it is not necessary to call EMS/9-1-1.



## Full Thickness Burns

### What to Look For



- Redness
- Pain (may not be present in the most severely burned areas due to nerve damage)
- Possible swelling
- Blisters
- Charred or waxy, white flesh

### What to Do



#### Care

1. Stop the burn from worsening by cooling the affected area with clean running or standing water for at least 10 minutes. A clean, cool or cold (but not freezing) compress can be used as a substitute.
2. Remove jewellery and clothing from the burn site, but do not attempt to move anything that is stuck to the skin.
3. Once it is cool, cover the burn loosely with a dry, sterile dressing, preferably non-stick gauze.
4. Have the person lie down until EMS personnel arrive.



## Chemical Burns

Chemical burns can be caused by a wet or dry caustic chemical. Use caution with dry caustic chemicals, as they may spread or react if they become wet.

### What to Do



#### Care

1. Wear protective equipment to avoid being burned yourself.
2. If there are dry chemicals present, brush them off the person's skin before flushing with water.
3. Flush the affected areas with large amounts of cool running water for at least 15 minutes, or until EMS personnel arrive. Flush the chemicals away from areas of the body that have not been contaminated.
4. Remove any clothing that is wet or that has been contaminated by the chemical.
5. Refer to the appropriate Material Safety Data Sheet (MSDS) for additional first aid measures, if it is available.

## Electrical Burns

Electrical burns are caused by powerful electrical currents. They are typically caused by a person either touching a live electrical circuit or being struck by lightning. Because electricity and lightning can affect the heart, it is important to monitor the person's ABCs closely.

### What to Do



#### Care

1. Treat the person as if he or she has a head and/or spinal injury.
2. Look for two burns (the entry and exit points) and care for them as you would care for thermal burns.

## Radiation Burns

The most common type of radiation burn is a sunburn, but they can also be caused by exposure to radioactive substances.

### What to Do



#### Care

1. If the burn results from exposure to a radioactive substance, consult the appropriate workplace safety system (e.g., WHMIS) for specific first aid steps.
2. Care for radiation burns as you would care for thermal burns.

## Special Considerations When Caring for Burns

Remember the following special considerations when providing care for burns:

- Don't use ointments on partial or full thickness burns.
- Blisters are a natural cooling system. Leave them in place.
- Touch a burn only with sterile or clean dressings.
- Do not use absorbent cotton or pull clothes over any burned area.

Pay close attention to the person's airway. Look for signs of burn injuries around the face. If you suspect that the airway or the lungs may be burned, monitor the person's breathing closely.



## Bruises

A bruise is a discoloured area of the skin that is created when blood and other fluids seep into nearby tissues.

### What to Look For

The following are signs and symptoms of bruising:

- Discoloured skin (red, purple, black, or blue areas)
- Swelling
- Pain

## What to Do



### Call

If the person is in severe pain or cannot move a body part without pain, or you think the force that caused the injury was great enough to cause serious damage or life-threatening bleeding, call EMS/9-1-1 and get an AED immediately. You may be dealing with internal bleeding, head and/or spinal injuries, or a bone, muscle, or joint injury. Care for the injury accordingly.




### Care

1. Using ice or another cold source, cool the injured area to help reduce pain and swelling. Place some sort of cloth, towel, or pad between the cold source and the skin to reduce the risk of freezing.



2. Apply the cold source for 20 minutes and then take it off for 20 to 30 minutes. Continue to do this for as long as the person keeps feeling pain.



## Splinters

A splinter (or "sliver") is a sharp, thin piece of material that is stuck in the surface of the skin. If the splinter is completely below the skin, encourage the person to seek medical attention rather than attempting to remove it yourself.

## What to Do



### Care

1. Gently grab the exposed end of the splinter with tweezers and carefully pull it out.
2. Wash the area with water and apply a bandage.





## Nosebleeds

### Common Causes

Nosebleeds may be caused by the following:

- Forceful nose blowing
- High blood pressure
- Dry weather conditions
- Trauma to the nose or head
- Bleeding disorders

If you know that the bleeding is caused by an object in the nose, refer to the care for foreign objects in the nose.

### Prevention

To prevent nosebleeds:

- Use a humidifier if the air indoors is dry.
- Wear protective athletic equipment when participating in sports that could cause injuries to the nose.
- Practice gentle nose blowing.
- Teach children not to pick their noses.

### What to Do



#### Call

Call EMS/9-1-1 if the bleeding continues for more than 15 minutes, the bleeding is the result of a head injury, or the person is losing a large amount of blood.



#### Care

1. Have the person sit with the head slightly forward, pinching the nostrils, for 10 to 15 minutes. If the nosebleed was caused by a severe head injury, do not pinch the nose.
2. Once you have controlled the bleeding, tell the person to avoid rubbing, blowing, or picking his or her nose because this could start the bleeding again.
3. If the person loses responsiveness, place him or her in the recovery position to allow blood to drain from the nose.



## Foreign Objects in the Nose

### What to Look For

An object in the nose may be visible, and may also cause the following signs and symptoms:

- Unusual noises when the person breathes through his or her nose
- Bleeding
- Nose deformity

### What to Do



#### Care

1. If you can easily see and grasp the object, remove it.
2. If the object cannot be dislodged easily, leave it in place for a medical professional to remove.
3. Prevent the person from attempting to remove the object and from blowing his or her nose.



## Knocked-Out Teeth

### Common Causes

A tooth can be knocked out by any kind of blow that involves the mouth.

### Prevention

The following tips may avoid an injury that results in knocked-out teeth:

- Wear appropriate equipment when playing sports, such as a mouth guard or face mask.
- Always wear a seat belt while in the car and do not eat or drink in a moving car.

### What to Look For

Aside from a tooth that is visibly missing, signs and symptoms of knocked-out teeth include the following:

- Bleeding (although this is often very minimal)
- Pain in the mouth

### What to Do



#### Call

Call EMS/9-1-1 and get an AED if the knocked-out tooth was caused by a forceful blow to the head, especially if the person is unresponsive, or you suspect that there may be other more serious injuries.



## Care

1. Control any bleeding by having the person bite down on a clean dressing.



2. Carefully pick up the tooth by the crown (the whiter part), not the root.



4. Get the person to a dentist as soon as possible. The greatest chance for repair is during the first hour after the tooth is knocked out.



An injury that knocks out a tooth can also be powerful enough to cause jaw, head, or spinal injuries.

3. Put the tooth in propolis, egg white, coconut water, whole milk, or saline, if available, and keep it with the person. If none of these are available, wrap the tooth in gauze (or a clean cloth) with some of the person's own saliva. Seal the container and label it with the name of the person, the date, and the time.



## Eye Injuries

Eye injuries can have long-term consequences for a person's vision, so they must always be treated with extreme care. The eye is an extremely sensitive organ: Avoid touching the eye or putting pressure on or around it, as this can cause further damage. Wounds involving the eyelids or eye sockets should also be treated as eye injuries.

### Common Causes

An eye injury may be caused by the following:

- Foreign objects or particles in the eye
- An impact to the eye
- Radiation or burns
- Chemicals or other caustic materials

## Prevention

The following may help to prevent an eye injury:

- Wearing appropriate equipment in the workplace (e.g., safety glasses)
- Wearing appropriate equipment when playing sports (e.g., wearing a helmet with a visor or face protector when playing hockey)

## What to Look For

An eye injury may include the following signs or symptoms:

- Pain and irritation in the eye
- Redness of the eye
- Difficulty opening the eye
- Problems with vision
- Watering of the eye
- Deformities of the eye

## What to Do



### Call

Call EMS/9-1-1 if there is an impaled object in or near the eye, the eye is out of the socket, or the eye has been exposed to a chemical or caustic substance.



### Care

If an object is impaled in or around the eye:


1. Have the person lie on his or her back and keep as still as possible.
2. Stabilize the object with bulky dressings.
3. Cover both eyes with gauze or another light material.
4. Provide comfort and reassurance until EMS personnel arrive.

If there is a foreign object in the eye but it is not impaled:

1. Try to remove the foreign object by having the person blink several times. The eye will produce tears that may wash out the object. An object that is not touching the eye itself (e.g., in the corner of the eye) may be removed with a moist cotton-tipped applicator.
2. Clean away any dirt around the eye and then gently flush the eye with running water (letting the water run away from the unaffected eye).
3. If these steps do not remove the object, the person should seek medical attention.

If there is a chemical or other caustic substance in the eye:

1. Gently flush the eye with running water (away from the unaffected eye) for at least 15 minutes, or until EMS personnel arrive.



If the eyes were flash burned (e.g., while welding):

1. Cover the eyes with a cool, wet cloth.
2. Make sure the person gets medical attention.

## Ear Injuries

### Prevention

The following tips can help prevent ear injuries:

- Wear proper protection when using loud equipment such as lawn mowers and chainsaws.
- Wear a helmet and other recommended safety gear when participating in sports and physical activities.
- Cover the ears when there is a risk of foreign objects or substances entering the ear (e.g., wear a swim cap when participating in water activities).

### Common Causes

An ear injury may be caused by the following:

- An impact to the head
- Cuts or tears to the ear(s)
- Loud noises
- Blast injuries
- Foreign objects or substances in the ear

### What to Look For

Signs and symptoms of an ear injury include the following:

- Blood or other fluid coming from within the ear
- Hearing problems
- Sudden or intense pain in the ear
- Swelling or deformity of the ear

### What to Do



#### Call

Call EMS/9-1-1 and get an AED if there is blood or other fluid draining from the ear, or if the ear injury is the result of an explosion or pressure (e.g., while scuba diving).



#### Care

1. If the injury is an external wound, treat it the same way you would treat a wound on any other part of the body.
2. If there is a foreign object in the ear but you don't suspect a head and/or spinal injury, and it looks as if the object can be easily removed:
  - Tilt the head to the affected side, then gently tap the ear to loosen the object.
  - Attempt to grasp the object and pull it out.





## Impaled Objects

An impaled object is anything that has penetrated through the skin and is stuck in the tissues below.

### Common Causes

An object can become impaled when any type of force causes it to penetrate the skin and underlying tissue.

### What to Look For

An impaled object will be visibly protruding from the person's body, and may cause the following signs and symptoms:

- Shock
- Pain
- Bleeding

### What to Do



#### Call

Call EMS/9-1-1.



#### Care

1. Keep the person still and leave the object in place.



2. Expose the object by carefully removing any clothing around it.
3. Check the circulation below the site of the injury.
4. Stabilize the object by using bulky dressings.



5. Use bandages to hold the dressings in place. Wrap the bandages snugly, but not so tightly as to cut off circulation.



6. Once the dressings are secure, recheck circulation below the site of the injury.
7. Make sure the person gets medical attention.



## Amputations

An amputation is a complete or partial severing of a body part. Although this causes a lot of damage to the surrounding tissues, bleeding may or may not be severe.

### Prevention

To reduce the risk of amputation:

- Keep hands clear of doors when they are closing.
- Use caution near train tracks: only cross at designated crossings, and never walk along the tracks.
- Be extremely careful when on or around any farm machinery, even when it is turned off or not operating.

### What to Do

If the body part is still partially connected to the body, put it back into place as best you can and treat the injury as an open wound or fracture.



#### Call

Call EMS/9-1-1.



#### Care

1. Try to control the bleeding with direct pressure. If you are not successful, or you are unable to apply direct pressure, apply a tourniquet.



2. Even if the body part has been fully amputated, there is still the possibility that it could be reattached as long as it is cared for properly. Remember to "protect it, bag it, cool it, tag it:"

**PROTECT IT:** Wrap the amputated part in gauze or a clean cloth.



**COOL IT:** Keep the amputated part cool by placing the bag on ice. Wrap the bag before cooling it so that the body part does not freeze.



**BAG IT:** Place the amputated part in a plastic bag.



**TAG IT:** Label the container with the person's name, the date, and the time.



3. Make sure the amputated part goes with the injured person to the hospital.



## Crush Injuries

### Common Causes

A crush injury occurs when there is a great deal of pressure on a part of the body, such as when a body part is squeezed between two heavy objects.

### What to Look For

If the person is still trapped under the object(s) or between two objects, it is likely that a crush injury has occurred. Other signs and symptoms of a crush injury include the following:

- Internal bleeding
- Shock
- Deformity
- Pain

### What to Do



#### Call

Call EMS/9-1-1 and get an AED.




#### Care

1. If the object is restricting the person's breathing, or the object is crushing the person's head, neck, chest, or abdomen, remove the object if it can be safely removed. If it is crushing another body part, leave the object where it is until EMS personnel arrive.


## Chest Injuries

Chest injuries can impact the ribs and/or the organs they protect. They can involve open wounds or internal injuries. If ribs are broken, they can penetrate the skin or the lungs, causing additional injury.


Injuries to the chest can result in conditions known as pneumothorax (air in the chest) or hemothorax (blood in the chest).



Pneumothorax is a condition in which air enters the chest cavity from the wound site but doesn't enter the lung. The air in the chest cavity presses against the lung, causing it to collapse.



Hemothorax is a condition in which blood accumulates in the chest cavity from the wound site but doesn't get into the lung. Because blood takes up space in the chest cavity, the lung can't expand effectively.



## Penetrating Chest Injuries

Penetrating chest injuries cause varying degrees of internal and external bleeding. Additionally, if the object penetrates the ribcage, air can pass freely in and out of the chest cavity and the person can no longer breathe normally. This can result in a breathing emergency.

### What to Look For

The following are signs and symptoms of a penetrating chest injury:

- Difficulty breathing
- Gasping
- Bleeding from an open chest wound that may bubble at the injury site
- A sucking sound coming from the wound with each breath
- Severe pain at the site of the injury
- Coughing up blood

### What to Do



#### Call

Call EMS/9-1-1 and get an AED.




#### Care

1. Have the person rest in a comfortable position. If the person wants to lie down, help the person into the recovery position with the injured side toward the ground.
2. If the wound is bleeding profusely, applying pressure or gauze dressings may be necessary to stop the bleeding. If bleeding is minor, do not apply pressure or a dressing.



To avoid the build-up of air between the chest wall and the lung (tension pneumothorax), air must be able to exit an open chest wound. If you must apply a dressing, ensure that it does not become saturated with blood, as this will prevent air from escaping. If the dressing becomes saturated, it must be changed.



## Blunt Chest Injuries

A blunt chest injury is caused by a crushing force that does not create a wound in the chest wall. The most common blunt chest injury is a broken rib. Incidents that may result in blunt chest injuries include motor vehicle collisions, falls, and sports injuries.

If multiple ribs break in multiple places, this can produce a loose section of the ribcage that does not move normally with the rest of the chest during

breathing. This is called flail chest. Normally, the loose section will move in the opposite direction to the rest of the chest. This is called paradoxical movement. Flail chest can also involve the sternum.

### What to Look For

The signs and symptom of a blunt chest injury include the following:

- Pain
- Deformity or swelling
- Guarded, shallow breathing
- Bruising at the site

### What to Do



Call

Call EMS/9-1-1.



Care

1. Keep the person as still as possible, as a blow to the chest may have caused head, neck, and/or spinal injuries.
2. Give the person something bulky (such as a towel) to hold against the chest, as this can make it easier to breathe.



## Abdominal Wounds

The abdomen is the area immediately under the chest and above the pelvis. It is easily injured because it is not surrounded by bones. The abdomen contains important organs such as the liver, which can be easily damaged. Injuries to the abdomen are more likely to produce life-threatening internal bleeding.

### What to Look For

The signs and symptoms of serious abdominal injuries include:

- Bruising
- Nausea and vomiting (sometimes vomit containing blood)
- Pain, tenderness, or a tight feeling in the abdomen
- Distension in the abdomen
- Organs possibly protruding from the abdomen

## What to Do



### Call

Call EMS/9-1-1 if you suspect life-threatening internal bleeding or if there are organs protruding from the abdomen.



### Care

If organs are not protruding, care for any injuries as you would care for them on any other part of the body.

If there are organs protruding:

1. Do not apply direct pressure or attempt to push organs back into the abdomen.
2. Cover the protruding organs with moist, sterile dressings.
3. Loosely cover the dressings with plastic, if available.
4. Cover the person with a towel or blanket to keep him or her warm



## Blast Injuries

Blast injuries occur when pressure waves generated by an explosion strike and pass through the body's surfaces. Blasts release large amounts of energy in the form of pressure and heat. Injuries can include loss of hearing, pneumothorax, internal bleeding, and organ damage. Thermal burns may also occur from exposure to heat. The extent of blast injuries may be difficult to identify because sometimes there

are no visible external injuries, and indicators of internal injuries may not be apparent. A person with suspected blast injuries should be monitored and reassessed frequently while waiting for EMS personnel to arrive.

There are four mechanisms of injury resulting from blasts:

1. Pressure or heat injuries from the blast itself.
2. Injuries from flying debris (shrapnel).
3. Trauma from being thrown by the blast.
4. Injuries or illnesses caused by hazardous material being dispersed by the blast.

## What to Do



### Call

Call EMS/9-1-1 and get an AED.



### Care

1. Keep the person still.
2. Treat any obvious external injuries.
3. Constantly monitor the person's condition as it may change rapidly.



# 9

# Head, Neck, and Spinal Injuries




Traumatic events such as falling from a height, getting hit by a motor vehicle, or sustaining a powerful blow to the head can cause head, neck, or spinal injuries. Injuries to the head or spine can damage bones, such as the skull or vertebrae, and tissues, such as the brain and the spinal cord. Head injuries are often accompanied by spinal injuries, but a person with a superficial head wound may also have an underlying head injury.

Depending on the nature and severity of the injury, the person may be left with permanent damage (e.g., paralysis, speech impairment, memory problems, or behavioural conditions). Although injuries to the head and spine account for a small percentage of all injuries, they cause more than half of the fatalities.

### The Neck and Spine

The spine is a strong, flexible column of small bones called vertebrae; it supports the head and the torso. The spine surrounds and protects the spinal cord, which is a bundle of nerves that runs between the brain and the lower back.



The spine is divided into four regions:

1. The cervical (neck) region, which consists of the C1 to C7 vertebrae.
2. The thoracic (chest) region, which consists of the T1 to T12 vertebrae.
3. The lumbar (lower back) region, which consists of the L1 to L5 vertebrae.
4. The sacrum (hip) and coccyx (tailbone) region, which consists of the S1 to S5 vertebrae and the four fused vertebrae that form the tailbone.

Severe injuries can move or break the vertebrae so that they squeeze or cut the spinal cord, causing temporary or permanent paralysis, or even death. Damage can also occur if the injury causes the soft tissue of the spinal cord to swell, compressing it against the hard bone that surrounds it.

Depending on the location and severity of the injury, the person may develop paralysis (the loss of movement, sensation or both) in body parts below the injury. Paraplegia is paralysis that affects both legs and the lower trunk. Quadriplegia is paralysis that affects both arms, the torso, and both legs. The higher up the spine the injury occurs, the more extensive the paralysis will be.

## The Brain

The brain is the centre of the nervous system. It is the master organ that controls all body functions. The brain has the consistency of gelatin, but it is cushioned from everyday bumps and jolts by the cerebrospinal fluid inside the skull. Because the skull is hard, there is not much room for the brain to move or swell.

Brain injuries can occur as a result of a blow to the head, a penetrating injury to the head (such as a bullet wound), or exposure to acceleration-deceleration forces that cause the head to snap forward and then back. A blow to the head can lead to a concussion (a traumatic brain injury that alters the way it functions), a brain contusion (bruising of the brain tissue) or a brain hematoma (bleeding into the space between the brain and the skull, which increases the pressure inside the skull and damages brain tissue).

### Common Causes

Many different situations can lead to head, neck, or spinal injuries:

- Motor vehicle collisions
- Recreation and sports injuries
- Falls
- Blow to the head or back
- Violent acts, such as assault

You should suspect a head, neck, and/or spinal injury in the following situations:

- A fall from any height greater than the height of the person
- Any diving injury (e.g., diving head-first into shallow water)
- A person found unresponsive for unknown reasons
- Any injury that involves a strong blow to the lower jaw, head, or torso (e.g., colliding with another participant while playing a sport)

- Any injury that causes a wound in the head or torso
- A motor vehicle collision or rollover
- Any injury in which a person's helmet is damaged
- A person has been struck by lightning
- A person has been electrocuted

It might be difficult to determine how much damage has taken place without advanced medical assessment and diagnosis, so always treat the injury as if it is serious. The size of a "goose egg" isn't necessarily an indication of the severity of a head injury. Watch for other signs of injury. You can put ice on a swollen area to reduce its size.



### Prevention

The following tips help to prevent head, neck, and spinal injuries:

- Wear safety belts with shoulder restraints when in a vehicle.
- Children and babies should always ride in approved safety seats. Make sure the seat is designed for the child's age and weight and make sure it is properly installed.
- Wear all recommended protective equipment for the activity you are engaged in (e.g., approved bicycle helmets).

- Never join in a new sport without knowing the rules and risks involved.
- Prevent falls around the home and workplace with non-slip floors, non-slip treads on stairs, handrails on staircases, rugs secured with double-sided adhesive tape, and handrails by the bathtub and toilet if necessary.
- Make sure that there is good lighting in stairways and hallways.
- If there are small children present, put gates at the top and bottom of the stairways.
- Make sure that your workplace is clean and tidy. Keep floors and aisles uncluttered and make sure that there is nothing blocking stairways, work sites, or exits.
- Drink responsibly. Alcohol is often a factor in serious motor vehicle collisions and water injuries. Alcohol slows down your reflexes and gives you a false feeling of confidence.
- Prescription drugs and common drugstore medications can make driving or operating machinery dangerous, so follow the directions on the package carefully.
- Check equipment (such as warehouse forklifts, ladders, and scaffolding) regularly for worn or loose parts.
- Use ladders carefully and correctly.
- Always be very careful around water:
  - Before diving, make sure that the water is deep enough, and check for objects below the surface, such as logs or pilings. Pools at homes, motels, or hotels may not be safe for diving.
  - Enter unknown water feet first.
  - Enter above-ground pools feet first.
  - When bodysurfing, keep your arms out in front of you to protect your head and neck.

## What to Look For

The signs and symptoms of a head, neck, or spinal injury depend on the nature and location of the injury. Bleeding inside the skull can happen slowly, so the signs may take time to appear. The severity of a head, neck, or spinal injury can only be determined through an evaluation by medical personnel: you should always assume that such an injury is serious.

The following signs and symptoms alone do not always indicate a serious head, neck, or spinal injury, but you should call EMS/9-1-1 if any of these are present:

- Changes in level of responsiveness, awareness, and behaviour
- Drowsiness, confusion, or disorientation
- Severe pain or pressure in the head, neck, or back
- Blood or other fluids in the ears or nose
- Heavy external bleeding from the head, neck, or back
- Unusual bumps, bruises, or depressions on the head, neck, or back
- Seizures
- Impaired breathing or vision
- Nausea or vomiting
- Unequal pupil size
- Persistent headache
- Partial or complete loss of movement of any body part without an obvious traumatic cause (e.g., inability to move a limb that does not appear to be injured)
- Back pain, weakness, tingling, or loss of sensation in the hands, fingers, feet, or toes
- Unusually positioned neck or back
- Dizziness and/or loss of balance
- Bruising of the head, especially around the eyes and behind the ears
- Loss of bladder or bowel control
- Behaviour similar to that of a person under the influence of alcohol or drugs (e.g., stumbling, memory loss, speech problems)

## What to Do

It is imperative that a person with a head, neck, or spinal injury remain as still as possible, unless keeping him or her in the position found would actually put the person's life at risk (if the scene is becoming unsafe, for example, or if you must roll the person over to put pressure on life-threatening bleeding). As always, an unresponsive person should be rolled into the recovery position. The risk of further damage to the spine is less serious than the risk of the person's airway being obstructed. You should take steps to discourage the person from moving reflexively. For example, if the person is lying on the ground, approach from a direction that allows the person to see you without turning his or her head. You should also tell the person to respond verbally to your questions rather than nodding or shaking his or her head.

If the person is unresponsive, you must check his or her ABCs, but this can often be done without moving the person. If you can see chest movement or can hear the person moaning, crying, or coughing, that means the person is breathing. Do not attempt to remove a person's helmet or remove a child from a car seat unless it prevents you from checking the ABCs or providing care for a life-threatening condition.



## Call

Call EMS/9-1-1 and get an AED.



## Care

1. Restrict movement of the head, neck, and spine as much as you can until EMS personnel arrive:
  - If the person is responsive, instruct him or her to keep as still as possible. If the person is unable to do so, manually support the head in a position that limits movement.



2. Keep the person in the position he or she was found, unless the person is in immediate danger or has life-threatening conditions that need immediate attention.
3. If blood or other fluid is in the ear canal or draining from the ear:
  - Let the ear drain. Do not apply direct pressure. Do not move the person unless it is absolutely necessary.
  - Cover the ear loosely with a sterile dressing.

## When to Move a Person with a Head, Neck, or Spinal Injury

You may need to move a person with a head, neck, and/or spinal injury in the following situations:

- The person's airway is blocked (for example, if the person vomits)
- The scene of the emergency becomes dangerous
- There is no organized EMS response in the region (for example, if the injury occurs in a very remote area)

When moving the person, try to keep the head, neck, and spine in the position you found them. Avoid unnecessary jostling and move the person only as much as is necessary.

Remember, it is more important to treat life-threatening conditions than it is to keep the person still. If the person has a potential spinal injury and is not breathing, opening the airway and starting CPR immediately are more important than protecting the spine from further damage.



### Concussion

A concussion is a subset of traumatic brain injuries (TBI) that involve a temporary alteration in brain function. An impact to the head or upper body can create forces that cause the brain to "shake" inside the skull. This shaking can cause the brain to collide with the bony structures that make up the skull's inner shell (sometimes causing swelling and/or bleeding), or to rotate within the skull, which can shear or tear the brain nerve fibres and sometimes stretch and damage the brain cells.

A concussion can result from even a seemingly minor bump, blow, or jolt, and may be difficult to recognize because the signs and symptoms may not be immediately obvious. Concussions are evolving injuries, with the effects intensifying, dissipating, or changing unexpectedly in the days and weeks following the event of the injury. Depending on the severity of the concussion, signs and symptoms can last for days, weeks, or even months. The majority of concussions, however, resolve in a short period of time.

It is not fully known exactly what happens to brain cells in a concussion, but a change in the brain's chemical function seems to be involved in the process. As a result, concussions are rarely visible in neuroimaging scans. The exact duration of this altered function is not yet known, but during the recovery period the brain is more susceptible to further injury. A person who has had one concussion is at increased risk for future concussions. As a result, it is critical that as soon as someone is suspected of having a concussion, they are removed from activity and seek care from a qualified medical professional. A person does not need to become unresponsive to have incurred a concussion.

### Common Causes

A concussion may be caused by:

- A blunt force to the head or jaw
- An explosion
- Whiplash
- A forceful blow to the body
- Shaking (in the case of a baby)

**What to Look For****SIGNS AND SYMPTOMS OF A CONCUSSION**

Thinking and Remembering	Physical	Emotional	Sleep
<ul style="list-style-type: none"><li>• Confusion</li><li>• Clouded or foggy mindset</li><li>• Seeming stunned or dazed</li><li>• Temporary memory loss regarding the event of the injury</li><li>• Difficulty concentrating</li><li>• Difficulty remembering or recalling events</li><li>• Slowed reaction times</li></ul>	<ul style="list-style-type: none"><li>• Neck pain, headache, or pressure within the head</li><li>• Fatigue, low energy, or feeling "foggy"</li><li>• Short-term loss of responsiveness</li><li>• Dizziness or loss of balance</li><li>• Double or blurred vision, or "seeing stars"</li><li>• Ringing in the ears</li><li>• Nausea or vomiting</li><li>• Mumbled or indistinct speech</li><li>• Sensitivity to light and/or noise</li><li>• Not feeling "right"</li><li>• Seizure or convulsion</li></ul>	<ul style="list-style-type: none"><li>• Irritability</li><li>• Sadness or depression</li><li>• Heightened emotions</li><li>• Nervousness or anxiety</li><li>• Personality changes</li></ul>	<ul style="list-style-type: none"><li>• Drowsiness</li><li>• Sleeping more or less than usual</li><li>• Difficulty sleeping</li></ul>

Concussions are often left untreated because the symptoms are not always recognized by the person who has been injured, and the signs are not always obvious to others. There may also be personal or external pressures to continue participating in the activity at hand (e.g., workplace requirements, social events, or sporting activities), instead of avoiding such aggravating factors. Continued participation in activities that provoke symptoms during either the acute or the recovery stage can aggravate the condition due to raised levels of exertion. To avoid this, it is important that the person seek medical attention following a blow to the head, neck, or upper body with any concussion symptoms. Cessation of physical activity and the promotion of cognitive rest (reducing "screen time" and other activities that require concentration) are critical to the resolution of concussion injuries.

### Concussion Signs and Symptoms in Children and Babies

Identifying signs and symptoms of a concussion in a child is more difficult than for an adult. It is particularly difficult for a child or baby, since the child or baby may not be able to speak to describe the symptoms. In addition to the standard signs of a concussion, the child or baby may exhibit the following signs, which are more specific to this age group:

- Disturbed sleeping and eating patterns
- Excessive crying
- Disinterest in activities or favourite toys

Children may also express simply "feeling off" or "not feeling right" following a physical impact. This is also a possible symptom of a concussion.

### What to Do



#### Call

Call EMS/9-1-1 if the person has the following more severe signs and symptoms of a concussion:

- Repeated or projectile vomiting
- Loss of responsiveness of any duration
- Lack of physical coordination (e.g., stumbling and unusual clumsiness)
- Confusion, disorientation, or memory loss
- Changes to normal speech
- Seizures
- Vision and ocular changes (e.g., double vision, dilated pupils, or unequal pupil size)
- Persistent dizziness or loss of balance
- Weakness or tingling in the arms or legs
- Severe or increasing headache



#### Care

If you think a person has sustained a concussion, advise the person to stop the activity he or she was engaged in when the incident occurred. If the person sustains an injury that causes a jolt or blow to the head or body, and then displays any signs and symptoms of a concussion, you should assume that a concussion has occurred. The person should follow up with a qualified healthcare provider as soon as possible for a full evaluation. A healthcare provider is best able to evaluate the severity of the injury and make recommendations about when the person can return to normal activities. While rare, permanent brain damage and death are potential consequences of failing to identify and respond to a concussion in a timely manner. Prompt recognition of when a potential concussion injury has occurred, immediate removal from the activity, and appropriate follow-up care are key elements in preventing catastrophic consequences.

Anyone who exhibits more concerning signs and symptoms requires immediate medical evaluation. An injury that causes a concussion may also trigger bleeding in or around the brain, leading to symptoms like extended drowsiness and confusion. This type of bleeding can be life-threatening. The person must be monitored in the immediate hours after the injury to ensure that the symptoms do not progress or worsen, as this can indicate additional brain injuries that require emergency care.

### Myth-Information

#### Myth: A person with a concussion who falls asleep could die.

It is generally considered safe for a person with a concussion to go to sleep. However, the person's healthcare provider may recommend that you wake the person periodically to make sure that his or her condition has not worsened.

### Follow-Up Care for Concussions

It is important that the person seek medical advice from a doctor, ideally one who specializes in concussions, since signs and symptoms can escalate in the days or weeks following the injury. A person who has suffered a concussion should not participate in any physical activities until a doctor gives permission to do so. Activities that require concentration (e.g., reading) and visual stimulation, such as working on a computer, texting, etc. should also be reduced to allow the brain the opportunity to recover from cognitive stressors. When given permission from a doctor, it is important to resume all of these activities gradually and be monitored by a medical professional.

A person recovering from a concussion may not be able to effectively monitor his or her own symptoms, and may not make the best decisions about things like rest, sleep, and limiting visual stimulation. It is crucial that the person's family members, caregivers, friends, or colleagues note any changes in his or her sleep habits or physical, cognitive, or emotional behaviour, as any of these can indicate that the recovery phase has

been compromised. These people can also have a positive impact on concussion recovery by supporting the person and encouraging him or her to follow the concussion recovery plan outlined by the doctor.

Proper management of the person's signs and symptoms by a qualified healthcare provider will help to avoid future problems that may lead to permanent issues and a diminished quality of life. If treated improperly, concussions can cause a wide range of functional long-term changes affecting thinking, sensation, language, and/or emotions. Concussions can also cause long-term neurological conditions, such as seizure disorders, and, in the most severe cases, death.

### Shaken Baby Syndrome

Shaken Baby Syndrome (SBS) refers to a variety of injuries that may result when a baby or a young child is violently shaken. This causes the brain to move within the skull, forcing blood vessels to stretch and tear. SBS is most associated with babies and toddlers, but can occur in children up to 5 years old.

SBS is the most common cause of mortality in babies. It is the most frequent cause of long-term disability in babies and young children, and it can cause permanent brain damage. SBS differs from other forms of physical abuse in that it is frequently a single event and there may be no obvious sign of injury.

Often there is no intent to harm the baby. Inconsolable crying is the most common trigger that leads a person to shake a baby. When trying to console a crying baby, never shake the baby, no matter what. Instead, place the baby face-up in a safe place and let the baby cry. Take a few deep breaths and then try again to soothe the baby. Other common triggers are feeding problems and difficulties with toilet training.

Feelings of anger and frustration can be overwhelming. However, shaking a child or baby is never appropriate. If you feel like you might lose control, stop! Take a break, never shake. Place the child in a safe

place and take a time-out for as long as it takes you to feel calm. Talk to someone you trust about your feelings. If you ever feel you may hurt a baby, call for help: a family member, neighbour, or local crisis line can provide emotional support and help you to find a solution.

### ***What to Look For***

While there may be no visible physical signs of injury, some signs of SBS include the following:

- Unexplained injuries (e.g., bruising or broken bones, especially skull, rib, and long-bone fractures)
- Bruising, usually in or around the eyes or mid-body area
- Bleeding or clear fluid coming from the ears and/or nose
- Minor neurological problems (e.g., irritability, lethargy, tremors, and vomiting)
- Major neurological problems (e.g., seizures, unresponsiveness, and death)

### ***What to Do***



#### **Call**

If you suspect a baby has SBS, call EMS/9-1-1.



#### **Care**

Treat any injuries you find. Avoid accusations and interrogation.



# 10 Bone, Muscle, and Joint Injuries

Bones are strong, hard, dense, tissues that come in many different sizes and shapes. More than 200 bones make up the skeleton, creating the framework for the body. Muscles are soft tissues that can contract and relax. Most muscles are skeletal muscles, which are attached to bones by tendons and move the body when they receive signals from the brain. A joint is formed where two or more bones are held together by ligaments that allow movement.

Bones, muscles, and joints have many jobs:

- Supporting the body
- Protecting internal organs
- Allowing movement
- Storing minerals
- Producing blood cells
- Producing heat through movement


There are four basic types of bone, muscle, and joint injuries:



**Sprain:** Occurs when a ligament is stretched, torn, or damaged. Ligaments connect bones together at the joints. Sprains most commonly affect the ankle, knee, wrist, and finger joints.



**Strain:** Occurs when a tendon or muscle is stretched, torn, or damaged. Tendons connect muscles to bones. Strains are often caused by overworking a muscle (e.g., by lifting an overly heavy object or by repetitively performing the same action). They usually involve the muscles in the neck, back, thigh, or the back of the lower leg.



**Dislocation:** Occurs when the bones that meet at a joint move out of their normal position. This type of injury is usually caused by a violent force that tears the ligaments, allowing the bones to move out of place.

**Fracture:** A complete break, chip, or crack in a bone. Fractures can be open (if the end of the broken bone breaks through the skin) or closed (the broken bone does not break through the skin). Open fractures are also called "compound fractures."

### Muscle Cramps

A muscle cramp is a painful condition that can be caused by heavy exercise or staying in the same position for too long. You can usually stop the pain by stretching and massaging the area with the cramp, resting, or changing position.

### Common Causes

Bone, muscle, and joint injuries can occur in many ways. The causes include the following:

- Falls
- Awkward or sudden movements
- Direct blows to the body
- Repetitive actions or forces, such as jogging

## Prevention

The following tips help prevent bone, muscle, and joint injuries:

- Always wear seat belts and shoulder restraints when in a vehicle, and ensure that all passengers do the same.
- Secure younger children in approved and properly installed child-restraint systems (e.g., car seats and booster seats).
- Wear all safety equipment that is required or recommended (e.g., helmets, goggles, and pads).
- Put non-slip adhesive strips or a mat in the bathtub and shower stall.
- Stretch before exercising.
- Avoid over-exertion and take breaks often. Children should learn their limits and rest when they are too tired or frustrated to continue safely.

Falls are the leading cause of injuries among the elderly. Reduce the risk with safety measures such as:

- Good lighting.
- Sturdy railings on staircases.
- Non-slip floors and rugs.

## What to Look For



The signs and symptoms of bone, muscle, and joint injuries may include the following:

- Pain, deformity, swelling, or bruising
- Limited or no use of the injured body part
- A broken bone or bone fragments sticking out of the skin
- A sensation or sound of bones grating
- Possible muscle cramps
- The sound of a snap or a pop when the injury happened

## What to Do

The person may be anxious and in a lot of pain. In most cases, the best thing that you can do is keep the person comfortable and prevent further injury until the person receives medical attention. To avoid damaging nerves, blood vessels, and tissues, do not perform traction or any other manipulation.

When caring for an open fracture, your first priority is to stop any bleeding. Provide care as you would for any other open wound.



Bone, muscle, and joint injuries range from very minor to life-threatening, so you must determine whether there is a need to call EMS/9-1-1. In general, if the injury seems severe or the person is in a significant amount of pain, you should call.

You should always call EMS/9-1-1 and get an AED if:

- There is a problem with the ABCs.
- The injury involves the head and/or spine.
- You suspect that there may be multiple injuries.
- There are injuries to the thigh bone or pelvis.
- The area below the injury is pale, blue, or cold to the touch.
- The area below the injury is numb.
- The person has an altered level of responsiveness.
- A broken bone is protruding through the skin.
- It is not possible to safely or comfortably move the person to a vehicle for transport to a healthcare facility.



## Care

Treat the injury using the RICE method:

**R REST** — Have the person stop any current activities and rest without moving or straightening the injured body part.

**I IMMOBILIZE** — If you need to move the person or if EMS response will be delayed, immobilize the injured area in the position in which it was found by creating a splint. Otherwise keep the person still and do not splint the injury.

**C COOL** — If it does not cause the person any discomfort, cool the injured area for 20 minutes of every hour to reduce swelling and pain. The part should be cooled for 20 minutes of every hour for up to 48 hours. Do not rub the ice or cold pack on the injured area. If you use ice, put some sort of thin, dry cloth or pad between it and the person's bare skin to avoid freezing the skin.

**E ELEVATE** — Keep the injured area above the level of the heart, if possible. However, do not raise the injured area if moving it will cause pain.

## Myth-Information

**Myth: You should apply heat to a muscle, bone, or joint injury to speed healing.**

Although heat is commonly used to relieve pain associated with chronic muscle, bone, and joint conditions such as arthritis, it is not the best treatment for an acute muscle, bone, or joint injury. Applying heat causes the blood vessels in the area to dilate (widen), bringing more blood to the area and increasing swelling. Cold, on the other hand, causes blood vessels to constrict (narrow), reducing blood flow to the area and helping to reduce swelling. In addition, applying cold slows the nerve impulses and helps to reduce pain.



## Splinting Guidelines

Splinting involves securing an injured bone or joint to keep it from moving, which reduces the risk of further injury and helps reduce pain. However, you should apply a splint only if you must move a person to get medical help or if the EMS response will be delayed.



Common items such as rolled newspapers, scarves, belts, and pillows can be used to improvise slings and splints if commercial versions are not available. Medical tape and medical braces can also be used to support strains and sprains.

There are four types of splint:



**Soft splints** are soft, bulky objects (e.g., a folded blanket, towel, pillow, or bandage).



**Rigid splints** are hard, fixed objects (e.g., a board, a rolled newspaper, a tree branch).




**Anatomical splints** use another body part for support (e.g., immobilize an injured leg by securing it to the uninjured leg).



**Slings** use cloth looped around the neck to support an upper extremity (e.g., the arm or wrist).

## Applying a Splint



No matter what type of splint you use, follow these steps:

- Check the temperature and colour of the skin below the injured area before and after splinting. The area should be warm, indicating good circulation.
  - If the area is cold before splinting, call EMS/9-1-1 if you have not already done so.
  - If the area is warm before splinting and cold afterwards, the splint may be too tight. Loosen it gently, and reassess to see if circulation has been restored.
- When possible, splint the injured part in the position in which it was found. Do not try to straighten or move the body part.
- Make sure the splint is long enough to extend above and below the injured area:
  - For bone injuries, immobilize the joint above and below the site of the injury.
  - For joint injuries, immobilize the bones above and below the site of the injury.
  - If you are not sure what is injured, include both the bones and the joints above and below the injured area in the splint.
- Always pad a rigid or anatomical splint to make the person more comfortable.
- Remove any jewellery (especially rings) that the person is wearing below the site of the injury, as swelling is likely to occur.

## Applying a Regular Sling for an Arm Injury

1. Check the person's circulation by comparing the warmth and colour of the fingers on the injured side with the other hand, and ask if the person has any numbness or tingling in the fingers.
2. Have the person support the injured arm, holding it across the body.



5. Tie the bandage toward one side of the neck, not at the back, to reduce discomfort from the knot.
6. Secure the elbow by twisting, tying, or pinning the corner of the bandage.
7. Apply a binder (as described below).



3. Slide the open triangular bandage between the injured arm and the body. The corner of the triangular bandage should extend past the elbow. The opposite side (the base of the triangle) should be running vertically, parallel to the edge of the body with the upper point over the shoulder.
4. Take the bottom end of the bandage and place it over the opposite shoulder.



8. Recheck circulation. Slings should fit tightly enough to restrict movement, but not be so tight that blood flow is affected. If fingers are bluish or cold or if the person feels numbness and tingling, loosen the bandages. If loosening the bandages does not improve circulation, call EMS/9-1-1 and get an AED immediately.




## **Applying a Tube Sling**

1. Check the person's circulation by comparing the warmth and colour of the fingers on the injured side with the other hand, and ask if the person has any numbness or tingling in the fingers.
2. Have the person hold the arm of the injured side across the body with the fingers touching the opposite shoulder.



5. Twist the end of the bandage hanging below the elbow several times to secure the arm, then bring it around and up the back to meet the other end of the bandage near the person's neck.
6. Adjust the height of the sling to make sure it is supporting the arm.
7. Secure the sling by tying the two ends together in the hollow of the neck on the uninjured side.



3. Place the open triangular bandage over the forearm and hand with the base parallel to the body and the opposite point extending past the elbow.
4. Support the forearm and gently tuck the lower edge of the bandage under the arm, from the hand to the elbow.



8. Pad between the arm and the body, in the natural hollow, with soft, firm material.
9. Apply a binder (as described on the following page).
10. Recheck circulation. Slings should fit tightly enough to restrict movement, but not be so tight that blood flow is affected. If fingers are bluish or cold or if the person feels numbness and tingling, loosen the bandages. If loosening the bandages does not improve circulation, call EMS/9-1-1 and get an AED immediately.



## Applying a Binder for an Arm Injury

A binder is used in addition to a sling and secures a slung upper extremity to the body. This provides additional support by creating anatomical support in addition to the sling.

1. Wrap a broad bandage around the injured arm and the body.
2. Tie the bandage snugly at the uninjured side.
3. Recheck circulation. The binder should fit tightly enough to restrict movement, but not be so tight that blood flow is affected. If fingers are bluish or cold or if the person feels numbness and tingling, loosen the bandages. If loosening the bandages does not improve circulation, call EMS/9-1-1 and get an AED immediately.

## Osteoporosis

Osteoporosis is a degenerative bone disorder that occurs when the amount of calcium in the bones decreases, causing low bone mass and the deterioration of bone tissue. Normally, bones are hard, dense tissues that can endure tremendous stress. Bone-building cells constantly repair damage that occurs as a result of everyday wear and tear, keeping bones strong. When the calcium content of bones decreases, the bones become frail, less dense, and less able to repair themselves after incurring stress and damage.

The loss of density and strength leaves bones more susceptible to fractures (especially of the hips, vertebrae, and wrists). Instead of being caused by tremendous force, fractures may now occur spontaneously, with little or no aggravation, trauma, or force. For example, the person may be taking a walk or washing dishes when the fracture occurs. Some hip fractures thought to be caused by falls are actually spontaneous fractures that cause the person's fall. Repeated fractures are also a sign of osteoporosis. Osteoporosis is a leading cause of bone and joint injuries in older people. It is much more common in women, affecting one in four.

You can help prevent osteoporosis by:

- Building strong bones at an early age.
- Making sure to get enough calcium.
- Making sure to get enough vitamin D (the body needs vitamin D so that it can absorb calcium).
- Exercising regularly.

A photograph of a young woman with long dark hair, wearing a grey hoodie, sleeping peacefully with her head resting on a wooden table. A golden-brown croissant sits on a small white plate in front of her. In the background, other people are seated at tables in what appears to be a cafeteria or breakroom setting.

# 11 Sudden Medical Emergencies

An acute illness is one that strikes suddenly and usually only lasts for a short period of time. A chronic illness is an illness that a person lives with on an ongoing basis and that often requires continuous treatment to manage. When a person becomes suddenly ill, it may be the result of an acute illness, or it may be an acute flare-up of a chronic condition.

Signs and symptoms such as difficulty breathing, pain that is persistent or severe, problems seeing or speaking, problems with sensation or movement, seizures, or unresponsiveness require a call to EMS/9-1-1. If you are unsure about the severity of the illness, it is better to call for help early than to wait for the situation to progress.

Fortunately, you do not need to know exactly what is wrong to provide appropriate first aid care in a sudden medical emergency. If your initial check of the person reveals any life-threatening conditions, make sure that someone calls EMS/9-1-1 right away, and then provide care according to the signs and symptoms that you find and your level of training. Follow the same general guidelines you would for any emergency.



## Fainting

Fainting is a brief period of unresponsiveness that happens when there is not enough blood flowing to the brain. If a person suddenly becomes unresponsive and then "comes to" after about a minute, he or she may have simply fainted.

## Common Causes

Fainting is caused by a sudden decrease in blood flow to the brain. Usually the cause of fainting is not serious. The following are common causes of fainting:

- Pregnancy
- Pain
- Heat
- Dehydration
- Decreased blood sugar (e.g., from missing a meal)
- Standing in one position for too long without moving
- Intense emotion
- Traumatic information (e.g., news of someone's death)

## Prevention

The following tips help prevent fainting:

- Keep hydrated.
- Eat at regular intervals to maintain a consistent blood sugar level.
- Get up slowly from sitting or laying down.

Fainting may be preceded by paleness, dizziness, sweating, or nausea. If you think that someone is about to faint, have the person sit or lie down.

## What to Do

### Call

Call EMS/9-1-1 and get an AED if the person is unresponsive for more than a few minutes, the person is pregnant, the person's medical history is unknown, or you suspect that the person fainted as a result of a serious injury or illness. If you are unsure, call EMS/9-1-1 and get an AED.



## Care

1. Place the person in the recovery position.
2. Encourage the person to follow up with his or her healthcare provider.



### Diabetic Emergencies

Diabetes is a chronic condition characterized by the body's inability to process glucose (sugar) in the bloodstream. An organ called the pancreas secretes insulin, a hormone that causes glucose to move from the bloodstream into the cells, where it is used for energy. In a person who has diabetes, either the pancreas fails to make enough insulin or the body's cells are unable to respond to insulin. Either situation causes glucose levels in the bloodstream to increase.

A person with diabetes may manage the condition with insulin injections or oral medication. Diet and exercise also play an important role. A person with diabetes must follow a well-balanced diet, with limited sweets and fats. The timing of meals and snacks relative to

exercise and medication is important as well. People with diabetes may need to check blood sugar levels more frequently when daily routines are disrupted—for example, by illnesses or vacations—as this can cause blood sugar levels to fluctuate more than usual.

If food intake, exercise, and medication are not in balance, the person may experience a diabetic emergency. A diabetic emergency happens when blood glucose level fluctuates outside the normal range, resulting in either hyperglycemia (too much sugar in the blood) or hypoglycemia (too little sugar in the blood).

**Hyperglycemia** is a condition in which lowered insulin levels result in elevated blood sugar. Hyperglycemia can result if a person eats too much food, takes too little medication, exercises less than usual, or experiences physical or emotional stress. Hyperglycemia develops slowly, so it is less likely to be a first aid emergency.

With hyperglycemia, the body cells cannot get the sugar they need, even when there is a lot of sugar in the body. To meet its energy needs, the body breaks down other food sources, causing waste products to build up in the body and thereby making the person ill. This may also cause the person's breath to have a fruity or sweet odour. This can lead to a serious form of diabetic emergency called a diabetic coma.

**Hypoglycemia** occurs when the insulin level is too high relative to the level of sugar in the blood. The small amount of sugar in the blood gets used up quickly. Hypoglycemia can result if a person misses a meal or snack, exercises more than usual, vomits, or takes too much medication. With hypoglycemia, the brain does not get enough sugar to work properly; the result is an acute condition called insulin shock. Hypoglycemia develops quickly, and is the cause of most diabetic emergencies.

## Common Causes

A diabetic emergency occurs when there is an imbalance between two or more of the following:

- Exercise
- Food intake
- Medication
- Insulin production



Not all people experiencing a blood sugar emergency have diabetes.

## What to Look For

The following are signs and symptoms of a diabetic emergency:

- Changes in the level of responsiveness
- Changes in behaviour, such as confusion, irritability, or aggression
- Rapid breathing
- Cool, sweaty skin
- Skin that is paler than normal
- Appearance of intoxication (e.g., slurred speech, difficulty walking)
- Feeling and looking ill
- Seizures



First aid is the same for both types of diabetic emergencies (hyperglycemia and hypoglycemia).

## What to Do



### Call

Call EMS/9-1-1 if:

- You are unable to give the person sugar.
- The person is not fully awake.
- The person has a seizure.
- The person's condition does not improve within 10 minutes of having sugar.



### Care

If the person is unable to follow simple commands, unable to swallow safely, or is unresponsive, ensure that EMS/9-1-1 has been called and place the person in the recovery position.

1. If the person is able to follow simple commands, answer questions, and swallow safely, offer 15 to 20 grams of sugar. Check the label on packaged products to determine how much of the package's contents to give. The following are the forms of sugar to give, listed in order of preference:

- Oral glucose tablets
- Chewable candy
- Fruit juice
- Fruit strips
- Milk
- Other forms of sugar, such as a non-diet soft drink or a spoonful of sugar mixed into a glass of water



2. Monitor the person's condition for 10 minutes:

- If the person's condition deteriorates or does not improve within 10 minutes, call EMS/9-1-1 and administer more sugar if it is still safe to do so.
- If the person's condition improves, recommend that he or she eat a complete meal.


If the person's condition is caused by low blood sugar, ingesting sugar will help improve his or her condition. Even if the person is experiencing hyperglycemia (too much glucose in the bloodstream), giving the person 15 to 20 grams of sugar will not cause additional harm. If a glucometer is available, having the person check his or her blood sugar will reveal whether he or she has hyper- or hypoglycemia.



Do not give the person insulin. Only a medical professional can diagnose the cause of the emergency and be certain that insulin is the correct treatment.

### ***Using a Glucagon Kit for a Diabetic Emergency***

Some people with diabetes may carry a prescribed glucagon kit to use in case of a severe hypoglycemic emergency. Glucagon is a hormone that stimulates the liver to release glucose into the bloodstream. The glucagon kit is used only when the person is unresponsive or has lost the ability to swallow. Those who spend a significant amount of time with the person (e.g., family members, teachers, coaches, or co-workers) may receive additional training to learn how to administer a glucagon injection. If you have not received specific training in the use of a glucagon kit, you should never attempt to administer this medication.



### **Seizures**

A seizure is an episode of abnormal electrical signals in the brain that result in temporary and involuntary disturbances in brain function, shaking or contraction of limbs (convulsions), changes in sensation, shifts in behaviour, and altered levels of responsiveness.

## Common Causes

Seizures can have many different causes. One common cause is epilepsy, a chronic seizure disorder that can often be controlled with medication. Other causes of seizures include the following:

- Fevers or infections
- Head injuries
- Heat stroke
- Poisons (including drugs)
- Drug or alcohol withdrawal
- Diabetic emergencies
- Audio-visual stimulation (e.g., flashing lights)

## What to Look For

The following are common signs and symptoms of seizures:

- Hallucinations
- Uncontrollable muscle movement
- Eyes rolling upward into the head
- Drool or foaming at the mouth
- Uncontrolled repetitive motions (partial seizures)



Any one of these signs can indicate a seizure—not all seizures involve convulsions.

A person may experience an aura (an unusual sensation or feeling) before the onset of the seizure. If the person recognizes the aura, he or she may have time to tell someone what is happening and sit down before the seizure occurs.

There are many different types of seizures, and their specific signs and symptoms vary. Common types of seizures include:

**Tonic-Clonic or Convulsive Seizures:** Formerly called "Grand Mal" seizures. The person becomes unresponsive and has convulsions. During the tonic phase, muscles will stiffen. Then, during the clonic phase, the extremities will jerk and twitch rhythmically. Saliva that has not been swallowed during the seizure may appear as froth at the mouth. Breathing may be irregular as the respiratory muscles can be affected. A tonic-clonic seizure usually lasts one to three minutes, but may last up to five minutes. The period after the seizure is called the post-ictal phase. During this time,

the body rests to recover from the seizure. The person will slowly regain responsiveness and may appear fatigued, confused, or disoriented.

**Absence Seizures:** Formerly called "Petit Mal" seizures. The person experiences a brief, sudden lapse in responsiveness. He or she may momentarily become very quiet and have a blank stare, or appear to be daydreaming. The person may also make chewing motions, breathe rapidly, blink rhythmically, or make slight movements such as tugging at clothing. There are no convulsions with absence seizures. Absence seizures are very brief, usually lasting between two and ten seconds. There is no confusion after the seizure, and the person can usually resume full activity immediately.

**Focal or Partial Seizures:** These seizures begin in one specific part of the brain. The signs and symptoms of the seizure vary depending on which region of the brain is affected. Typically a focal or partial seizure will affect only one part of the body (e.g., one arm may convulse) or aspect of experience (e.g., a person's vision could be disturbed, or a person could experience a sudden change in his or her emotional state). The person may or may not remain aware during the seizure. Some focal or partial seizures can evolve into a generalized tonic-clonic or convulsive seizure.

## What to Do



### Call

Call EMS/9-1-1 and get an AED if:

- You do not know the person or the person's medical history.
- The seizure lasts more than a few minutes.
- The person has several seizures in a row.
- The person appears to be injured.
- The person is pregnant.
- The person is experiencing a diabetic emergency.
- The seizure takes place in water.
- This is the person's first seizure, or the cause of the seizure is unknown.
- The person does not wake up after the seizure or is unresponsive for an extended period.

## Calling EMS/9-1-1 for People with Seizure Disorders

Some people have multiple seizures during a day, and will likely have a seizure treatment plan in place. The treatment plan will outline when a seizure no longer follows the normal pattern or duration for the person and when to call EMS/9-1-1. Follow the person's seizure plan if you are familiar and comfortable with it.

## Care

### During the seizure:



#### 1. Protect the person from injury by:

- Moving furniture and other objects that could cause injury out of the way.
  - Protecting the person's head with a soft object (such as a blanket).
2. Do not try to hold the person down or stop the seizure from happening.
3. Roll the person onto his or her side, if you are able to do so safely



### After the seizure:

The person may be drowsy and disoriented for up to 20 minutes.

1. Check the person's ABCs.
2. Place the person in the recovery position.



Although a seizure can be frightening to see, it is easy to care for a person who is having a seizure. Most seizures only last a few minutes, and the person usually recovers fully without any complications.

## Myth-Information

**Myth:** You should put something between the teeth of a person who is having a seizure to prevent the person from biting or swallowing his or her tongue.

This practice is unsafe and unnecessary. It is impossible to swallow one's own tongue. Although the person may bite down on the tongue, causing it to bleed, this is a minor problem compared to the risks of attempting to put an object in the person's mouth. The person could chip a tooth or knock a tooth loose, creating a choking hazard. The person may also bite down with enough force to break the object and then choke on one of the pieces. Additionally, attempting to place an object in the person's mouth puts you at risk of being bitten.

## Febrile Seizures

Babies and young children may have seizures if their body temperatures suddenly rise. These are called "febrile seizure" and are most commonly associated with sudden high fevers, normally with temperatures over 39°C (102°F). In most cases, these seizures are non-life-threatening and do not last long.

To reduce the risk of febrile seizures in a child or baby with a high fever, you must lower his or her body temperature:

1. Remove any excess clothing or blankets.
2. Give the child or baby a sponge bath with water that is room temperature (not icy cold).
3. Give the child or baby plenty of fluids to drink to help prevent dehydration.
4. Consider giving the child or baby fever-reducing medication such as acetaminophen or ibuprofen.

If these steps do not reduce the child's or baby's temperature, seek medical attention.



### Call

Call EMS/9-1-1 for a febrile seizure if:

- It is the first time that the child or baby has had a febrile seizure.
- The seizure lasts longer than 5 minutes or is repeated.
- The seizure is followed by a quick increase in body temperature.



## Mental Health Crisis

Mental health first aid is the initial care given by a First Aider to a person in a mental health crisis. Like any first aid, it includes the recognition of signs and symptoms and the care that can be given until either appropriate professional follow-up care is received or the crisis is resolved.

A First Aider is not a therapist, but rather a listener. Your role is to look, listen, and feel empathy for the person's situation and provide both physical and mental health first aid as necessary. You may also need to refer the person to the appropriate follow-up care, just as you might when caring for physical conditions.

## Mental Health Conditions

Mental health conditions are difficult to identify. They may go unrecognized unless the person shares the information with you. People in distress can be suffering from a number of disorders including substance-related disorders, mood disorders, anxiety disorders or, less commonly, psychotic disorders.

A person with a mental health condition may experience one, many, or none of the following symptoms:

- Inability to think clearly
- Inability to concentrate or focus on a task
- Hallucinations (e.g., hearing voices, seeing, or feeling things that aren't there)
- Delusions
- Depression or acute mood swings (e.g., from happy to depressed with no clear reason for the change)
- Poor memory
- Obvious lack of motivation

## What to Do

Helping a person through a mental health crisis can be achieved by using active listening practices:

- Reduce distractions and encourage the person to sit down to promote conversation.
- Keep the person as calm as possible.
- Listen empathetically.
- Acknowledge the person's feelings and emotions without judgment.
- If the person is delusional, do not dismiss his or her beliefs: accept that they are real to the person (but not to you).
- Speak quietly and firmly.



### Call

Call EMS/9-1-1 immediately if you suspect that the person poses a risk to you or others, or that the person is contemplating suicide or other self-harm. If you are aware of support systems that are in place for the person, contact them as well.



### Care

1. Use active listening practices to provide reassurance, comfort, and support.
2. Offer self-help strategies such as breathing exercises.



## Self-Inflicted Injuries

Self-harm ranges from minor self-inflicted injuries to suicide. Suicide is one of the ten most common causes of death in Canada, affecting people from a variety of backgrounds and age groups. Anyone expressing intentions of self-harm requires immediate intervention, and therefore EMS/9-1-1 must always be called.

Possible indicators of self-harm include the following:

- A person expressing negative (especially suicidal) thoughts and comments about him-or herself
- A person expressing his or her intent to die, especially if he or she has a plan to do so
- A person expressing his or her final wishes to someone close by (e.g., a bystander at scene)
- Items associated with a suicide attempt found at the scene (e.g., a gun, sleeping pills)



## Childbirth

You may find yourself in the position of helping a pregnant woman who is in labour. Childbirth is a natural process and the woman's body knows what to do: Your primary role will be to provide comfort and reassurance and to facilitate the process while you wait for EMS personnel to arrive. The labour and delivery process will happen without much intervention on your part. Remember, the woman who is pregnant delivers the baby. Be patient and let the birth happen naturally.

## What to Look For

Signs and symptoms that signal imminent childbirth include the following:

- Contractions that are 2 minutes apart or less
- The woman says that the baby is coming
- The woman feels the urge to push
- The woman feels like she needs to have a bowel movement
- The baby is crowning

## What to Do



### Call

Have someone call EMS/9-1-1 if birth is imminent.



### Care

1. Clear the area of unnecessary bystanders.
2. Help the woman into a position of comfort.
3. Create a clean birthing area by placing clean blankets and/or towels under the woman. For the sake of privacy, place a clean sheet or towel over the woman's abdomen.

4. As the baby is being birthed:
  - Support the head as it is delivered. Do not push or pull the baby.
  - Once the shoulders emerge, the rest of the baby will be delivered very quickly.
  - Newborns are slippery, so hold the baby firmly but do not squeeze him or her.



5. Position the baby face down and wipe any fluids or mucus away from his or her mouth and nose.
6. Check the baby's airway and breathing. Babies may not breathe and cry immediately after they are born. Usually, actively drying them with a towel will stimulate them to breathe and cry. If the baby does not begin to breathe or cry following stimulation, begin CPR.
7. Keep the baby warm by placing him or her directly onto the mother's chest and covering with a blanket or towel, or by wrapping the baby in a clean towel or blanket.
8. Let the placenta and cord drop onto a clean towel and keep this near the baby. Do not cut the cord. Handle the placenta and cord as little as possible.
9. Record the time of birth.
10. If the mother is bleeding, you may need to apply gentle pressure to any bleeding tears.
11. Provide continual care for both the mother and the baby until EMS personnel arrive.

## Assisting with Childbirth: What Not to Do

When providing care for a woman during childbirth:

- Do not let the woman leave to use the restroom. (The woman could deliver the baby into the toilet, putting the baby at risk for injury.)
- Do not try to physically delay delivery, as this can cause serious injuries to both the mother and the baby.
- Do not place your fingers in the woman's vagina for any reason. This can introduce pathogens that can cause an infection.
- Do not pull on the baby.

## Miscarriage

Miscarriage is the spontaneous end of a pregnancy any time during the first 20 weeks after conception. This occurs in about 1 in 10 pregnancies. The risk of miscarriage drops as the pregnancy progresses.

## Common Causes

There are a number of causes of miscarriage, including the following:

- Hormonal or genetic complications
- Abnormalities in the womb
- Infection and certain illnesses
- Trauma

## What to Look For

The signs and symptoms of a miscarriage include the following:

- Anxiety and apprehensiveness
- Vaginal bleeding, which may be minor or profuse
- Cramp-like pain that is similar to labour or menstruation

## What to Do



### Call

Have someone call EMS/9-1-1.



### Care

1. If possible, save any tissue from the miscarriage for EMS personnel.
2. Provide comfort and continual care until EMS personnel arrive.



# 12 Environmental Illness

## How Body Temperature Is Controlled

The human body's core temperature (i.e., the temperature of the heart, lungs, and brain) is normally around 37°C (98.6°F) and is maintained by balancing heat loss with heat gain.

The body generates heat by either converting food into energy or by contracting muscles (e.g., shivering, exercising). The heat produced by routine activities such as walking is usually enough to balance normal heat loss.

## The Hypothalamus and Body Temperature

The hypothalamus is the part of the brain that controls thermoregulation, which is the body's ability to maintain a temperature that is within the ideal range. The hypothalamus gets temperature information from the skin and central receptors. If the body is too warm, thermoregulatory "heat loss" responses include increasing the skin's blood flow and/or sweating to facilitate evaporation. If the body is too cold, responses include decreasing the skin's blood flow and shivering, which produces heat by contracting and relaxing muscles.


## HUMAN THERMOREGULATION RESPONSES

### Warm / Hot Environment

- Heat Loss (vasodilation)
- Heat Loss (sweating)

### Cool / Cold Environment

- Heat Conserving (vasoconstriction)
- Heat Production (shivering)




## The Four Mechanisms of Heating and Cooling the Body

Heat moves from warmer areas to cooler ones through four mechanisms:

1. **Radiation:** (Warming or cooling) Involves the direct loss or absorption of heat energy through electromagnetic waves.
2. **Conduction:** (Warming or cooling) Occurs through direct contact with a solid or liquid. Heat loss due to direct contact with cold snow could be decreased by minimizing contact or increasing insulation (e.g., sitting on an insulated pad on the snow).
3. **Convection:** (Warming or cooling) Occurs when air or liquid moves across the skin. In cold air, this is known as the wind chill factor. Convective heating can also occur if warm air is blown across the skin.
4. **Evaporation:** (Cooling only) Occurs when a liquid changes to a vapour. This physical process requires heat, which means that evaporation from the skin takes heat from the surface of the body. When sweat or water (either on the skin or in clothing that is in contact with the skin) evaporates, the skin is cooled. Sweating occurs when either the skin and/or body core temperatures increase. In a cold environment, sweating should be minimized (e.g., layers of clothing should be removed before or during hard work). In a hot environment, evaporative heat loss is beneficial. However, any sweat that does not evaporate will not cause cooling. For example, when humidity is high, sweat does not evaporate as efficiently and may simply drip off the body.

## FOUR MECHANISMS OF HEAT LOSS

- Conduction: Through direct contact with a solid or liquid.
- ~~~ Convection: Flowing water or air removes heat.
- - - Radiation: Through space (air) by infrared radiation.
- • • Evaporation: Change from liquid to gas requires energy.



## Heat-Related Illnesses

Hyperthermia occurs when the body's core temperature rises above the normal range. It results when at least one of the four mechanisms that cool the body is impaired by clothing, drugs, or disease, or is overwhelmed by internal heat production and/or external heat exposure. Heat-related illnesses will get worse without treatment and can change from one level to another very quickly.

### Prevention

One of the most important actions a person can take to prevent heat-related illnesses is to drink plenty of fluids. You should drink eight 250 mL (8 oz.) glasses of fluid a day for low-exertion, routine activities and more as exertion levels increase. It is important to rehydrate regularly during longer events. If you feel thirsty, you are already dehydrated, so develop the habit of drinking fluids regularly rather than waiting for the warning signs of dehydration to appear.

The following tips will also help to prevent heat-related illness:

- Maintain a healthy level of cardiovascular fitness.
- Prior to a major event in the heat, let the body gradually acclimatize by being exposed to 1 to 2 hours of heat exertion for at least 8 days.
- Avoid being outdoors during the hottest part of the day.
- Reduce the intensity of activities as it gets hotter and don't work or exercise for too long at a time.
- Take frequent breaks in a cool or shaded area to cool off. This will help the body cope with short periods of extreme heat.
- Dress for the heat and for the intended activity.
- Make adjustments to exertion levels and work/rest cycles on hot days (e.g., schedule football practices for cooler parts of the day).
- Wear a light hat when under the sun. Wear loose-fitting light clothing that allows liquid and vapour to escape.
- Choose hydrating drinks, such as water or juice, over dehydrating ones like alcohol.

## COMMON CAUSES OF ENVIRONMENTAL ILLNESS

Environmental	Physical	Behavioural
<ul style="list-style-type: none"> <li>• Heat waves, especially if there hasn't been one in recent years</li> <li>• High humidity (above 75%), which decreases the ability of sweat to evaporate</li> </ul>	<ul style="list-style-type: none"> <li>• Age: Babies, children, and the elderly are less able to sweat and adjust to changes in temperature</li> <li>• Body size and mass: Larger bodies with more fat tissue retain heat and warm up more quickly</li> <li>• Clothing that prevents the loss of heat and moisture (e.g., football padding)</li> <li>• Chronic illness</li> <li>• Heart disease</li> <li>• Skin, hormone, or nervous system diseases</li> <li>• Burns</li> <li>• Poor physical fitness</li> </ul>	<ul style="list-style-type: none"> <li>• Working or exercising too much in hot weather</li> <li>• Not drinking enough fluids to replace the water lost by sweating (dehydration)</li> <li>• Drinking too much alcohol in hot weather</li> <li>• Taking stimulants such as cocaine or amphetamines</li> <li>• Salt depletion</li> <li>• Fatigue</li> </ul>



## Heat Cramps

Heat cramps are painful muscle spasms, usually in the legs and abdomen, caused by loss of fluids and electrolytes as a result of sweating. While they are usually not serious, they are often the first sign that the body is beginning to overheat.

### What to Look For

The following signs and symptoms can indicate heat cramps:

- Mild muscle contractions that can become severe, usually in the legs and abdomen
- Moist skin

A person with heat cramps will typically have a normal or slightly elevated core temperature (37°C or 98.6°F).

### What to Do



#### Care

1. Reduce heat exposure (e.g., get into the shade) and, if possible, move the person to a cooler environment.
2. Have the person slowly sip a cool electrolyte-replacement beverage (e.g., commercial sports drink, coconut water, fruit juice, or milk). If a drink containing electrolytes is not available, have the person drink water.
3. Gently stretch and massage the cramped muscles.

When the cramps stop, the person can usually resume his or her activity as long as there are no other signs or symptoms of illness.

## Myth-Information

**Myth: When a person has heat cramps, you should give salt tablets to replenish lost sodium.**

Salt tablets are not an effective treatment for heat cramps. Consuming a concentrated form of salt can actually promote loss of fluid from the body, which will make the person's condition worse instead of better.



## Heat Exhaustion

Heat exhaustion occurs when the body begins to overheat, especially when it loses more fluids and electrolytes through sweating than it gains through rehydration. It usually happens after long periods of strenuous activity in a

hot environment, especially if it is humid or lacking in air circulation, as these conditions make it difficult for sweat to evaporate.

Heat exhaustion is an early indicator that the body's temperature-regulating mechanisms are being overwhelmed. As more blood flows to the skin to remove heat from the body's core, not enough blood flows to the vital organs. This causes a mild form of shock. Heat exhaustion can progress to heat stroke, so monitor the person's condition carefully.

## What to Look For

The following signs and symptoms can indicate heat exhaustion:

- Normal or slightly raised core temperature (37 to 39°C, or 98.6 to 102.2°F)
- Moist skin that is flushed (red) immediately after exertion, and then turns pale or ashen (grey)
- Dehydration or intense thirst
- Headache, nausea, dizziness or fainting
- Weakness or exhaustion

## What to Do



### Call

Call EMS/9-1-1 if the person is vomiting, has an altered level of responsiveness, or is unable to drink fluids.



### Care

1. Reduce heat exposure (e.g., get into the shade) and if possible, move the person to a cooler environment.
2. Loosen any tight clothing and remove any padded clothing or equipment.
3. Pour cool water on the person's clothing and/or on towels or cloths and place them on the person's chest. Fan the person to increase evaporation.
4. Apply ice or cold packs to the armpits and chest.
5. If the person is responsive and able to swallow, have him or her slowly sip a cool electrolyte-replacement beverage (e.g., commercial sports drink, coconut water, fruit juice, or milk). If a drink with electrolytes is not available, have the person drink water.
6. Advise the person not to do any more activities in the heat that day.



## Heat Stroke

Heat stroke is the least common but most severe heat-related illness. It is a life-threatening emergency that occurs when the body's cooling system is completely overwhelmed and stops working.

The body stops sweating because the levels of fluid are too low. When sweating stops, the body's temperature rises quickly. When it gets too high, the brain and other vital organs cannot work properly. The person becomes unresponsive, has seizures, and can die without immediate first aid.

### What to Look For

- High core temperature, above 40°C (104°F)
- Hot, dry skin
- Flushed (red) or pale skin
- Headache
- Altered mental status (e.g., confusion)
- Irritable, bizarre, or aggressive behaviour
- Progressive loss of responsiveness
- Rapid, weak pulse that becomes irregular
- Rapid, shallow breathing
- Vision problems
- Seizures or coma

### What to Do

#### Call

Call EMS/9-1-1 and get an AED.

#### Care

1. Move the person to a cooler environment.
2. Quickly lower the person's core temperature with one of the following methods:
  - Immerse the person in cool water from the neck down. Do not remove the person's clothing.
  - Immerse the person's forearms and hands in cool water.
  - Pour cool water on the person's clothing and/or on towels or cloths and place them on the person's chest. Fan the person to increase evaporation.
  - Apply ice or cold packs to the person's armpits and chest.
3. Continue to use the rapid cooling methods listed above until either the person's condition improves or EMS personnel arrive.
4. If the person is responsive and able to swallow, have him or her slowly sip a cool electrolyte-replacement beverage (e.g., commercial sports drink, coconut water, fruit juice, or milk). If a drink with electrolytes is not available, give the person water.

ASSESSMENT OF A HEAT-STRESSED PERSON					
	SKIN	PHYSICAL	MENTAL	PULSE	BREATHING
HEAT CRAMPS	MOIST WARM	MUSCLE CONTRACTIONS (MILD TO SEVERE)	NORMAL	NORMAL	NORMAL
HEAT EXHAUSTION	MOIST WARM	HEADACHE WEAKNESS/EXHAUSTION NAUSEA, VOMITING FAINTING	ANXIETY DIZZINESS	NORMAL	NORMAL
HEAT STROKE	DRY HOT	SEIZURES COMA SEVERE HEADACHE	ALTERED BEHAVIOUR: IRRITABLE AGGRESSIVE BIZARRE	RAPID WEAK	RAPID SHALLOW

CAUTION    CALL EMS/9-1-1

**Heat Cramps**

Remove from heat



Loosen tight clothing, remove padding from torso



Gentle stretching



If person is alert, provide cool drink

**Heat Exhaustion**

Remove from heat



Loosen tight clothing, remove padding from torso

**ACTIVE COOLING**

Pour water on torso



Fan skin



If person is alert, provide cool drink

**Heat Stroke**

Remove from heat



Loosen tight clothing, remove padding from torso

**AGGRESSIVE COOLING  
(ORDER OF PREFERENCE)**

Immerse body in cool water



Immerse forearms in cool water



Pour water on torso



Fan skin



If person is alert, provide cool drink

## Cold-Related Illness

Cold-related illnesses are any conditions that are caused by exposure to colder temperatures. They range in severity from superficial frostbite to life-threatening hypothermia.



### Frostbite

Frostbite is a local, superficial injury caused by freezing of the skin and, in more extreme cases, the underlying tissues. Extremities, such as the toes, feet, fingers, hands, ears, and nose are particularly prone to frostbite.

There are two levels of frostbite: **Superficial frostbite** occurs when the skin freezes but the underlying tissues are unaffected. It may produce clear blisters after the tissue thaws, but little or no tissue loss typically occurs. **Deep frostbite** occurs when tissues beneath the skin (including tendons, muscles, and blood vessels) freeze. Inside the body, ice crystals and swelling begin to damage or destroy the body's cells, blood vessels, and nerves. Deep frostbite may produce dark, hemorrhagic blisters when the tissue thaws, and is more likely to result in tissue loss.

Before frostbite occurs, a person's skin may appear shiny and rosy (a condition sometimes called "frost nip"). This is a warning that frostbite is imminent. If you see these signs, advise the person to move to a warmer environment or protect his or her skin with layers of clothing.

### Common Causes

Factors that increase the risk of frostbite include the following:

- Moisture on the skin
- Exposure to wind
- Insufficient insulation
- Clothing or equipment that reduces or constricts blood flow

### Prevention

The following tips will help to prevent frostbite:

- Never ignore numbness: If an extremity feels numb or tingly, you must take steps to warm it immediately (e.g., put your hands under your armpits or pull your arms inside your jacket for direct skin-to-skin contact).
- Cover up vulnerable areas such as the cheeks, nose, and ears by wearing clothing that is appropriate for the weather (e.g., scarf, toque, mittens).
- Wear mittens instead of gloves when possible, as mittens provide better insulation.
- Maximize foot insulation but avoid a tight fit around the toes.
- Wear clothing in layers and adjust as necessary so that you feel warm but are not overheating and sweating.
- Keep well hydrated.
- Keep clothing dry and change out of wet clothing as soon as possible.

### What to Look For

#### Superficial Frostbite

Signs and symptoms of superficial frostbite include:

- Hardened skin
- Skin that looks paler than the area around it
- Pain or stinging in the area, followed by numbness

#### Deep Frostbite

Signs and symptoms of deep frostbite include:

- Waxy skin that is colder than the area around it
- Skin and underlying tissue that is hard and solid to the touch
- Skin that is white, blue, black, or mottled
- Complete loss of feeling

## What to Do

Because both hypothermia and frostbite are caused by exposure to the cold, you may be providing care for someone with signs and symptoms of both conditions. Because hypothermia is more serious, however, you should care for it first before caring for the frostbite.



### Care

1. If possible, remove jewellery or other extraneous material that may restrict blood flow to the affected area.
2. Thaw the area only if you are sure it will not freeze again.
  
3. Warm the affected area using warm water, if available (approximately 38 to 40°C (100.4 to 104°F)) or body heat (e.g., by placing the person's hands in his or her armpits or on the abdomen). You should be able to place your own hands in the water for a minute without feeling too warm.



4. If the frozen area has thawed, don't break any blisters. Protect them with loose, dry dressings. Place gauze between the fingers or toes if they are affected.



5. If possible, elevate any thawed extremities above the level of the heart.
6. Because thawing frostbite can be extremely painful, the person may wish to take an over-the-counter painkiller.
7. Rehydrate the person by providing plenty of fluids.
8. Encourage the person to seek medical attention.

### Myth-Information

**Myth:** When a person has frostbite, you should rub the frozen area or apply snow to it.

When providing care for frostbite, never rub the frozen area or put snow on it. Rubbing the area can cause the ice crystals within the cells to damage the surrounding tissues, and rubbing snow on the area will only worsen the condition.

## Hypothermia

Hypothermia occurs when the body's core temperature (i.e., the temperature of the heart, lungs, and brain) drops to 35°C (95°F) or lower. Hypothermia becomes a life-threatening condition when the core temperature drops below 28°C (82.4°F). People with smaller body types, lower overall body weights, and lower body-fat content are much more likely to experience hypothermia. Hypothermia can occur at any time of year, with young children and the elderly being most at risk.

### Common Causes

The following may lead to hypothermia:

- Sweating while outdoors in cold weather
- Cold, wet, windy conditions
- Cold-water immersion, especially in water below 10°C (50°F)
- Dehydration
- Taking medications to eliminate water from the body

### Prevention

The following tips will help to prevent hypothermia:

- Prepare for activities in cold environments by wearing appropriate clothing:
  - Wear a hat and clothing made of tightly woven fibres (e.g., wool, fleece), which provide insulation and allow moisture transmission from the skin to the external environment.
  - Avoid cotton because it soaks up water and stays wet.
  - Wear clothing in layers so that they can be added or removed according to the weather conditions and exercise intensity. Remove some clothing before working to reduce the amount of sweat and keep clothing dry.
- Keep clothing dry. If clothing gets wet, change into dry clothing as soon as possible.
- Carry and consume high-energy foods that have a lot of sugar.

- Drink plenty of warm, high-calorie drinks to help fuel heat production. If high-calorie drinks are not available, drink plenty of water to at least keep from becoming dehydrated.
- Do not drink alcohol.
- Take frequent breaks from the cold to let the body warm up.
- Increase your activity level, add insulation, add a heat source, or get out of the cold if shivering occurs.
- Be careful around cold water and always wear a personal flotation device (PFD).

### What to Look For

#### Levels of Cold Stress

##### Cold Stress (Not Hypothermic)

- Shivering
- Normal mental status
- Able to care for self

##### Mild Hypothermia

- Vigorous shivering; complaining of the cold
- Decreased physical function
- Difficulty taking care of self

##### Moderate Hypothermia

- Weak and intermittent shivering or shivering that later stops
- Sometimes complaining of the cold
- Lack of coordination or speech; confused or unusual behaviour
- Impaired judgment
- Possible unresponsiveness

##### Severe Hypothermia

- Shivering has stopped
- Unresponsiveness; breathing has slowed down or stopped
- Body feels stiff
- No pulse

## What to Do

### Check

Hypothermia can slow a person's breathing, so you may need to spend longer than usual checking the person's ABCs. If the person is unresponsive and you suspect hypothermia, check for signs of breathing for 60 seconds.



### Call

Call EMS/9-1-1 and get an AED if the person has moderate to severe hypothermia.



### Care

When caring for hypothermia, handle the person very gently and keep him or her horizontal. You should only move the person if it is absolutely necessary. Take care to avoid jostling or bumping him or her. Remove wet clothing, but only in an area sheltered from the cold environment.

1. Insulate the person (e.g., by applying a hypothermia wrap) to protect him or her against further cold exposure.
2. Warm the person by applying warm water bottles, heating pads, or electric blankets to the person's upper torso (i.e., the armpits, chest, and upper back). Body heat from yourself or a bystander can also work in an emergency. Never put the person in a warm bath or shower or try to warm the person with a fire, a stove, or hot rocks.
3. If the person is responsive, provide warm, sugary, non-alcoholic liquids to drink.


1. From outside ring to centre: assess responsiveness, movement, shivering, alertness
2. Assess whether **normal function**, or **impaired or no function**
3. Treat according to appropriate result-quadrant

### Cold Stressed, Not Hypothermic

1. Reduce heat loss (e.g., add dry clothing)
2. Provide high-calorie food or drink
3. Increase heat production (e.g., exercise)

### Mild Hypothermia

1. Handle gently
2. Keep horizontal
3. No standing/walking for at least 30 min.
4. Insulate/vapour barrier
5. Apply heat to upper trunk
6. Provide high calorie food/drink
7. Monitor until improvement (at least 30 min.)
8. Call EMS/9-1-1 if no improvement



### Severe Hypothermia

1. Treat as moderate hypothermia, and
  - a) IF no obvious vital signs, **THEN 60-second breathing check**
  - b) IF no breathing, **THEN start CPR**
2. Call EMS/9-1-1

### Moderate Hypothermia

1. Handle gently
2. Keep horizontal
3. No standing/walking
4. No drink or food
5. Insulate/vapour barrier
6. Apply heat to upper trunk
7. Call EMS/9-1-1

### SUGGESTED SUPPLIES FOR A HYPOTHERMIA WRAP:

- |   |  |
|---|--|
| 1 - Tarp or plastic sheet for vapour barrier outside sleeping bag | 1 - Plastic or foil sheet (2 x 3 m) for vapour barrier placed inside sleeping bag                                |
| 1 - Insulated ground pad  | 1 - Source of heat (e.g., <i>chemical heating pads/blankets or warm water in a bottle or hydration bladder</i> ) |
| 1 - Hooded sleeping bag (or equivalent)                           |  |


### INSTRUCTIONS FOR HYPOTHERMIA WRAP

**1. Dry or damp clothing:** *Leave clothing on*

*IF shelter/transport is **less than** 30 minutes away,  
**THEN wrap immediately***

**2. Very wet clothing:**

*IF shelter/transport is **more than** 30 minutes away,  
**THEN protect person from environment, remove wet clothing, and wrap***



### Applying a Hypothermia Wrap

Applying a hypothermia wrap is one way to minimize the heat lost by a person with hypothermia. It consists of a number of layers of insulation and heat-reflecting material.

To apply a hypothermia wrap:

1. Place an insulation pad (or pads) between the person and the ground.
2. Apply as much insulation as possible. Add extra clothing and wrap the person in blankets or sleeping bags.
3. Cover the person's head and neck with a toque, heavy hat, or hood.
4. Place a vapour barrier (plastic or foil blanket) outside the insulation wrap if the person is dry. If the person is still wet, place the vapour barrier inside the insulation wrap. If two vapour barriers are available, place one inside and one outside the insulation wrap.

If the person's clothing is wet and shelter or transport is less than 30 minutes away:

1. Leave the person in the hypothermia wrap.
2. Seek protection from the wind.
3. Create shelter if necessary.
4. Remove wet clothing. Cut the clothing to remove it, if necessary.
5. Dry (blot, don't rub, the skin) and rewrap the person.
6. Provide continual care until EMS personnel arrive.

## Freezing of Skin to Metal Objects

A person's skin (especially the tongue and lips) can freeze to cold metal objects. This is especially a risk when skin is wet or moist. While this is a concern mainly in cold weather, skin can also freeze to metal objects in other situations (e.g., skin freezing to cold appliances such as freezers).

### What to Do



#### Call

Call EMS/9-1-1 if you cannot safely remove the body part.



#### Care

1. Do not pull or tug the frozen body part.
2. Pour warm (not hot) water on the surface of the object or the skin that is stuck to the object.
3. As the skin begins to come free, gently help release the person from the metal object.
4. Treat any torn skin as an open wound.



## Snow Blindness

### Common Causes

Snow blindness occurs when a person's eyes are exposed to ultraviolet rays. It most commonly occurs when the sun's light is reflected from snow, ice, sand, or water into a person's eyes. Snow blindness can occur even on cloudy days.

## Prevention

To help prevent snow blindness, wear sunglasses that block 100% of UV rays and prevent light from shining in from below, above, or the sides.

### What to Look For

Signs and symptoms of snow blindness may include:

- Redness of the eyes
- Swelling of the tissue around the eyes
- Pain, itchiness, or a burning sensation in the eyes that may become intense
- Temporary colour changes in vision, or even vision loss



The signs and symptoms of snow blindness may not appear for several hours following exposure to bright, snowy conditions.

### What to Do



#### Care

1. Place the person in a darker environment, if possible, or cover his or her eyes.
2. Apply a cool, damp cloth to reduce pain and burning.
3. If the person's vision is affected, seek medical attention.



## Cold-Water Immersion

Cold-water immersion can occur in open water (e.g., when a boat capsizes) or after breaking through the ice. Although hypothermia occurs faster in cold water than cold air, the signs, symptoms, and care are essentially the same.

Cold-water immersion has four distinct phases:

- 1. Cold-Shock Response:** Rapid cooling of the skin causes a gasp followed by hyperventilation (lasting 1 to 2 minutes).
- 2. Cold Incapacitation:** Further cooling of nerve and muscle fibres causes weakness and a loss of coordination, leading to incapacitation within approximately 15 minutes.
- 3. Hypothermia:** A person dressed appropriately for the weather conditions will take 30 minutes or more to become mildly hypothermic.
- 4. Circum-Rescue Collapse:** Can occur just prior to, during, or after rescue. The signs and symptoms range from collapsing, to fainting, to cardiac arrest. When rescuing a person from cold water, monitor the person's condition carefully and be prepared to provide emergency care.

Most deaths in cold water result from the Cold-Shock Response (when the head is under water) or drowning due to Cold Incapacitation. A properly worn personal flotation device (PFD) will keep a person afloat after he or she loses the ability to swim during the Cold Incapacitation phase.

## What to Do (Rescue)

### Self-Rescue

- If you are in open water, do a self-rescue by getting into a boat, raft, or other mode of marine transportation, if available.
- If you have fallen through the ice, place your arms on the surface of the ice, kick your legs until your body is horizontal to the water's surface, and then kick and pull forward until you are on the ice. Roll away from the hole before standing up.

### Survive to Await Rescue

If you are in open water and self-rescue is not possible, increase your survival time by doing the following:

- Get out of the water as much as possible by using a capsized boat or other floating object as a flotation device.
- If you are wearing a personal flotation device (PFD), adopt the Heat Escape Lessening Position (HELP):
  - Press your arms against your armpits.
  - Place your forearms across your chest.
  - Squeeze your thighs together and raise your knees toward your chest.
- If several people are together, adopt the HUDDLE position:
  - Face each other.
  - Have each person squeeze his or her thighs together.
  - Form a circle and hug each other. Each person should place one arm above and one arm below the arms of those adjacent, and pull the sides of the chests together.
  - Sandwich any children or persons without a PFD inside the HUDDLE.

If you have fallen through the ice and are unable to exit by yourself, stop struggling. Place your arms on the ice and let them freeze in place. This will prevent you from drowning if the cold exposure causes incapacitation or unresponsiveness.

### First Aider Rescue

To help rescue a person from cold water:

- 1. Talk** – Instruct the person on self-rescue.
- 2. Throw** – Throw a rope or buoyant object to the person.
- 3. Reach** – Reach out to the person with a rigid object like a ladder, paddle, or tree branch.



Any person rescued from cold water should be treated for hypothermia.



Unless you have received specific training in water rescue, never enter the water or approach a hole in the ice to attempt to help a person.

### Drowning

Drowning occurs when a person's airway is submerged in water, cutting off the oxygen supply to body.

Drowning is a leading cause of death for children and babies. An average of 400 Canadians drown each year. Young children aged 1 to 4 and men aged 15 to 44 are at the greatest risk.

### Prevention

The following tips will reduce the risk of drowning:

- Always wear a personal flotation device (PFD) when in a boat or other mode of marine transportation.
- Supervise children in, on, and around any body of water.
- Those who do not swim or are weak swimmers should wear a PFD when in, on, and around any body of water.
- Check water depth before swimming or diving.
- Have appropriate safety equipment available when in or on the water.
- Take Canadian Red Cross Swimming and Water Safety lessons.

### Responsive Drowning Person

#### What to Look for

These common warning signs can indicate that someone is drowning:

- The person is vertical in the water, and is not using his or her legs to move forward or tread water.
- The person may have an expression of fear.
- The person's arms may be flailing up and down as he or she tries to rise out of the water to breathe, rather than trying to swim forward.
- The person's movements are uncontrolled.

## What to Do



### Call

Call for a lifeguard or other trained person (if one is nearby), or call EMS/9-1-1.



### Care

1. Remove the person quickly and safely from the water, but do not put yourself in danger. Follow the same steps you use to perform a First Aider Rescue for cold-water immersion. Remember, any person rescued from cold water should be treated for hypothermia.

## Unresponsive Drowning Person

### What to Do



### Call

Call EMS/9-1-1 and get an AED.



### Care

1. If the person is not breathing, begin CPR.



## Seasickness

Seasickness (motion sickness) is a sudden illness largely due to the motion of a ship or vessel. People who are not accustomed to the sea are most susceptible, but even experienced seafarers may be affected in rough water conditions.

Signs and symptoms of seasickness vary, but include:

- Loss of appetite, nausea, vomiting, or abdominal cramps
- Headaches, dizziness, or exhaustion
- Cold sweat
- Dry mouth



### Care

- In mild cases, the condition will gradually wear off (often during sleep) and no specific treatment is necessary
- Providing small portions of dry food (such as crackers) can help settle the person's stomach
- Sucking on ice chips instead of drinking water will reduce the risk of vomiting while still preventing dehydration
- More severe cases of prolonged vomiting may be managed with over-the-counter preventive medications.



# 13 Poisons

A poison is a substance that has a harmful effect within the body if it is inhaled, swallowed (ingested), absorbed, or injected. Poisons are immediately life-threatening if they affect breathing or circulation. Practically anything can be a poison if it is not meant to be taken into the body. Even some substances that are meant to be taken into the body, such as medications, can be poisonous if they are taken by the wrong person, or if the person takes too much. Combining certain substances can also result in poisoning.

Poisoning can happen anywhere, but most poisonings take place in the home. Children younger than 5 years, especially toddlers, are at the highest risk for poisoning. Children may be attracted to pretty liquids in bottles, sweet-smelling powders, berries on plants that look like they are edible, or medications or vitamins that look like candy. Additionally, very young children explore their world by touching and tasting things around them, so even substances that do not look or smell attractive are poisoning hazards among this age group. Older adults who have medical conditions that cause confusion (such as dementia) or who have impaired vision are also at a higher risk of unintentional poisoning.

When providing first aid for a poisoned person, use caution to avoid contact with the poison.

Your local Poison Control Centre can provide you with specific first aid instructions for a variety of types of poison, but remember that you should always call EMS/9-1-1 if the person has an altered level of responsiveness or is having difficulty breathing. Keep the local Poison Control Centre number by your telephone and save it as a contact in your mobile phone.

## TYPES OF POISONS

### Swallowed Poison



Enters the body through the mouth, lips, esophagus, or stomach. Drinking bleach is an example of swallowed poisoning.

### Absorbed Poison



Enters the body through the skin. Plants (such as poison ivy) and chemicals can cause absorbed poisoning.

### Injected Poison



Enters the body through bites or stings or as drugs injected with a needle.

### Inhaled Poison



Is breathed into the body. Breathing in carbon monoxide from a car's exhaust is an example of inhaled poisoning.

## HOUSEHOLD POISONS

Many everyday household items can be poisonous if they are used incorrectly. Common causes of unintentional poisonings at home include:

- Alcohol (found in many products, including hand sanitizer, mouthwash, perfume, cologne, aftershave, and vanilla extract)
- Medications (over-the-counter and prescription) and vitamins
- Cleaning products (detergent "pods" are especially attractive to children)
- Glues and paints (lead paint remains toxic even when it is applied, and children may put peeling paint into their mouths)
- Insect and weed killers
- Car products (e.g., antifreeze, windshield washer fluid)
- Plants (both houseplants and outdoor plants)
- Oils, lubricants, and polishes
- Tobacco

### Identifying Hazardous Materials

The international hazard symbols on product labels make it easier to identify harmful materials.

If you must handle a product that is labelled with any of the international hazard symbols:

- Carefully read the warnings and follow the instructions.
- Wear protective equipment (such as eye goggles, a face mask, and disposable gloves).
- Ensure the product is properly sealed and stored after use.

## THE INTERNATIONAL HAZARD SYMBOLS



### Flammable

These materials catch fire easily and burn quickly when exposed to any form of ignition (such as fire or heat).



### Toxic

These materials can harm the respiratory (breathing) system, nervous system, and other systems if they are absorbed through the skin, inhaled, or ingested.



### Explosive

These materials can explode when exposed to heat, flame, or pressure.



### Corrosive

These materials can burn the skin and cause permanent blindness.

## Prevention

### Prevention at Home

The following tips will help reduce the risk of poisoning:

- Keep all medications, household cleaning products, poisonous plants, and other toxic substances well out of the reach of children. Use locked cupboards or child-resistant latches.
- Use child-resistant safety caps on medications and other potentially toxic products.
- Never call medicine "candy" to persuade a child to take it.
- Teach children to check with an adult before eating an unknown substance.
- Keep potentially poisonous products in their original containers with their original labels.
- Ensure that prescription medicine is taken only by the person whose name is on the label.
- Carefully dispose of expired medications through your local hazardous waste process.
- Be aware that many cleaning products and aerosol sprays have toxic fumes. Use potentially poisonous chemicals in a well-ventilated area.
- Wear proper protective clothing any time you are working with or around a poisonous substance.
- Run gas and other combustion engines only in open, well-ventilated areas.
- Learn about poisonous plants in your area and before going into a new environment.

### Prevention in the Workplace

The following tips will reduce the risk of poisoning in the workplace:

- Clearly label all toxic substances and never use a product that is not clearly identified.
- Have a detailed MSDS (Material Safety Data Sheet) available for every hazardous substance in the workplace.
- Ensure that workers who might be exposed to hazardous materials receive the proper training in safety measures and emergency procedures.
- Check all warning labels, tags, and posters in the workplace and follow their instructions carefully.
- Read labels and the MSDS to find out the risks of each hazardous material, the safety measures that must be taken to prevent poisoning, and the first aid required if poisoning occurs.

### General Care for Poisoning

Your check of the scene and the person might give clues that point to poisoning as the cause of the person's illness. For example, you may note an open or spilled container, an unusual odour, burns around the person's mouth, a strange odour on the person's breath, or other people in the area who are also ill.

If you think that a person has been poisoned, try to find out the following key points so that you and others can give the most appropriate care:

- The type of poison
- The quantity taken
- When it was taken

Calling EMS/9-1-1 or a Poison Control Centre is the most important thing you can do if a poisoning of any type is suspected. Other general first aid care tips for poisoning include the following:

- Limit further exposure by moving either the person or the source of the poison.
- If the poison's container is found nearby, give the information from the label to the Poison Control Centre or EMS/9-1-1 dispatcher.
- Do not give the person anything to eat or drink unless an EMS dispatcher or Poison Control Centre staff member tells you to do so.
- If you do not know what the poison was and the person vomits, save a sample to give to EMS personnel.

## Swallowed Poisons



### What to Look For

The following may indicate that a poisonous substance has been swallowed:

- An open container of poison nearby
- Burns around the mouth
- Increased production of saliva or salivation that is an abnormal colour
- Abdominal cramps, vomiting, or diarrhea
- Seizures
- Dizziness or drowsiness
- Unresponsiveness
- A burning sensation in the mouth, throat, or stomach

## What to Do

### Call

If the person is responsive and alert and his or her ABCs are unaffected, call the local Poison Control Centre. Call EMS/9-1-1 and get an AED if the person has an altered level of responsiveness or has difficulty breathing.

### Care

1. If the person is not breathing, start CPR. Use a barrier device so that you don't contaminate yourself with the poison.
2. Check the packaging of the poison, if possible, so that you know what it is.
3. Induce vomiting only if told to do so by the EMS dispatcher or the Poison Control Centre.
4. If the person needs to go to the hospital, bring a sample of the poison (or its original container).

### Myth-Information: Induce Vomiting in a Poisoned Person

**Myth: Inducing vomiting in a poisoned person will remove the harmful substance from the person's body.**

Inducing vomiting in a person who has been poisoned often causes additional harm and is not recommended. Sometimes the person may vomit on his or her own, but you should never give the person anything to make him or her vomit unless you are specifically instructed to do so by an EMS dispatcher or a Poison Control Centre staff member.

## Inhaled Poisons

### What to Look For



The following may indicate that a poisonous substance has been inhaled:

- Breathing difficulties
- Irritated eyes, nose, or throat
- Dizziness
- Vomiting
- Seizures
- Bluish colour around the mouth
- Unresponsiveness
- An unusual smell in the air

### What to Do



#### Call

If the person is responsive and alert and his or her ABCs are unaffected, call the local Poison Control Centre. Call EMS/9-1-1 and get an AED if the person has an altered level of responsiveness or has difficulty breathing.



#### Care

1. If the person is not breathing, start CPR. Use a barrier device so that you don't contaminate yourself with the poison.
2. Get the person into fresh air but do not enter into a hazardous atmosphere in order to do so.

Inhaled poisons can affect everyone in an area. Stay out of the area if you suspect that the poison may still be in the air.

## Carbon Monoxide Poisoning

Carbon monoxide (CO) is a gas that has no smell, colour, or taste. CO poisoning is often called a "silent killer" because it is not detectable to any of the body's senses. CO bonds to red blood cells 200 times better than oxygen, preventing oxygen from attaching and therefore starving the body of oxygen.

It releases when fuel is burned in small engines, lanterns, fireplaces, stoves, grills, gas ranges, furnaces, cars, and trucks. When equipment that burns these fuels is properly ventilated, CO poisoning is not a problem. But if the equipment or ventilation system is faulty, or if outdoor equipment is used in an enclosed area, toxic levels of CO can build up quickly, creating the risk of CO poisoning. Concentrated CO is poisonous and life-threatening to those who inhale it.

### Prevention

The following tips help prevent carbon monoxide (CO) poisoning:

- Ensure that a battery-operated or battery back-up CO detector is installed in the home or workplace where the alarm will be easily heard.
- Check the CO detector's batteries twice a year, ideally once in the fall and once in the spring.
- Have all fuel-based appliances and equipment installed and repaired by a qualified technician.
- Have all fuel-based appliances and equipment serviced annually by a qualified professional.
- Use fuel-based appliances and materials as intended (e.g., do not use generators or portable flameless chemical heaters indoors, do not use a gas stove or oven for heating, do not burn charcoal indoors).
- Ensure that chimneys are inspected and cleaned annually by a professional.
- Never run a car or truck inside an attached garage. For detached garages, always leave the garage door open when running a car or truck inside.

## What to Look For

The signs and symptoms of acute carbon monoxide (CO) poisoning are commonly mistaken for a stomach virus, the flu, or food poisoning. Signs and symptoms include the following:

- Headache
- Dizziness or light-headedness
- Confusion
- Impaired hearing and vision
- Weakness or fatigue
- Muscle cramps
- Nausea and vomiting
- Chest pain
- Altered level of responsiveness

## What to Do



### Care

1. Treat the person as you would for any other type of inhaled poison.

## Absorbed Poisons

### What to Look For



The following signs and symptoms can indicate that a poisonous substance has been absorbed:

- Rash or hives (raised, itchy areas of skin)
- Burning or itching skin
- Swelling
- Blisters
- Burns
- Unresponsiveness

## What to Do



### Call

If the person is responsive and alert, and the person's ABCs are unaffected, call the local Poison Control Centre. Call EMS/9-1-1 and get an AED if the person has an altered level of responsiveness or has difficulty breathing.



### Care

1. If the person is not breathing, start CPR.
2. If the poison is a dry powder, brush it off the person's skin. Be careful to avoid contaminating yourself.
3. Remove any clothing or items covered in the poison.
4. Flush the skin with running water for at least 15 minutes. To prevent any further injury, make sure the water flushes away from any unaffected areas of the body.

## Rash-Causing Plants

Some plants produce chemicals that can cause absorbed poisoning on contact. These plants vary in appearance depending on the location, species of the plant, and time of year. As prevention is the best strategy, it is a good idea to become familiar with the appearance of rash-causing plants in your area.

### Prevention

When there is the potential for exposure to rash-causing plants, follow these guidelines:

- Wear a long-sleeved shirt, long pants, and closed-toed shoes.
- Apply a pre-contact barrier cream or lotion before going outside.
- Wash skin with a specialized skin cleanser designed to remove plant oils or a degreasing soap (such as dishwashing liquid) and plenty of water immediately after returning indoors.
- Wash tools, work gloves, and clothing that may have contacted rash-causing plants, as oils can remain on these items and transfer to the skin the next time they are used.

If you suspect that your skin has come into contact with a rash-causing plant, wash the entire area with a degreasing soap immediately.

### Poison Ivy, Sumac, and Oak

Plants such as poison ivy, poison sumac, and poison oak are covered with an oil called urushiol, which causes a skin reaction in most people.



### What to Look For

Contact with urushiol can cause the following signs and symptoms:

- Itchy skin
- Reddening of the skin
- Bumps
- Blisters

The rash can range from irritating to unbearable, depending on the person's sensitivity, the amount of skin exposed, and the rash's location.



Do not burn plants containing urushiol. The oil is carried in the smoke, and inhalation can cause swelling of the throat and irritation of the lungs.

### What to Do



Call EMS/9-1-1 only if the person experiences a severe allergic reaction or is having trouble breathing.



1. Encourage the person to apply a cream or ointment designed to reduce itching and blistering (e.g., calamine).
2. Suggest that the person take an oral antihistamine to help relieve itching.
3. If the rash is severe or on a sensitive part of the body (such as the face or groin), the person should see a healthcare provider.

## Giant Hogweed and Wild Parsnip

The sap of giant hogweed and wild parsnip contains toxic photosensitizing compounds called furanocoumarins. Contact with the plant transfers these compounds to the skin. While they do not have an immediately visible effect, they cause the skin to react when exposed to the sun's UV radiation to produce severe inflammation, intense burning, and weeping blisters. If any of these plants are found, their location should be reported to the local environmental or municipal authorities.



### What to Look For

If the person's skin has come into contact with the sap of giant hogweed or wild parsnip, and is then exposed to the sun's UV radiation, the following signs and symptoms may occur:

#### 24 hours after exposure:

- Swelling of the skin
- Reddening of the skin

#### 48 hours after exposure:

- Painful blistering
- Purplish scarring of the skin

### What to Do

#### Call

Call EMS/9-1-1 if the person is having trouble breathing, or if the sap is on the person's eyes, face, or groin.

#### Care

1. Protect the area from sunlight.
2. If sap gets into the eyes, rinse them thoroughly with water for at least 15 minutes, or until EMS personnel arrive.
3. Encourage the person to seek medical attention.

If a reaction occurs, the affected area may be sensitive to sunlight for months or even years. Protect the area by keeping it covered, wearing sunglasses, and applying sunscreen.

## RASH-CAUSING PLANTS

### Poison Ivy



Found in all the provinces except Newfoundland and Labrador, but has not been reported in any of the territories.

### Poison Oak



Found in British Columbia

### Poison Sumac



Found in Southern Ontario and Quebec

### Giant Hogweed



Found in Nova Scotia, New Brunswick, Quebec, Ontario, and British Columbia

### Wild Parsnip



Found in all provinces and territories except for Nunavut

## Injected Poisons

### What to Look For



### What to Do



#### Call

If the person is responsive and alert, and the person's ABCs are unaffected, call the local Poison Control Centre. Call EMS/9-1-1 and get an AED if the person has an altered level of responsiveness or has difficulty breathing.



#### Care

1. Clean the puncture site with clean running water.
2. Keep the person as still as possible.



## Poisoning Caused by Alcohol or Drugs

### Drug Overdose

Drugs (whether over-the-counter, prescription, or illegal) are a frequent cause of death by poisoning. Drug overdoses may be accidental or intentional. Signs and symptoms will vary depending on the drug.

#### Stimulants

- Stimulants affect the brain and nerves to speed up physical and mental activity.
- Many stimulants are swallowed as pills, but some can be absorbed or inhaled.

#### Hallucinogens

- Hallucinogens cause changes in mood, sensation, thought, emotion, and self-awareness.
- They can cause intense fear, panic, paranoid delusions, vivid hallucinations, deep depression, tension, and anxiety.

#### Depressants

- Depressants send signals to the brain and nerves that slow down physical and mental activity.
- They cause drowsiness and impair coordination and judgment.
- Alcohol is the most widely used and abused depressant in Canada.

#### Designer Drugs

- Designer drugs don't fit into any of the categories mentioned above.
- They are often chemically altered versions of medical drugs, such as narcotics and amphetamines.
- Their effects can be unpredictable and dangerous.
- Ecstasy, GHB, ketamine, and LSD are examples of designer drugs.

### What to Look For

The signs and symptoms of drug poisoning will vary depending on the drug but often include the following:

- Moist or flushed skin
- Sweating
- Chills or fever
- Nausea or vomiting
- Vomiting blood or passing blood
- Changes in breathing or difficulty breathing
- Racing or pounding heart
- Continuous pain or pressure in the chest or abdomen
- Seizures
- Severe headache
- Slurred speech
- Changes in level of responsiveness
- Changes in behaviour (e.g., aggression or fear)

### What to Do



#### Call

Call EMS/9-1-1 and get an AED if you suspect a drug overdose or if the person is having seizures, has difficulty breathing, is unresponsive, or has a change in behaviour.



#### Care

1. Try to find out from others at the scene what substance or substances the person may have taken, and then communicate that information to EMS personnel or the Poison Control Centre worker.

## Opioid Overdose

Opioid drugs, such as heroin, oxycodone, and fentanyl, are a common cause of drug overdose in Canada. Many prescription drugs are or contain opiates. EMS personnel use naloxone (Narcan®) to reverse the effects of opioid drugs. In some provinces, First Aiders, first responders, and people in high-risk addiction situations can receive training in administering naloxone. Naloxone can be supplied as a nasal spray or an injectable solution.



### Naloxone and Opioid Drug Overdoses

Naloxone is used to quickly reverse the effects of an opioid drug overdose, specifically counteracting central nervous system depression and respiratory depression.

Administering naloxone requires special training. If you are trained and naloxone is available, you can give naloxone to someone suffering from an opioid drug overdose, as this is an effective and lifesaving treatment.

### What to Look For

Signs and symptoms of an opioid overdose include the following:

- Slowed breathing (or no breathing)
- Chest tightness
- Extreme drowsiness or unresponsiveness
- Small pupils

### What to Do



Call EMS/9-1-1 and get an AED.



1. If the person is unresponsive and not breathing, begin CPR. Do not delay CPR treatment to give the person naloxone.
2. If the person is unresponsive but is breathing, and if you have received training in its use, give the person naloxone by following the instructions on the label. Always follow specific policies and the training you have received in providing naloxone.
3. Stay with the person and provide continual care until EMS personnel arrive. Ensure that the person receives professional medical care, as naloxone alone is not enough to care for a person who has overdosed on opioids.

## Alcohol Poisoning

Alcohol is a depressant that affects the central nervous system. It impairs judgment, slows down reflexes, and makes driving unsafe. Very high levels of alcohol in the bloodstream can affect the brain's ability to control the body's breathing, heart rate, and temperature, which can lead to death. Alcohol poisoning occurs when large quantities of alcohol are consumed in a short period of time, resulting in toxic levels of alcohol in the body.

There are many factors that can contribute to an unsafe blood alcohol level:

- Speed of drinking
- Full or empty stomach
- Type of alcohol
- The person's weight and sex

## What to Look For

The signs and symptoms of alcohol poisoning include the following:

- Confusion
- Vomiting
- Seizures
- Slow or irregular breathing
- Low body temperature (hypothermia)
- Unresponsiveness
- Blue-tinged skin or skin that is paler than normal

## What to Do



### Call

Call EMS/9-1-1 and get an AED.



### Care

1. Roll the person into the recovery position.
2. Carefully monitor the person until EMS personnel arrive, as his or her condition can deteriorate rapidly.

## Bites and Stings

Bites and stings can range in severity from mildly irritating to life-threatening. When a person is bitten or stung, proper first aid care can help to limit complications and speed healing, and it may even be lifesaving.



## Animal Bites

Any animal that has teeth, whether domesticated (e.g., pets or livestock) or wild, can be the source of a bite wound. If the person does not know the animal, rabies may be a concern. Most animal bites carry a high risk of infection, so they should be monitored closely in the hours and days after the incident.

## What to Do



### Call

Call EMS/9-1-1 if bleeding is life-threatening. Call your local animal control department if the animal is wild or a stray.



### Care

1. Try to get the person safely away from the animal without putting yourself in danger.
2. Do not try to capture the animal.
3. If the wound is bleeding heavily, provide care for external bleeding.
4. If the wound is minor:
  - Wash the wound thoroughly with clean water, ideally by running it under a tap, as the pressure from the running water lets it penetrate more deeply into the wound.
  - Control any bleeding.
  - Encourage the person to apply antibiotic ointment or cream to the wound, if they have no known allergies or sensitivities.
  - Cover the wound with a dressing or bandage.

5. Seek medical attention if the animal is wild, stray, or unknown to you, or if you suspect the animal might have rabies.
6. Watch for signs and symptoms of infection and ensure that the person continues to monitor the wound over the next several days.

## Rabies

Rabies is a serious infection that attacks the brain and spinal cord. It is fatal if it is not treated. The virus that causes rabies is spread when an animal that has the disease bites another animal or a person. Wild animals (such as foxes, skunks, bats, and raccoons) can carry rabies, as can pets and livestock that have not been vaccinated against it. Only mammals can be infected with rabies.

Animals with rabies may act strangely. For example, an animal that is usually active at night may appear in the daytime, or a normally shy wild animal might not run away when people approach. Rabid animals may drool, appear to be partially paralyzed, or be aggressive or strangely quiet. Call EMS/9-1-1 if a person is bitten by an animal that you suspect might have rabies. If possible, try to remember details about the animal's behaviour and appearance, and where you last saw it. When you call EMS/9-1-1, the dispatcher will direct the proper authorities (e.g., animal control) to the scene.

A person who is bitten by an animal that might have rabies must get medical attention immediately. Treatment for rabies includes a series of injections to build up immunity that will help fight the disease.



## Insect Stings

Most of the time, insect stings are merely uncomfortable. However, allergic reactions and anaphylaxis are always a concern. Some insects, such as bees, leave their stingers embedded in the person's skin. Others, such as wasps, can sting multiple times, but leave no stinger behind.

## Prevention

To reduce the risk of insect stings:

- Wear a long-sleeved shirt, long pants, and closed-toe shoes.
- Don't wear perfume, cologne, or other products with strong scents.
- Remove or cover items that will attract insects, such as garbage or food.

## What to Look For


The following can indicate an insect sting, especially if you see insects nearby:

- Quick, sharp pain at the site of the sting
- Pain, redness, or swelling at the site of the sting
- A stinger embedded in the skin

## What to Do



Call EMS/9-1-1 and get an AED if there are any signs of a severe allergic reaction.



### Tick Bites

Ticks are found in wooded, bushy areas, in tall grass, and in leaf litter on the ground. They can attach themselves to any warm-blooded animal that they come into direct contact with, including people. When ticks attach themselves to the skin, they can spread pathogens

from their mouths into the person's body. These pathogens can cause serious illnesses, such as Lyme disease. Promptly remove any ticks that you find before they become swollen with blood. Ticks are most active from April to September.



### Care

1. If the stinger is still in the person's skin, remove it by scraping it away from the skin. Use a plastic card (such as a debit card) to do this.
2. Wash the area with clean water.
3. Cover the site with an adhesive bandage.
4. Apply ice or a cold pack to help control swelling. Put a thin cloth between the cold source and the person's skin to avoid freezing the skin.
5. Continue to watch for signs of infection, an allergic reaction, or anaphylaxis.



### Prevention

Tips to help prevent tick bites include the following:

- Avoid wooded areas and areas with high grass and leaf litter.
- Walk in the centre of hiking trails.
- Limit the amount of exposed skin. Wear long-sleeved shirts and long pants. Tuck your shirt into your pants and your pant legs into your socks or boots.
- Use insect repellents with 10 to 30% DEET on skin and clothing. A parent, guardian, or caregiver should apply repellents to children, carefully avoiding the eyes, mouth, and hands.
- Check gear and pets for ticks when coming in from outside.
- Immediately after being outdoors, do a head-to-toe check for ticks using a hand-held or full-length mirror. Check the scalp, under the arms, in and around the ears, inside the navel, around the waist, behind the knees, and between the legs. If you are outdoors for an extended period of time, check several times throughout the day. Caregivers and parents or guardians should check children for ticks.
- Put outdoor clothing in a tumble dryer on high heat for 1 hour to kill any ticks that might be attached.
- When walking with children in a tick-infested area, prevent ticks from entering their clothing by using a rubber band or tape to seal the area where children's pants and socks meet.

### Using Insect Repellents to Prevent Tick Bites

Consider using an insect repellent if you will be in a grassy or wooded area for a long period of time, or if you know that the tick population in the area is high. Use repellents sparingly, as one application will last 4 to 8 hours and heavier or more frequent applications do not increase effectiveness.

## What to Do




### Care

If the tick hasn't started to dig into the flesh, remove it by brushing it off the skin.

If the tick has begun to bite:

1. Use tweezers to grasp the tick by the head as close to the person's skin as possible.
2. Pull upward slowly and steadily without twisting until the tick releases its hold. If you cannot remove the tick or if its mouthparts stay in the skin, the person should seek medical attention.



3. If the tick is removed, wash the area with clean tap water. The pressure of the running water lets it penetrate more deeply into the wound.
4. If possible, save the tick in a resealable bag and record the date of the bite.
5. Advise the person to monitor the bite for several days for signs and symptoms of infection. If the area becomes infected (i.e., it becomes red, warm, or painful), or the person develops a fever, he or she should seek medical attention.



Ticks can be infected with more than one type of bacteria that can cause human illness. Signs can appear weeks to months after a bite. If a person feels unwell after a tick bite, he or she should seek medical attention.

### Myth-Information: Properly Removing a Tick

**Myth: You can remove a tick safely by burning it off with a flame or smothering it with petroleum jelly or nail polish.**

These folk remedies are not the most effective methods. They rely on the tick detaching itself, which could take hours. As long as the tick's mouthparts are in contact with the skin, the tick is potentially transmitting disease. The goal is to remove the tick in one piece as quickly as possible. The best tool for doing this is a pair of fine-tipped tweezers or a special tick removal tool, such as a tick key.



### Lyme Disease

Lyme disease is caused by bacteria that are transmitted to humans through bites from infected ticks. If Lyme disease is not treated, it can become a chronic condition with serious symptoms such as cognitive dysfunction, joint and muscle pain, and neurological disorders.

## What to Look For

The most distinctive sign of Lyme disease is a small red rash centred on the location of the bite in the shape of a bullseye (a red dot on a paler circle with an outer red ring). These rashes are typically 13 to 18 cm (5 to 7 in.) across.

Other signs and symptoms of Lyme disease include:

- Fatigue
- Headache
- Fever or chills
- Swollen lymph nodes
- Numbness or tingling
- Muscle spasms or weakness
- Joint and muscle pain (similar in feeling to the flu)

## What to Do

Seek medical attention as soon as possible. If possible, save the tick responsible for the bite in a resealable bag or empty pill bottle, and bring it to the medical appointment. It may help the doctor diagnose the illness.



## Snakebites

In the wild, snakes like to occupy rock outcrops, swamps, undergrowth, and abandoned human structures. When disturbed, some snakes can inject venom as they bite.

The vast majority of Canada's snakes are non-venomous, but there are three venomous snakes currently native to Canada. All three are rattlesnakes:

1. Northern Pacific Rattlesnake
2. Massasauga Rattlesnake
3. Prairie Rattlesnake

Other venomous snakes in North America include copperheads, cottonmouths (water moccasons), and coral snakes. Most deaths from venomous snakebites occur because:

- Too much time passed before the person received medical care.
- The person had an allergic reaction to the venom.
- The snake bite compounded an existing health condition in the person.

## Prevention

The following steps will help prevent snakebites:

- Do not aggravate a snake.
- Wear proper footwear and watch where you put your feet when hiking.



If you hear a rattlesnake, remain still until the snake is located, then back away slowly: avoid stomping or jumping as this may startle the snake.

## A Snake's Striking Range

A snake's striking range is about two-thirds of its length forward and one-third upward. (In water, snakes have a shorter striking range.) If you are within striking distance, slowly back out of range.

## What to Look For

Signs and symptoms of a possibly venomous snakebite include:

- A pair of puncture wounds in the skin
- Localized redness of the skin
- Pain and swelling in the area of the bite

## What to Do



### Call

Call EMS/9-1-1 and get an AED. If you are not sure whether the snakebite was caused by a venomous snake, call EMS/9-1-1 anyway. Do not wait for life-threatening signs and symptoms of poisoning to appear.



### Care

Before providing care, ensure that the snake is no longer present. If you see the snake, remember what it looks like so that you can describe it to EMS personnel. This information will help them provide the most appropriate treatment. Never attempt to capture or handle a potentially venomous snake.

1. Keep the injured site still and level with the heart, if possible.
2. If the bite is on a limb, remove any jewellery or tight clothing from the limb and watch for swelling.
3. Wash the wound with water.
4. Cover the bite with a clean, dry dressing.

## VENOMOUS SNAKES IN CANADA

### Northern Pacific Rattlesnake



Found in British Columbia

### Massasauga Rattlesnake



Found in Ontario

### Prairie Rattlesnake



Found in Alberta and Saskatchewan

## Myth-Information

**Myth:** You can help to slow the spread of venom through the body by cutting the wound or applying suction, ice, electricity, or a tourniquet.

These measures are not effective for slowing the spread of venom. In fact, they are likely to cause pain and injury. Your time is better spent seeking medical attention as quickly as possible.

## Spider Bites

Very few spiders in Canada can cause serious illness or death. The bites of harmless spiders often cause reactions similar to those of bee stings (e.g., swelling, redness, and stinging or pain at the site). However, two types of venomous spiders are occasionally encountered in Canada: the brown recluse and the black widow. The bite of either spider can, in rare cases, be life-threatening.



### Black Widow Spiders

The black widow spider is black with a reddish hourglass shape on the underside of its body. Its bite is the most painful and deadly of all the widow spiders, especially for very young children and older adults. The bite usually causes an immediate, sharp, "pin-prick" pain, followed by dull pain in

the area of the bite. However, the person often does not know that he or she has been bitten until he or she starts to feel ill or notices a bite mark or swelling.

The signs and symptoms of a black widow spider bite (e.g., abdominal cramps), can mimic those of other medical emergencies. Anyone with a suspected bite should be seen by a doctor for correct diagnosis and treatment.



### Brown Recluse Spiders

The brown recluse spider has a distinctive violin-shaped pattern on the back of its front body section. At first, the bite may produce little or no pain. Pain in the area of the bite develops an hour or more later. A blood-filled blister forms under the surface of the skin, sometimes in a target or bull's-eye pattern. Over several hours, the blister increases in size and eventually ruptures, leading to tissue destruction and a black scab.

### What to Look For

Signs and symptoms of spider bites depend on the amount of venom injected and the person's sensitivity to the venom. Most spider bites heal with no adverse effects or scarring. Signs and symptoms of venomous spider bites can seem identical to those of other conditions and therefore can be difficult to recognize. The only way for a First Aider to be certain that a venomous spider has bitten a person is to have witnessed it.

### Black Widow Spider Bite

Signs and symptoms of a black widow spider bite include the following:

- A raised, round, red mark
- Cramping pain in the thighs, shoulders, back, and abdominal muscles
- Restlessness and anxiety
- Dizziness
- Headache
- Excessive sweating
- Weakness

## Brown Recluse Spider Bite

Signs and symptoms of a brown recluse spider bite include the following:

- A slight stinging sensation (though bites may not be initially felt)
- A blood-filled blister that appears within 2 to 8 hours
- A bull's-eye pattern around the bite

Signs and symptoms of a severe reaction to a brown recluse spider bite occur within 72 hours of the bite and include nausea, vomiting, and joint pain.

### What to Do



#### Call

Call EMS/9-1-1 only if you suspect the person has been bitten by a black widow or brown recluse spider.



#### Care

To care for a bite from a non-venomous spider:

1. Wash the area with water.
2. Suggest that the person apply an antibiotic ointment to the wound if the person has no known allergies or sensitivities.
3. Apply a cold pack wrapped in a thin, dry towel to help reduce pain and swelling.

To care for a spider bite from a black widow spider or a brown recluse spider:

1. Wash the area with water.
2. Apply a cold pack wrapped in a thin, dry towel.



## Stings from Marine Life

Many forms of marine life (e.g., jellyfish, stingrays, sea urchins, stinging coral, and spiny fish) cause stinging wounds. In Canada, stings from marine life are usually from jellyfish, though stingrays can also cause injury. Stings from marine life can have effects that range from merely painful to potentially life-threatening.

### What to Look For

The signs and symptoms of a sting from marine life include the following:

- Pain
- Rash
- Redness
- Swelling
- Puncture wounds
- Lacerations

## What to Do



### Call

Call EMS/9-1-1 and get an AED if the person is having airway or breathing problems, the person was stung on the face or neck, or you do not know what caused the sting.



### Care

To care for a person who has been stung by a jellyfish:

1. Get the person out of the water as soon as possible.
2. Flush the injured area with vinegar for at least 30 seconds to counteract the toxin. If vinegar is not available, mix baking soda and water into a paste and leave it on the area for 20 minutes.
3. While wearing gloves or using a towel, carefully remove any stingers, tentacles, or pieces of the animal.
4. Immerse the affected area in water as hot as the person can tolerate (no more than about 45°C (113°F)) for at least 20 minutes or until the pain is relieved.
5. If hot water is not available, use dry hot packs, or, as a second choice, dry cold packs to help decrease the pain. Remember to wrap the hot or cold packs in a thin, dry towel or cloth to protect the skin. Do not rub the area or apply a bandage.



If the person has been stung by a Portuguese man-of-war (also called a bluebottle jellyfish), flush the injured areas with ocean water instead of vinegar. Do not flush any jellyfish sting with fresh water, ammonia, or rubbing alcohol because these substances can increase the person's pain.

To care for a person who has been stung by a stingray, sea urchin, or spiny fish:

1. Get the person out of the water as soon as possible.
2. Flush the affected area with tap water. Ocean water may also be used.
3. Keep the injured part still and soak the affected area in water as hot as the person can tolerate for at least 20 minutes or until the pain is relieved.
4. Check with a healthcare provider to determine if a tetanus shot is necessary and monitor the wound for signs and symptoms of infection.



# 14 Childhood Illnesses

Caring for children, either at home or in a child care setting, involves special considerations. Children often have a higher risk of complications from illnesses and there are certain diseases that children are more susceptible to. Young children, especially, may not yet be vaccinated against diseases that most adults are immunized against. In general, children's immune systems are less developed and less able to resist infection.

Infections can spread quickly between children, as they don't always follow proper hygiene procedures (such as blowing their noses and washing their hands). Child care settings have special notification and disinfection procedures, so ensure that you are aware of the protocols that apply to your workplace and follow them carefully.

### Protecting Children from Infection

In a child care setting, protect children from infectious diseases by:

- Insisting that staff members who are sick not come in to work.
- Encouraging parents and guardians to keep sick children at home.
- Having an isolation room for children who unexpectedly become ill.
- Washing your hands before and after contact with a child who has diarrhea or is vomiting.
- Washing bedding and equipment at least once a week, or every day when children are sick.
- Teaching children the importance of covering their mouths (preferably with the inside of their elbow) when they cough or sneeze and washing their hands afterward.

In a child care setting, follow these guidelines for cleaning the bathroom:

- Clean and sanitize the bathroom sink, countertop, toilet, and floor with a bleach solution once a day.
- If you use the bathroom sink to clean toilet trainers, use a bleach solution to sanitize the sink afterward.

### Questions to Ask Yourself If You Think a Child May Be Ill

It can be more difficult to identify illness in a child than in an adult, because children may not be as good at communicating what is wrong. If you suspect that a child may be ill, ask yourself the questions in the following table. If you answer "yes" to any, it may be a sign of illness.

Area to Watch	Questions to Ask
<b>Behaviour</b>	<p>Is the child:</p> <ul style="list-style-type: none"> <li>• Confused?</li> <li>• Unusually sleepy?</li> <li>• Unusually irritable or fussy?</li> <li>• More active or more subdued than normal?</li> <li>• Not interested in other children or play?</li> <li>• Crying nonstop, even when cuddled?</li> </ul>
<b>Face</b>	<p>Does the child:</p> <ul style="list-style-type: none"> <li>• Appear pale or flushed?</li> <li>• Show signs of pain or anxiety?</li> <li>• Have bluish lips?</li> <li>• Have any swelling?</li> </ul>
<b>Skin</b>	<p>Does the child have:</p> <ul style="list-style-type: none"> <li>• Hot and dry skin or cold and moist skin?</li> <li>• A rash or spots?</li> <li>• An unusual skin colour?</li> <li>• Itchy skin?</li> <li>• Any bruising or swelling?</li> </ul>

Area to Watch	Questions to Ask	Area to Watch	Questions to Ask
<b>Eyes</b>	<p>Does the child:</p> <ul style="list-style-type: none"> <li>• Rub and scratch his or her eyes?</li> <li>• Have red and inflamed eyes?</li> <li>• Have discharge in his or her eyes?</li> <li>• Have dull or unusually bright eyes?</li> <li>• Have swollen or puffy eyes?</li> <li>• Have yellow eyes?</li> <li>• Complain of seeing spots?</li> </ul>	<b>Breathing</b>	<p>Does the child have:</p> <ul style="list-style-type: none"> <li>• Rapid shallow breathing?</li> <li>• Painful breathing?</li> <li>• A strange odour on his or her breath?</li> </ul>
<b>Ears</b>	<p>Does the child have:</p> <ul style="list-style-type: none"> <li>• Trouble hearing?</li> <li>• Swelling in or around the ears?</li> <li>• Ringing in the ears?</li> <li>• An earache?</li> <li>• Any discharge?</li> <li>• Loss of balance?</li> <li>• A tendency to pull, cup, or poke his or her ears?</li> </ul>	<b>Throat</b>	<p>Does the child have:</p> <ul style="list-style-type: none"> <li>• A sore throat?</li> <li>• Difficulty swallowing?</li> <li>• Unusual drooling?</li> <li>• A red and inflamed throat?</li> <li>• A voice that sounds different?</li> </ul>
<b>Tongue</b>	<p>Does the child have a:</p> <ul style="list-style-type: none"> <li>• Dry and cracked tongue?</li> <li>• Red and raw tongue?</li> <li>• White or yellow coating on his or her tongue?</li> </ul>	<b>Cough</b>	<p>Does the cough:</p> <ul style="list-style-type: none"> <li>• Occur frequently, and is it dry?</li> <li>• Bring up sputum?</li> <li>• Sound unusual?</li> </ul>
			<p>Does the child have:</p> <ul style="list-style-type: none"> <li>• Little or no appetite?</li> <li>• An unusual level of thirst?</li> </ul>

Area to Watch	Questions to Ask
<b>Vomiting</b>	<p>Is the child:</p> <ul style="list-style-type: none"> <li>• Unable to keep food or water down?</li> <li>• Nauseated?</li> <li>• Frequently vomiting?</li> <li>• Projectile vomiting?</li> </ul>
<b>Temperature</b>	<p>Does the child:</p> <ul style="list-style-type: none"> <li>• Complain about feeling very cold?</li> <li>• Complain about feeling very hot?</li> <li>• Shiver uncontrollably?</li> </ul>
<b>Bowel Movements</b>	<p>Are the child's bowel movements:</p> <ul style="list-style-type: none"> <li>• Abnormally frequent and liquid?</li> <li>• Abnormally infrequent, dry, and hard?</li> <li>• Abnormal in content, such as undigested food, mucus, or blood?</li> <li>• An unusual colour or odour?</li> </ul>

### ***When to Call the Parent or Guardian of a Child in Your Care***

Call the child's parent or guardian if:

- The child has a fever.
- The child has diarrhea more than twice in a day.
- The child has been vomiting.
- The child has an injury that requires medical attention.

For minor issues, such as a small cut, a change in behaviour (e.g., not playing as usual), or a change in appetite, you should inform the parent or guardian when he or she picks the child up.

### ***Common Childhood Conditions: Guidelines for Care***

Some childhood illnesses and conditions are contagious. Some conditions cause irritation and discomfort, while others can cause lifelong complications or even death. Preventative practices include proper hygiene (including routine hand washing) and limiting exposure to people who are ill. Babies who are 6 months or older are the most vulnerable to these diseases.

Vaccination is also a fundamental preventive strategy. Vaccines have been incredibly effective in preventing childhood diseases and decreasing child mortality rates. Common childhood vaccines include diphtheria, tetanus, pertussis, measles, mumps, and rubella.

Provincial/territorial legislation dictates which contagious diseases must be reported to the local health unit. For some contagious diseases, parents and guardians have a responsibility to ensure that the child care facility or school is informed, in addition to the local health unit.

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Eczema</b> <ul style="list-style-type: none"> <li>Not contagious</li> </ul> <p>Note: Eczema sores can become infected.</p>	<ul style="list-style-type: none"> <li>Inflamed skin</li> <li>One or a combination of rashes, pimples, scaly skin, and scabs</li> <li>Dry skin, or skin with a watery discharge</li> <li>Itchy or burning skin</li> </ul>	<ul style="list-style-type: none"> <li>Soften crusts and dry skin with cold cream or oil. This often relieves itching.</li> <li>Prepare a lukewarm bath (<math>39.6^{\circ}\text{C}</math> (<math>103^{\circ}\text{F}</math>)) for the child.</li> <li>Do not scrub the affected area with soap.</li> <li>Eczema can often be helped with medication prescribed by a doctor.</li> </ul>	<p>Eczema may be caused by stress or it may be an allergic reaction to something in the child's environment or diet.</p> <p>Eczema in babies is often related to milk allergies. This type of eczema does not respond to creams and does not heal easily.</p>
<b>Impetigo</b> <ul style="list-style-type: none"> <li>Bacterial infection</li> <li>Contagious</li> </ul>	<ul style="list-style-type: none"> <li>Inflamed skin</li> <li>Clusters of pimples filled with straw-coloured fluid that become crusted and break, found around the mouth and nose; may be flat and pitted</li> </ul>	<ul style="list-style-type: none"> <li>Wash the infected area frequently with soap and water.</li> <li>Doctors often prescribe antibiotic ointments or creams for the treatment of impetigo.</li> </ul>	<p>Impetigo can spread through contact with infected skin or items that have contacted infected skin (e.g., clothing, towels, and bed linens). Careful handwashing by both the child and the caregivers is the best prevention.</p> <p>A child who has had impetigo should not return to the child care setting or school until prescribed medication has been taken for at least one full day.</p>

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Ringworm</b> <ul style="list-style-type: none"><li>• Fungal infection</li><li>• Very contagious</li></ul>	<ul style="list-style-type: none"><li>• Red, scaling rings on the skin</li></ul>	<ul style="list-style-type: none"><li>• Clean the area twice a day with mild soap and water.</li><li>• Doctors often prescribe anti-fungal medication for ringworm, which should be applied after cleaning the affected area.</li></ul>	A child with ringworm is contagious until 48 hours after treatment begins, and so should not return to the child care setting or school before this time has passed.
<b>Scabies</b> <ul style="list-style-type: none"><li>• Skin parasite</li><li>• Very contagious</li><li>• This is a reportable condition.</li></ul>	<ul style="list-style-type: none"><li>• Many tiny blisters, scratch marks, and scaly crusts found mainly in skin folds (e.g., between fingers, on wrists, on the torso, and on genitals)</li><li>• Extreme itchiness</li></ul>	<ul style="list-style-type: none"><li>• Bathe the child prior to medication application.</li><li>• After the bath, apply the special lotion prescribed by the doctor.</li></ul>	<p>Scabies is a skin parasite called the "human itch mite."</p> <p>The scabies mite is passed by direct skin-to-skin contact, so everyone in the household should be treated at the same time.</p> <p>After treatment, wash all bed linen, underclothes, sleepwear, and blankets, or press them with a hot iron.</p> <p>A child who has scabies should not return to the child care setting or school until treatment has been completed.</p>

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Prickly Heat Rash</b> <ul style="list-style-type: none"> <li>• Not contagious</li> </ul>	<ul style="list-style-type: none"> <li>• A rash of tiny, pinpoint blisters surrounded by blotches of pink skin</li> <li>• A rash on the face or on the parts of the body that are most heavily clothed</li> <li>• Occurs in hot weather or whenever a child or baby is overdressed</li> </ul>	<ul style="list-style-type: none"> <li>• If the child or baby is feverish, give him or her a sponge bath.</li> <li>• Pat the skin dry to avoid rubbing the skin.</li> <li>• Dress the child or baby in layers, and remove layers as necessary.</li> <li>• Replace wet or sweaty clothing immediately.</li> </ul>	Prickly heat rash rarely bothers the child, but if there is a fever at the same time, the child may be irritable.
<b>Diaper Rash</b> <ul style="list-style-type: none"> <li>• May be contagious, depending on the underlying cause</li> </ul>	<ul style="list-style-type: none"> <li>• Red, scalded appearance that does not fade when left un-diapered and exposed to air</li> <li>• Pimples or sores found in the diaper area</li> </ul>	<ul style="list-style-type: none"> <li>• Wash and dry the area well.</li> <li>• Expose the area to the air.</li> <li>• Change diapers often.</li> <li>• Use a zinc-based cream.</li> <li>• If the rash does not clear up within 48 hours, or you are concerned about the underlying cause, seek medical attention.</li> </ul>	Some diaper rashes are yeast infections. Watch for whitish, curd-like deposits found in genital folds that are not easily wiped away. These may need to be treated with prescription ointment. A yeast infection can be contagious. Use proper precautions.

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Lice</b> <ul style="list-style-type: none"><li>• Contagious</li><li>• This is a reportable condition.</li></ul>	<ul style="list-style-type: none"><li>• Small white eggs (nits) on the scalp or in body hair (nits look like dandruff but cannot be washed off)</li><li>• Live lice (dark and slightly bigger than nits) on the scalp or in body hair</li><li>• Pustules and scabs on the scalp due to scratching</li></ul>	<ul style="list-style-type: none"><li>• Wash the child's hair with the medicated shampoo recommended by a doctor, then comb it with a fine-toothed comb to remove nits.</li><li>• Disinfect all clothing and bedding by having them dry-cleaned, washing them in very hot water, or freezing them for 72 hours. Items such as stuffed animals should also be disinfected, and then sealed in a plastic bag for 3 weeks.</li><li>• The whole family may need to be treated.</li><li>• Vacuum the environment regularly during the treatment period.</li></ul>	<p>Lice are very common among children because they play closely together. However, anyone can get lice. Lice is not caused by a lack of personal cleanliness.</p> <p>Always store children's headgear separately and do not allow children to share brushes and combs.</p> <p>Children may return to the child care setting or school after the first treatment.</p>

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Pink Eye (Conjunctivitis)</b> <ul style="list-style-type: none"> <li>• Bacterial infection</li> <li>• Contagious</li> </ul>	<ul style="list-style-type: none"> <li>• Pink colouration of the white of the eye</li> <li>• Swollen eyelids</li> <li>• Pus on eyelids</li> <li>• Itchy or sensitive eyes</li> </ul>	<ul style="list-style-type: none"> <li>• Seek medical attention: doctors often prescribe antibiotics.</li> <li>• Protect the eyes from bright light.</li> <li>• Do not let children share towels or washcloths.</li> <li>• The child should not rub or touch his or her eyes.</li> <li>• Use warm water compresses to remove pus.</li> </ul>	<p>Pink eye can be bacterial or viral, is transmitted through direct contact, and is very contagious.</p> <p>If the child's eyes have pus, keep the child home from the child care setting or school for 24 hours.</p>
<b>Pinworms</b> <ul style="list-style-type: none"> <li>• Very contagious</li> </ul>	<ul style="list-style-type: none"> <li>• Constant scratching around the anus</li> <li>• Unusual irritability</li> <li>• Restlessness or inability to sleep at night</li> </ul>	<ul style="list-style-type: none"> <li>• Wash the child's hands and nails (using a nail brush).</li> <li>• Keep the child's fingernails short and discourage nail biting.</li> <li>• Make sure the child has a daily bath or shower and a daily change of towels, sheets, and underwear.</li> <li>• Doctors often prescribe oral medication.</li> <li>• The whole family must be treated.</li> </ul>	<p>Pinworms are intestinal parasitic worms. Tiny eggs deposited around the anus by a female worm spread the infection.</p> <p>Keep the child home from the child care setting or school until after starting medication.</p>

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Chickenpox</b> <ul style="list-style-type: none"> <li>• Viral infection</li> <li>• Contagious</li> </ul>	<ul style="list-style-type: none"> <li>• A rash of tiny red spots covering the skin</li> <li>• Fluid-filled blisters and scabbing on the skin</li> <li>• Fever</li> <li>• Itchiness</li> </ul>	<ul style="list-style-type: none"> <li>• Try to make sure the child does not scratch.</li> <li>• Give fever-reducing medications as recommended by a doctor.</li> <li>• The doctor may also recommend a cream to relieve itching.</li> </ul>	<p>Chickenpox is most contagious 1 to 2 days before the rash appears, and for approximately 5 days after onset, or until the lesions have become crusted.</p> <p>Chickenpox is transmitted by direct contact with blisters or body fluids, or through coughing and sneezing.</p> <p>The child may have to stay home until lesions become crusted, depending on the protocols of the school or child care centre.</p>
<b>Diarrhea</b> <ul style="list-style-type: none"> <li>• Usually a symptom of an underlying condition, which may or may not be contagious</li> </ul>	<ul style="list-style-type: none"> <li>• Unusually frequent and liquid bowel movements</li> <li>• Pain or bloating in the abdomen</li> <li>• Dehydration</li> </ul>	<ul style="list-style-type: none"> <li>• Seek medical attention if there is vomiting or blood in the stool, or if diarrhea persists for more than 72 hours.</li> <li>• Disinfect diaper-change areas and toilet areas very carefully.</li> <li>• Isolate the child.</li> <li>• Give the child plenty of clear fluids for 24 hours.</li> <li>• If the child has two or more episodes of diarrhea, give fluids with electrolytes (e.g., sports drinks or coconut water).</li> </ul>	<p>Diarrhea may be a symptom of an infection. Food poisoning or a change in diet can also cause diarrhea.</p> <p>For child care workers and babysitters: Call the parent or guardian if a child has diarrhea more than twice during the day.</p>

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Dehydration</b> <ul style="list-style-type: none"><li>• Not contagious</li></ul>	<ul style="list-style-type: none"><li>• Dry mouth and tongue</li><li>• No tears when crying</li><li>• Sunken eyes</li><li>• Less frequent urination or urine that is darker than usual</li><li>• In babies, the soft spot on the top of the head is sunken</li></ul>	<ul style="list-style-type: none"><li>• Seek medical attention.</li><li>• Continue to give fluids.</li></ul>	<p>Dehydration results from a loss of bodily fluids.</p> <p>Dehydration in children is most commonly caused by prolonged vomiting and diarrhea. It can be life-threatening.</p>
<b>Earaches</b> <ul style="list-style-type: none"><li>• Not contagious</li></ul>	<ul style="list-style-type: none"><li>• Fever or chills</li><li>• Deafness</li><li>• Dizziness</li><li>• Nausea</li><li>• Fluid from the ear</li><li>• A worried appearance</li><li>• Pulling at the ear or covering it with the hand</li><li>• Pain</li></ul>	<ul style="list-style-type: none"><li>• Seek medical attention.</li><li>• The doctor may recommend an antibiotic.</li></ul>	<p>Never put anything into the ear canal (e.g., a cotton swab), even if fluid or discharge appears.</p>

## CHILDHOOD CONDITIONS

Conditions	What to Look For	What to Do	Remember
<b>Persistent Vomiting</b> <ul style="list-style-type: none"> <li>Usually a symptom of an underlying condition, which may or may not be contagious</li> </ul>	<ul style="list-style-type: none"> <li>Nausea</li> <li>Stomach pain</li> </ul>	<ul style="list-style-type: none"> <li>Give the child plenty of fluids.</li> <li>When the vomiting subsides, give small amounts of easily digestible food, such as toast.</li> <li>If the vomiting continues for more than 24 hours, signs of dehydration appear, or the vomiting is associated with diarrhea or pain, seek medical attention.</li> </ul>	Persistent vomiting could be caused by illness, emotional upset, food poisoning, or a reaction to medication or a new food.

Note: Although these conditions are most common in children, adults who are in regular contact with them (e.g., teachers) should check themselves often and follow the same care steps outlined above.

### Comparison Chart for Epiglottitis and Croup

Both epiglottitis and croup affect the throat, and can be difficult to distinguish.

**Epiglottitis** is a bacterial infection that causes severe inflammation of the epiglottis. The epiglottis is a flap of tissue above the vocal cords that protects the airway during swallowing. When it becomes infected, it can swell until the airway is completely obstructed. A child with epiglottitis may:

- Appear severely ill and have a high fever.
- Need to be sitting up.
- Strain to breathe.
- Appear to be very frightened.
- Drool from the mouth, as swelling may prevent swallowing.

**Croup** is a respiratory condition, usually triggered by an acute viral infection of the upper airways. The infection causes swelling of the throat and of the tissues below the vocal cords. Croup is generally non-life-threatening, but it can lead to severe shortness of breath and hypoxia. Croup is commonly identified by its distinctive harsh, barking cough, often described as being like the bark of a seal. Croup is often preceded by 1 or 2 days of illness, sometimes with a fever. Croup occurs more often in the winter months and the signs and symptoms of croup are often more evident in the evening. The child may show improvement when exposed to cool air.

## THE DIFFERENCES BETWEEN EPIGLOTTITIS AND CROUP

	<b>Epiglottitis</b>	<b>Croup</b>
<b>What Is It?</b>	Inflammation and swelling of the epiglottis. When it strikes, it is life-threatening.	Inflammation of the throat and vocal cords
<b>Cause</b>	Bacterial infection	Viral infection
<b>Ages Most Likely to Be Affected</b>	3 to 7 years old	6 months to 5 years old Called laryngitis among children 5 years and older
<b>Onset</b>	Comes on suddenly, from 6 to 24 hours after infection	Develops more gradually, from 24 to 72 hours after infection
<b>Child's Appearance</b>	Looks very ill and anxious	Looks only mildly or moderately ill
<b>Drooling</b>	Frequent: child has trouble swallowing saliva	No drooling
<b>Hoarseness</b>	Not usually	Very marked hoarseness
<b>Coughing</b>	Rare	Frequent, harsh "crowding" cough
<b>What to Do</b>	Call EMS/9-1-1 immediately. Report the child's condition to the parent or guardian. Note: Do not inspect the mouth and/or throat, as this may increase irritation.	Call EMS/9-1-1 immediately if the child shows signs of respiratory distress. The child may show improvement when exposed to cool air. Report the child's condition to the parent or guardian.

## Childhood Fevers

A fever is one of the body's defense mechanisms. A mild fever should not be a concern unless it continues for more than 3 days. If the child's temperature rises to 39°C (102°F) or is not easily controlled, seek medical attention.

A fever alone is not a clear indicator of how sick a child is. A child may be very sick and not have a fever, or may be running a fever and playing happily. The best way to gauge the severity of an illness is to note changes in the child's behaviour.



### Taking a Child's Temperature

Normal body temperature is 37°C (98.6°F). Anything higher than this is considered a fever. To determine a child's temperature you must use a thermometer. Many types of digital thermometers are available, including those intended for use in the mouth and ear. Follow these general principles when taking a child's temperature, along with any specific manufacturer's instructions for your thermometer.

## What to Do

1. Wash your hands thoroughly.
2. If taking the temperature orally, make sure that the child has not had anything hot or cold to eat or drink in the previous 10 minutes.
3. Clean and reset the thermometer as per the manufacturer's guidelines.
4. Place the thermometer in the child's mouth (under the tongue) or in the child's ear.
5. Leave the thermometer in place until it beeps, or for up to 1 minute (if your model does not have an audio notification).
6. If you are using a thermometer for more than one child (e.g., in a child care setting), use protective covers and throw them away after every use.
7. Record the temperature, the time, and the method of taking the child's temperature (e.g., "oral").
8. Clean the thermometer after every use as instructed by the manufacturer.



Do not take the temperature rectally as this can damage the bowel lining.

## Reducing a Child's Temperature:

1. Children with fevers should be dressed lightly because they may feel warm. If they get too cold or start to shiver, cover them until the skin feels warm when you touch it.
2. Sponge the child with room temperature water.
3. Give the child plenty of fluids to ensure that he or she stays hydrated.
4. If the fever rises above 39°C (102°F) or persists for more than 3 days, seek medical attention.
5. Give fever-reducing medications, as discussed with the healthcare practitioner.

## Complications Related to Fever

### Reye's Syndrome

A child or teenager with a viral infection, such as chickenpox or influenza, may develop Reye's syndrome if given ASA (e.g., Aspirin®). Reye's syndrome is a disease of the brain and liver that can lead to death. Never give ASA to a child or teenager with a viral infection without first consulting a healthcare practitioner.

### Seizure

A child or baby may suffer a seizure during a high fever. If this occurs, follow the care steps for febrile seizures.

## Giving Medication

### When to Give Medications

Give medication to a child or baby in your care:

- Only if you have written permission from a parent or guardian for each medication that is to be given
- Only when the medication is in the original container with the original label
- Only when the medication is properly labelled with:
  - The child's name.
  - The date the prescription was issued.
  - The instructions.
  - The time period during which the medication is to be taken.

If a child is being given non-prescription medication, ensure you have written permission from a parent or guardian for each medication that is to be given.



Parents or guardians should give the first dose of a new medication so that they can watch for any unusual signs, symptoms, reactions, or behaviours in their children or babies.



## General Rules for Medication

To reduce the risk of miscommunication and error, it is a good idea to have just one person in the child care setting who is responsible for giving all medications. Store all medication out of the reach of children.

### When giving medication:

- Wash your hands thoroughly.
- Check the medication three times:
  - When you take the medication out from the cupboard or refrigerator
  - When you take the medication out of the package
  - Just before you administer the medication
- Keep a written record of the date and time the medication was given, the name of the child, the name of the medication, the person who gave the medication, and the amount given. Record all this information when you give the medication.
- Call the parent or guardian if the child cannot keep the medication down.
- Report to the parent, guardian, or doctor any reactions or effects not usually experienced by the child.
- At the end of each day, tell the parent or guardian what medication was given and at what time.
- Follow any specific directions for the medication (e.g., "do not take with food").
- When applying topical medication, protect yourself by using an applicator or by wearing disposable gloves.
- Emergency medicine, such as inhalers or epinephrine auto-injectors, should be accessible at all times but out of the reach of children.



Some provinces and territories have legislation regarding medication administration in child care settings. Always follow local protocols.

## How to Give Specific Medications

### Medication in the Eye

1. Have the child look up while in a lying or sitting position.
2. Gently pull down the lower eyelid.
3. Hold the dropper horizontally about one inch out from the child's eye.
4. Drop the medication into the little pocket between the lower lid and the eyeball.
5. Ask the child to close the eye then briefly hold a cotton ball against the inside corner of the eye. Use a separate cotton ball for each eye.
6. Wipe away any excess medication.

### Medication in the Nose

1. Have the child lie on a flat surface with his or her head hanging over one edge. If the child must sit up, have the child tilt his or her head back as far as possible.
2. Place the exact number of drops indicated in the prescription into each nostril.
3. Have the child remain with his or her head back for a few minutes.



# 15 Caring for Children

Children are very vulnerable. Their bodies are more susceptible to injury, and they are typically less aware of risks and hazards in their environments. When caring for children, prevention, preparation, and constant supervision are critical to preventing injury.

## Toy and Equipment Safety

Products designed for children have many features to keep them safe, but these products must be selected and used properly.

You should always discard broken equipment and toys immediately, and always follow the manufacturer's directions for safe use. When considering second-hand equipment or toys, first check the label and contact Health Canada to find out if there has been a recall or safety alert on that specific model.

## Equipment for Babies

### Safe Use of Cribs

- Cribs manufactured before September 1986 do not meet current safety standards and should be disassembled and either thrown away or taken to your local recycling centre.
- Manufacturers of cribs are required to include a label showing the date of manufacture. If there is no label, assume that the crib is not safe to use.
- Allow only one small toy inside the crib and remove all mobiles or activity centres as soon as the baby can pull him- or herself up.
- Place the crib well away from potential hazards such as blind cords, windows, pictures, and shelves.
- When a child shows signs of trying to climb out of a crib, it is time to move the child to a bed or mattress on the floor.
- Keep the wheels on a crib locked whenever it is in use.
- Make sure that the crib slats are no more than 6 cm (2 3/8 in.) apart.

- Make sure that the corner posts extend no more than 3 cm (1 1/4 in.) above the headboard and footboard.
- The height from the mattress support to the top rail must be at least 66 cm (26 in.) when the support is in its lowest position.
- Make sure the mattress fits snugly inside the crib. Use only mattresses sold for use in cribs. Any other substitutes, such as foam pads, may allow for pockets or spaces in which a child may get stuck or be smothered.

### Safe Use of Playpens

- Manufacturers of playpens are required to include a Canadian Standards Association (CSA) label showing that the playpen has been approved by CSA. The label also shows the playpen's date of manufacture. Any playpen manufactured before 1985 has not been approved by the CSA. If there is no label, assume that the playpen is not safe to use.
- If a child shows signs of trying to climb out of a playpen, the playpen is no longer safe for that child.
- Ensure that the playpen's wheels are locked whenever it is in use.
- Make sure that playpen walls are sturdy and at least 48 cm (19 in.) high.
- Inspect the playpen regularly to ensure that there are no hazards such as rips or exposed sharp edges.

### Safe Use of High Chairs

- Do not allow older children to climb on the chair, even when it is unoccupied.
- A safe chair is stable and has a wide base to reduce the risk of tipping.
- The harness should consist of a strap that fits between the child's legs and a waist belt that is easy to fasten and is in good condition.

- Ensure that the child's hands, arms, and legs are clear of any moving parts before making adjustments to the chair or the tray.
- Keep the chair a safe distance away from hazards such as windows, blind cords, and mirrors, or surfaces that a child could push off of such as walls, doors, or furniture.

### Safe Use of Baby Strollers

- Manufacturers of baby strollers are required to include a Canadian Standards Association (CSA) label showing that the baby stroller has been approved by CSA. The label also shows the stroller's date of manufacture. Any stroller manufactured before 1985 has not been approved by the CSA. If there is no label, assume that the stroller is not safe to use.
- Do not load heavy parcels on the back or on the handles of the baby stroller. These could cause the stroller to tip backward.
- Use a stroller with safety straps that go around the child's waist and between the legs.
- Keep children's fingers away from moving parts.

### Safe Use of Pacifiers

- When purchasing a pacifier, make sure it is from a reputable manufacturer.
- Check pacifiers regularly for wear and tear. Discard them immediately if there is any sign of deterioration, such as discolouration, hardening, or cracking of the nipple.
- If a doctor diagnoses a child or baby with a yeast infection of the mouth, immediately discard any pacifier the baby is using. Give the baby a new one only after the infection has cleared.
- Use a pacifier clip, as opposed to a cord, to secure the pacifier to the child.
- A teething ring should be used instead of a pacifier once the baby's teeth start to appear.

## Equipment for Children

### Safe Use of Child Bike Trailers

- Child bike trailers are recommended over child bike carriers because they reduce the risk of falls, do not cause as many steering and balance problems, and increase protection from the weather. They are also farther from the bicycle wheels, ensuring that the child cannot reach the spokes.
- If possible, choose a bike trailer that has a five-point harness and roll bar.
- Attach a tall, bright flag to the trailer.
- Connect the trailer with a flexible joint so that it will not tip over, even if the bike does.
- As a bike trailer is wider than a bike, ensure that the wheels do not go over the road edge.
- Ride in low-traffic areas.
- Ensure that children wear Canadian Standards Association (CSA) approved and properly fitting helmets.

### Safe Use of Safety Gates

- Use only approved safety gates. Avoid makeshift alternatives, such as pieces of furniture.
- Manufacturers of safety gates are required to include a label showing the date of manufacture. If there is no label, assume that it is not safe to use.
- If you can fit a pop can between the railings, a child could get stuck and the gate is not safe.
- Install safety gates wherever stairs are exposed (at the top and bottom of the staircase). Make sure they are properly fitted and secure.
- If the gate has a pressure bar, make sure it is on the side away from children.

- Always open and close gates to pass through. Climbing over a gate is unsafe and may cause an injury. It also models unsafe behaviour to children.
- Safety gates are not a substitute for constant supervision. Always monitor children closely, even if a safety gate is in place.

## Toys for Children and Babies

The following are guidelines to keep children safe when playing with toys:

- Follow the manufacturer's directions for safe use, including suggested age level.
- Choose toys that are appropriate for the children who will be using them. The Canadian Toy Testing Council issues a list of guidelines for choosing toys.
- Toys must be cleaned daily in child care centres. To clean toys and equipment, use a solution of 6 mL (1 tsp.) of bleach to 400 mL (1½ cups) of water. Label the bottle and make a fresh solution every day.
- When there are older children in the same space, their toys may have smaller parts: These toys should be separated and stored out of common play areas.
- Avoid toys with sharp edges.
- Paints, crayons, and markers must be completely washable and non-toxic.
- Ensure that toys (and all of their detachable pieces) are too big to fit into the baby's or child's mouth, ears, or nose.
- Provide only toys without strings because strings may cause strangulation.
- Inspect toys regularly. Discard any pieces that are loose. If the toy seems unsafe, discard it immediately.
- Wooden toys should be glued or screwed together instead of nailed.
- Ride-around or ride-on toys must be selected according to the size of the child.
- Children should wear all safety equipment that is recommended for the activity. "Toy" protective gear (e.g., plastic helmets) is for imaginative play only.

- Electrical toys must bear the Canadian Standards Association (CSA) label.
- Choose costumes that are made of flame-retardant materials. Use face paints instead of face masks to avoid blocking vision.
- Large toy boxes and other containers must have air holes and self-supporting hinges if they have lids. This is necessary in case children decide to hide inside.

## Playground Safety

Playgrounds are the site of many serious injuries, mainly falls. Some playground equipment can be unsafe or inappropriate for certain age groups. The following points help to keep children safe on the playground:

- Examine park and school playgrounds for hazards such as broken glass or exposed sharp edges.
- Ensure that children use playground equipment as intended (e.g., ensure that they do not climb on the outside of a tubular slide).
- Ensure that there are enough adults present to watch all children carefully.
- If you see unsafe or broken equipment in a playground, report this to the managing agency (e.g., community centre, park board, school board).
- If you are installing or maintaining playground equipment, consider the age and size of the children who will be using it, and follow the manufacturer's directions.
- Ideally, sandboxes should be fitted with a cover to keep dogs, cats, and other animals from using it as a litter box.



Provincial/territorial legislation specifies regulations for child care centre play equipment.

## Car Safety

Motor vehicle injuries are the number one cause of death of children in Canada. Follow the guidelines on the safe use of car seats and the passenger rules for children to help ensure children's safety in and around motor vehicles.

### Safe Use of Car Seats

The law requires that all children use appropriate and approved child-restraint systems, such as child or baby seats and booster seats. These must have a label stating that they conform to the Canada Motor Vehicle Safety Standards (CMVSS), administered by Transport Canada.

Use caution when buying or using a pre-owned child car seat. Buy the seat from someone you know, and check it carefully. Make sure the car seat has:

- Instructions and all necessary hardware
- Not been in a collision
- A CMVSS sticker and is less than 10 years old
- No discoloured (stress) marks or cracks
- No signs of wear or tears on the harness

Always follow the manufacturer's instructions for installation and use.

When driving with children who do not require a car seat, remember:

- Children under the age of 13 should ride in the back seat.
- If you absolutely must put a child in the front seat of a car that has an airbag, push the seat back as far as it will go.

Make sure that any child who is no longer in a car seat is secured with a seat belt. Show the child how to wear the belt. The lap belt should be low across the hips, and the shoulder belt should be over the shoulder and across the chest. If the seat belt does not fit properly, a booster seat should be used.



It is against the law to leave a child or baby alone in a car, even for a short time.

### Passenger Rules for Children: Safety Education

Teach children the following rules for safety in and around motor vehicles:

- Always enter and exit the car on the curb side.
- Remain seated with your seat belt on while the car is moving. Stay that way until the driver has reached your destination and the car is completely turned off or the driver tells you it is safe to remove your seat belt.
- Remember that the driver needs to be able to concentrate on driving, so minimize distractions inside the car.
- Keep your head, arms, and feet in the car at all times.
- Remember that the car's controls are for the driver only. They are not for playing with, even when the car is parked.



As a caregiver, ensure that everyone's fingers, hands, and feet are clear of doors before the doors are closed. Ensure that any person driving children has adequate insurance coverage.

## Safe Use of Wheeled Equipment

### Choosing a Safe Bicycle

When choosing a bicycle for a child, ensure that it is the proper size:

1. Have the child stand over the bicycle seat with both feet flat on the ground.
2. Lift the bicycle up to the child's body (both tires off the floor). There should be at least 2.5 cm (1 in.) between the tires and the floor. Adjust the height of the seat as needed.

While on the seat, the child should be able to put the balls of his or her feet on the ground.

Regularly check the following for any bicycle used by a child:

- The brakes must work properly, and the child must be able to use them safely.
- The tires must be fully inflated.
- The wheels must be tightly secured and truly round.
- The chain must be oiled and tight.
- The spokes must be tight, straight, and unbroken.
- The seat post and handlebars must be securely attached.
- The bike must have a working bell or horn.
- The bicycle must pass the "bounce test:" Bounce the bicycle up and down on its wheels on a paved surface and listen for shakes and rattles. A safe bike is a quiet bike.



Local legislation may dictate what needs to be on your bike for safety (e.g., bike lights).

### Choosing a Safe Helmet

Mishaps during sports that require a helmet are the leading cause of head injuries for children in Canada. Wearing helmets can reduce the risk of a serious or fatal head injury by up to 80%.

Keep the following points in mind when choosing a helmet for a child:

- Use helmets approved by the Canadian Standards Association (CSA), Snell Memorial Foundation, or American National Standards Institute (ANSI).
- A helmet should fit squarely on the child's head and have a snug chin strap.

If a helmet is involved in a fall, buy another one. Even if no damage is visible, the helmet may have lost some of its ability to absorb shock.



As a role model and for safety reasons, adults should always wear a helmet while riding a bicycle.

### Safe Use of Bicycles

Ensure that children follow these rules when using a bicycle:

- Walk bicycles across roads only at intersections and pedestrian crosswalks.
- Before you cross, make sure that traffic crossing your path is fully stopped at stop signs or traffic signals.
- Ride in a straight line when you go down a road or street. Do not swerve between parked cars or onto driveways. Always look and signal in plenty of time before turning.
- Wear bright clothes even during the day. When motorists can see a cyclist, it is easier for them to co-operate with the cyclist.
- Do not ride a bicycle after dark.



Even the safest cyclist must watch for drivers who aren't paying attention!



It is important to teach children balancing, turning, hand signalling, and stopping on a bicycle before they go out riding. Practice these skills in the yard, a school yard, a park, or other low-traffic area.

## Safe Use of Skateboards and Rollerblades

To keep children safe while skateboarding and rollerblading:

- Restrict skateboarding and in-line skating to children over 5 years of age.
- Allow children to use only playgrounds and paths built for skateboarding and in-line skating.
- Insist that children wear the correct equipment: a bicycle helmet approved by the Canadian Standards Association (CSA), Snell Memorial Foundation, or American National Standards Institute (ANSI), as well as elbow, wrist, and knee pads.
- Ensure that children wear bright clothing while skateboarding or in-line skating. This will make the children more visible and reduce the risk of collisions.

## Fire Safety

### Reducing Risk of Fires and Burns

The following tips can reduce the risk of fires and burns:

- Keep children away from heat sources and appliances such as cooking ovens, barbecues, space heaters, wood stoves, and fireplaces.
- Make sure all fireplaces and wood stoves are cleaned and inspected regularly to follow safety codes.
- Keep space heaters at least 1 metre (3 ft.) away from flammable objects such as curtains and furniture.
- Turn pot handles in toward the centre of the stove.
- Keep cooking areas free of clutter. This will reduce the risk of igniting pot holders, aprons, and other kitchen items.
- Snuff out candles immediately after use.
- Keep all hot drinks out of children's reach.
- For nightwear, use only clothing designated as sleepwear. Other materials are sometimes more flammable. The government sets guidelines as to what is classified as sleepwear.
- Contact your local fire department for information on inspecting your home or workplace (including child care settings) for fire hazards and reviewing your fire escape plan.
- Make sure that smoke detectors are placed in all areas required by your local fire department or whichever agency regulates fire safety in your area. Change the batteries in all smoke detectors twice a year.
- Keep bedroom and sleeping-area doors closed when children are sleeping. Doors are important barriers to smoke.
- Make sure that all breakers and fuses on the electrical panel in your home, workplace, or child care setting are clearly labelled.
- Use only single-cord plugs, as multiple-cord or octopus plugs may overheat. Keep loose cords secured and out of the way to prevent tripping and to keep children from pulling them out.
- Keep an approved, all-purpose chemical fire extinguisher in the kitchen.

## Fire Escape Plan

Draw a floor plan. Mark the normal exit from each room. Then mark an emergency exit, such as a window. This exit can be used if fire blocks the doorway.

Identify a location where everyone will meet if they must escape from a fire. Decide who will assist those who cannot get themselves out, such as young children or people with difficulty walking.

In professional settings, legislation may determine how often this escape plan must be practiced.

## What to Do in the Case of Fire During the Night

1. When you hear the smoke detector, roll out of bed and onto the floor. Shout "Fire, fire, fire!" Be sure to shout it 3 times. Shout "Get out and stay out!" Teach children to do this as well.
2. Crawl to the door and touch it. If it is warm, do not open it. Unlock the door so that the firefighters can open it. Seal off any cracks with blankets or towels. Wet the blankets or towels if possible.
3. Escape via the emergency exit.
  - If you cannot get out, take refuge in a room with a window. Open it for ventilation. Hang a sheet out the window as a signal for help.
4. Join everyone at the special meeting spot.
5. After everyone is together, go to a neighbour's home to call the fire department.
6. Meet the fire trucks and tell the firefighters that everyone has escaped the burning building.

## Ice Safety

### Safe Skating or Walking on Ice

Teach children these safety rules for skating or walking on ice:

- Ice must be a minimum of 15 cm (6 in.) thick before it can support one person. Ice must be solid, clear blue, and not covered with snow.
- Always skate with a buddy.
- If you cannot be sure how thick a section of ice is, do not walk or skate on it.

Ensure that children know what to do if they fall through the ice:

- In shallow water, feel for the bottom with your feet. Then walk toward the shore or bank, breaking ice as you go. Keep going until you can slide up onto thicker ice.
- In deep water, flutter-kick your feet and extend your hands forward along the ice until you can slide up onto it. Slowly crawl to safety.
- After you are out of danger, get to shelter and change into warm clothing. This will help prevent hypothermia.

## Water Safety

### General Tips for the Prevention of Drowning

- Pay constant attention to babies or children when they are in, on, or around water.
- Turn wading pools upside down when not in use.
- Empty buckets of water immediately after use.
- Keep toilet lids down and keep the bathroom door closed.

## Safe Use of Bathtubs

- Always be present when a child or baby is in a bathtub or bathing basin. Drowning can occur even in a few centimetres of water.
- Check bathwater temperature by splashing water on your inner wrist. If it feels too hot or cold to you, do not place the child or baby in the bath until you have adjusted the temperature.
- If your bathtub is a Jacuzzi-type or is equipped with water jets, do not turn these on with the child or baby in the tub.
- Turn off all water taps tightly.
- Place a non-slip mat in the bathtub.

## Safe Use of Backyard Pools and Hot Tubs

- Keep children under constant supervision when they are playing in or around a backyard pool or hot tub.
- Make sure basic lifesaving equipment is available at all times. Examples of basic lifesaving equipment are: (1) a strong, lightweight pole with blunt ends; and (2) a ring buoy with a long throwing rope. Be aware of equipment requirements outlined in the legislation that applies to your situation.
- Cover hot tubs securely and lock them when they are not in use.
- Keep decks around the pool or hot tub clean and free of debris.
- Pay close attention to the surface of a diving board. Make sure the slip-resistant surface is always in good repair. Ensure that the water is deep in the entire area that divers use (mark the slope of the pool to avoid hitting the upslope).
- Install a rope with buoyant markers across the pool where the deep end slope begins.
- Use only unbreakable cups, dishes, and other utensils at the poolside.
- Keep electrical appliances away from backyard pools.
- Remove all floating toys and store them in a secure area when pool time is over.

- Make sure electrical equipment used to operate the pool or hot tub conforms to electrical code requirements.
- Have a phone available on the pool deck. Post a list of emergency numbers beside it.
- Ensure that an outdoor pool is surrounded by a fence or wall that children cannot get through or over. Gates should have self-closing latches above the reach of toddlers. They should also include hardware for permanent locking. Obey the fencing requirements of your local municipality.
- Make sure all pool chemicals are stored in a secure area.

## Safe Swimming

- Make sure that children and adults learn to swim. For swimming lessons, take a Red Cross Swim course.
- Ensure children are always supervised by a responsible adult.
- Establish sensible safety rules before children do any swimming. Enforce these rules consistently and firmly.
- Supervise even strong swimmers at all times. Sudden cramps can sink the most accomplished swimmer.
- Allow swimming only during daylight hours and in good weather.
- Remind non-swimmers and swimming supervisors of the dangers of relying on inflatable toys (e.g., inner tubes). Inflatable toys can leak and no longer provide support, and they can also create barriers to effective supervision.
- Ensure children stay within specified boundaries.
- When distance swimming at the beach, make sure that children swim parallel to the shore.
- Always know the depth of the water and ensure it is clear of all obstacles before diving.
- Post "No Diving" signs around above-ground pools.
- Allow only one person at a time on diving boards and waterslides.
- Ensure that swimmers stay away from the area underneath diving boards.

## Safe Boating

- Make sure that all boat occupants are wearing approved lifejackets or personal flotation devices (PFDs). Look for lifejackets or PFDs with labels stating that they have been approved by Transport Canada or the Canada Coast Guard.
- As a parent, guardian, or caregiver of a child, set a good example by wearing your lifejacket or PFD and by following boating safety rules.
- Small crafts can be unstable. Teach children to stay low in the boat.
- Teach children how to use a boat radio and fire extinguisher in case of an emergency.
- Discuss and practice survival techniques often.
- Choose PFDs or lifejackets carefully to match the wearer's size and weight.
- Children should use PFDs fitted with crotch straps. The strap must be used when wearing the PFD.
- All pleasure crafts, power vessels, sailing vessels, canoes, kayaks, and rowboats must carry safety equipment that is in good working order. To find out exactly what is needed for your boat, contact Transport Canada.
- As of September 15, 2009, anyone operating a powerboat on Canadian waters must have a pleasure craft operator card. Contact Transport Canada for more information on where to obtain your certification.

## Teaching Children Safety Awareness

Teach children how to get help in an emergency:

- Teach them to go to a trusted neighbour if they need help.
- Teach them about the Block Parent Program of Canada in your area, if applicable.

- Make a list of emergency phone numbers and teach children what to say when they call EMS/9-1-1:
  - Their name
  - The address they are calling from
  - What the problem is
  - The telephone number they are calling from
- Teach children to "hang up last" when they call EMS/9-1-1. This way the dispatcher will have all the necessary information.
- Teach children about traffic safety.

## Child and Youth Protection

There are different types of abuse. It is important to be able to understand the different situations in which abuse toward children and youth may occur. Many of these situations are preventable, and if you are able to recognize the signs, you may be able to intervene on a child or youth's behalf.

### Vulnerable Children and Youth

All children and youth live with some risk of experiencing abuse, violence, neglect, and bullying or harassment.

### Abuse and Violence

Abuse and violence may take different forms: emotional, physical, and sexual.

**Emotional abuse** is defined as chronic attacks on a child or youth's self-esteem.

**Physical abuse** occurs when a person in a position of power or trust purposefully injures or threatens to injure a child or youth.

**Sexual abuse** occurs when a younger or less powerful child is used by an older or more powerful child, adolescent, or adult for sexual gratification.

**Violence** is the intentional threat or use of physical force or power. It can be directed against oneself, another person, a group, or a community. It usually results in injury, death, psychological harm, abnormal development, or deprivation.

## Neglect

**Neglect** is the **chronic** inattention to the basic necessities of life, such as clothing, shelter, nutritious diet, education, good hygiene, supervision, medical and dental care, adequate rest, a safe environment, moral guidance and discipline, exercise, and fresh air.

## Understanding Bullying and Harassment

**Bullying** involves a person expressing his or her power through the humiliation of another person.

### Types of Bullying

- **Physical:** Hitting or kicking people, or taking or damaging their property or possessions.
- **Verbal:** Name-calling, insults, negative comments, and constant teasing.
- **Relational:** Trying to cut targets off from social connection by convincing peers to exclude or reject them.
- **Reactive:** Engaging in bullying, as well as provoking bullies into attacking, by taunting.
- **Cyberbullying:** Sending mean, cruel, and/or defamatory messages or images by electronic means (e.g., email, text messaging, instant messaging (IM), or personal and social websites).

## Harassment

According to the Canadian Human Rights Commission, discrimination is treating people "differently, negatively or adversely" on the basis of a prohibited ground of discrimination under federal, provincial, or territorial human rights legislation.

## Types of Harassment

- **Personal harassment:** Based on an individual's personal characteristics that are prohibited grounds for discrimination (e.g., racism).
- **Sexual harassment:** Unwelcome behaviour of a sexual or gender-based nature that negatively affects the person or the environment.
- **Criminal harassment:** When the harassing behaviour contravenes Canada's Criminal Code; also referred to as stalking.



Abuse is a protection issue.

Bullying is a relationship issue.

Harassment is a human rights issue.

All need intervention and support.

## Responding to Disclosures

A disclosure occurs when someone shares something with you in confidence. If someone discloses that they have experienced abuse, violence, neglect, or bullying, you must *always* act.

### How to Respond to Disclosures

If you are responding to a disclosure, you need to ask yourself three basic questions:

1. **What do I need to know?**
2. **What do I need to do?**
3. **When do I need to do it?**



Your jurisdiction's Child Protection Act may govern how you must respond to a disclosure.

When abuse is suspected or disclosed, you have a responsibility to **ACT**:

**A** **ACKNOWLEDGE** the child's situation and feelings.  
Access support

**C** **COMFORT** the child and take him or her to a safe place.  
Carefully listen to what the child says.

**T** **TAKE** notes and document what the child says and/or what you see. Take action—report the abuse immediately. Depending on the circumstances of the situation, use your judgment to decide if the child's parent or guardian should be contacted first (e.g., if he or she is being bullied or harassed by another child) or the Child Protection Authorities and police in your area (e.g., if the abuse is coming from an adult).

For more information on child and youth protection or to take a Respect Education course, please visit our website at [redcross.ca](http://redcross.ca).

**REMEMBER:** You do not have to be 100% certain that abuse has occurred. If you suspect it, report it. The safety of the child may be at risk. The authorities have the responsibility to determine the facts and evidence, not you.

## Children's Hygiene

### Diapering

Follow these steps to ensure that diapers are changed in a sanitary way:

- Use disposable gloves and remove them before sanitizing the area.
- Set up your changing area close to running water, away from the kitchen or eating area.
- Ensure that the changing surface and area is cleaned for at least 30 seconds with a bleach solution after each diapering.

- Dispose of dirty cloth diapers in a pail with a snug-fitting lid. Diaper pails and garbage cans need to be put out of children's reach.
- Place disposable diapers into the garbage can immediately.
- Avoid washing soiled clothes. Place them in a sealed plastic bag to be picked up by the parent or guardian at the end of the day.

## Dental Health

- Encourage children to brush their teeth every day.
  - A good time to brush is right after lunch. Use a pea-sized amount of toothpaste that contains fluoride.
  - Rinse the toothbrush well. When you put it away, make sure it is not touching other toothbrushes.
  - Replace the brush when the bristles become flattened.
  - Supervise young children when they are brushing.
- Children and babies' teeth can decay if they nap with a bottle because their soft teeth are constantly exposed to liquid with sugar. This is called nursing bottle syndrome, and it can be caused by:
  - Cow's milk
  - Juice
  - Formula
  - Sweetened liquids
- When children are between the ages of 1 and 2, encourage them to start drinking from a cup instead of from a bottle.
- Don't put honey, sugar, or syrup on soothers.
- Serve well-balanced meals. Too much sugary food is not healthy.
- Watch for the signs of tooth decay: dull, white or brown spots on the teeth.



## Example of an Emergency Supplies Kit

Have supplies ready for an emergency. Store them in a backpack or a duffle bag so you can take them with you if you have to evacuate the area. The following are examples of the items to include in your emergency supplies kit:

- 4 L (1 gal.) of water per person per day (use sealed, unbreakable containers and replace the supply every 6 months). Have enough for at least 3 days.
- Non-perishable packaged or canned food and a can opener. Replace the food once a year.
- Walking shoes, rain gear, and a change of clothing.
- Blankets or sleeping bags.
- A first aid kit and prescription medications (check the medications every 6 months to make sure they haven't expired).
- Toilet paper and other personal supplies.
- An extra pair of glasses.
- A battery-powered radio and flashlight, along with extra batteries.
- Spare cash.
- An extra set of car keys.
- A list of your family doctors.
- Important family information such as a list of any medical conditions or medical devices, including pacemakers.
- Photocopies of all important identification for you and your family, including health card numbers.
- Special items for babies, elderly, or disabled household members.
- Mobile phone and contact information for family and friends.



## Example of an Emergency Car Kit

Keep an emergency kit in your car. The following are examples of the items to include in your emergency car kit:

- A battery-powered radio and flashlight, with extra batteries
- A blanket
- Booster (jumper) cables
- A fire extinguisher
- A Canadian Red Cross first aid kit
- Bottled water and non-perishable high-energy foods (replace the water every 6 months and the food once a year)
- Maps of the area
- A shovel
- Flares
- A tire repair kit and pump
- Matches and a "survival" candle in a deep can that will burn for many hours

Generally speaking, you should avoid moving an injured or ill person to give care. Unnecessary movement can cause additional injury and pain and may complicate the person's recovery.

Three general situations require you to move a person:

1. There is an immediate danger (either a danger to you or to the person being rescued) e.g., from fire, a lack of oxygen, risk of drowning, a possible explosion, a collapsing structure, or uncontrolled traffic hazards.
2. You have difficulty accessing ill or injured people. For example, a person with minor injuries may need to be moved quickly to allow access to other people who may have life-threatening conditions.
3. There is a barrier that makes it difficult to provide proper care. For example, a person with a medical emergency, such as a cardiac arrest or heat stroke, may need to be moved to provide proper care. For example, a person in cardiac arrest requires CPR, which should be performed on a firm, flat surface. If the person collapses on a bed or in a small bathroom, the surface or space may not be adequate to provide appropriate care, and the person will need to be moved.

Usually, you will not face hazards that require you to move the person immediately. In most cases, you can give the necessary care wherever you find the person. Needless movement of a person can lead to further injury, and therefore should be avoided. For example, moving someone with a closed fracture of the leg, without taking the time to splint it, could result in an open fracture if the end of the bone tears the skin. This may result in soft-tissue damage, damage to nerves, blood loss, and infection.

Some rescue situations make gaining access to people especially challenging, and as such require first responders who have specialized training and equipment. One such situation is when the ill or injured person is in a confined space. Confined spaces are particularly challenging because they:

- Are not designed or intended to have people in them
- Have restricted entrances or exits due to their location, size, or structure
- Can present a risk to the health and safety of anyone who enters (including first responders), due to one or more of the following factors:
  - Their design, construction, location, or atmosphere
  - The materials or substances inside
  - Work activities being carried out inside

Before you act, you must consider the limitations of the situation. Considering the following factors will help you ensure that you move a person quickly and safely:

- Dangerous conditions at the scene
- The size of the person
- Your own physical ability
- Whether others can assist you
- The person's condition

Failing to consider these factors could result in an injury to the First Aider—you. If you become injured, you may be unable to move the person and may risk making the situation worse. Back injuries (and other musculoskeletal injuries) can occur as a result of lifting and carrying a great deal of equipment during an emergency. If this happens at a scene, you will not be able to help and another First Aider will have to provide care for you, in addition to the other ill or injured person/people.

To protect yourself and the ill or injured person, follow these guidelines when moving someone:

- Attempt to move a person only when you are sure you can comfortably handle the rescue.
- Walk carefully, using short steps.
- Walk forward with the person, instead of backward, wherever possible.
- Always take the shortest, most direct route to your destination, as long as it is safe.
- Scan the pathway you want to use before moving so that you can identify potential hazards (such as slip hazards or poor lighting).



There are specific ways to deal with confined spaces (e.g., a grain silo, vat, or pipe). Without appropriate training, you are putting yourself at risk if you enter them. Ensure that the appropriate responders are on the way and manage the scene until they arrive. Only people with specialized training and equipment should enter confined spaces.

## Body Mechanics

To reduce the risk of personal injury, everyone involved in a lift or carry should use proper body mechanics (also called “biomechanics”). The basic principles of body mechanics that can be used for all lifts and moves include the following:

- Use your legs, not your back, to lift. When lifting, use the muscles of your legs, hips, buttocks, and abdomen. Never use the muscles of your back to move or lift a heavy load.
- Keep the weight as close to you as possible. Reduce the distance you have to reach.
- Keep your body aligned. Imagine a straight line running from your shoulders through your hips and down to your feet, and then move

them as a unit. This will reduce twisting forces.

- Reduce the height or distance you need to move a weight and lift in stages, if necessary.
- Keep your wrists and knees in normal alignment.

If performed incorrectly, reaching for a person or object can also injure the ligaments of your back. When reaching, keep your back locked (avoid hyperextending it) and never twist your back. You should avoid reaching more than 15 to 20 centimetres (6 to 8 in.) in front of the body as the muscles in your upper back and shoulders can only stay stretched in that position for a matter of seconds before they become fatigued and the risk of injury increases.

## How to Move an Ill or Injured Person

There are many different ways to move someone to safety. The move is successful as long as you can move the person without injuring yourself, causing further injury to the person, or taking unnecessary risks.

Assists, carries, and drags are the different ways you can safely move an injured person. The most common of these moves include the following:

- Walking assist
- Two-person seat carry
- Clothes drag
- Blanket drag
- Extremity lift

These moves can be performed by one or two people without specialized equipment. This is important because equipment is often not immediately available and time is usually critical when a person must be moved.

## Walking Assist

The most basic move is the walking assist. It is used to help a responsive person who requires simple assistance with walking. Either one or two First Aiders can use this method.

To perform a walking assist:

1. Have the person stand up and position yourself on the person's weaker or injured side.
2. Place the person's arm across your shoulders and hold it in place with one hand.
3. Place your other hand around the person's waist. Your body acts as a "crutch," supporting the person's weight while you both walk.
4. A second First Aider can support the person in the same way from the other side.



## Two-Person Seat Carry

The two-person seat carry is a method for moving a person who cannot walk and who is not likely to have a head and/or spinal injury. It requires two First Aiders.

To perform the two-person seat carry:

1. Stand on one side of the person and have the other First Aider stand on the opposite side.
2. Put one arm under the person's thighs and the other across the person's back, while the other First Aider does the same.
3. Grab and hold the wrists of the second First Aider under the person's legs and behind the person's back while the second First Aider does the same to you.
4. Lift the person in the "seat" formed by your joined arms.



## Clothes Drag

The clothes drag is an emergency move that is appropriate for someone suspected of having a head and/or spinal injury, as it helps keep the head and neck stabilized.

To perform a clothes drag:

1. Gather the person's clothing, such as a jacket or shirt, behind the person's neck.
2. Cradle the person's head using both his or her clothing and your hands, and keep the person's head, neck, and back as straight as possible.
3. Pull the person to safety.



This type of emergency move is exhausting and may result in back strain for the First Aider, even when done properly.

## Blanket Drag

The blanket drag is an appropriate emergency move if you do not have a stretcher.

To perform a blanket drag:

1. Kneel beside the person and spread out a blanket on the person's other side.
2. Gather half of the blanket and place it up against the person's side.
3. Roll the person toward you, moving the person's body as one unit.
4. Tuck the gathered half of the blanket under the person as far as you can, and then roll the person back onto the blanket.
5. Wrap the blanket around the person and then drag him or her, holding the blanket around the person's head.



## Extremity Lift

The extremity lift (also called a fore-and-aft lift) is performed with a partner. The extremity lift is not appropriate for a person with a suspected head or spine injury, or with injuries to the pelvis, arms, or legs. This move can be used to lift an unresponsive person from the floor to a chair.

To perform an extremity lift:

1. Crouch at the person's head. Have the other First Aider kneel, either beside or between the person's knees.
2. Place one hand under each of the person's shoulders and reach through to grab the person's wrists. Ensure that the person's back is close to your chest.
3. Have the other First Aider slip his or her hands under the person's knees.
4. On your signal, both of you should lift simultaneously.



## Stretchers and Lifting Devices

There are many kinds of stretchers and lifting devices. The more common types are:

- Scoop stretcher (clamshell)
- Backboard
- Basket stretcher
- Multi-level stretcher (wheeled cot)
- Flexible litter
- Stair chair

Any of these stretchers will carry a person's entire body.



### Scoop Stretcher

The scoop stretcher (also called a clamshell) is a lifting device suitable for lifting a person from the ground to another kind of stretcher or backboard. The scoop stretcher is only used for lifting and is not intended for carrying people across long distances. It allows people to be lifted without moving or rolling them. One disadvantage is that the person's back cannot be examined unless the person is rolled.

### Backboard

The backboard (also referred to as the spine board or full board) is only used for lifting and is not designed to carry people across long distances. It is constructed of either plywood or plastic. Backboards are either rectangular in shape or tapered at one end. The board has holes cut along the sides through which straps may be placed to secure the person's body to the board. Most boards are approximately 1.8 metres (6 feet) long. If a backboard is used with a basket stretcher, the board must fit inside the stretcher.



### Basket Stretcher

The basket stretcher, also called the Stokes basket, comes in various types and strengths. Traditional basket stretchers are constructed of metal support frames, usually combined with a wire mesh lining. Other basket stretchers have metal frames and high-impact plastic liners. Basket stretchers are commonly used for rescues and also as lifting devices.

Follow these guidelines when using a basket stretcher:

- Place blankets under the person for comfort.
- Place a backboard or litter board on the bottom to make loading and unloading easier.
- If used in combination with a scoop stretcher, the basket stretcher must be able to accommodate the scoop stretcher.
- Secure the person if the stretcher is to be carried over a long distance or the person is to be lowered or raised off the ground as in a low- or high-angle rescue.





### Multi-Level Stretcher

This type of stretcher can be lowered or raised using release handles found at the end or side. This device is equipped with wheels. It has an adjustable head section and leg section. Each side of the stretcher has a safety rail. The stretcher should not be lifted using these rails. This stretcher is commonly used in transportation vehicles, such as ambulances.

### Flexible Litter

A flexible litter has no rigid structure of its own. These devices are made of synthetic materials that require some type of spinal immobilization device to provide rigidity. Because they are flexible, they work well when moving people through narrow passageways. They wrap around the person, so the litter is little more than the circumference of the person's body. Flexible litters also have the distinct advantage of being usable with people of a variety of sizes, including those for whom a basket stretcher may be restrictive.



### Stair Chair

A stair chair is used for transporting a person in a sitting position. It is used when a wheeled stretcher is deemed to be too long for the rescue or extrication. It is especially useful when there is a small elevator or staircase in which a long stretcher will not fit. The front wheels swivel to manoeuvre around tight corners and landings. Some stair chairs have a track that make navigating stairs or steep terrains easier for the First Aiders. When using the stair chair, it is recommended that three First Aiders be present to ensure a person's safety: two First Aiders to act as carriers and one to serve as a spotter for potential difficulties.

As a First Aider, you need a basic understanding of normal human structure and function. Knowing the body's structures and how they work will help you more easily recognize and understand illnesses and injuries. Body systems do not function independently. Each system depends on other systems to function properly. When your body is healthy, your body systems are working well together. But an injury or illness in one body part or system will often cause problems in others. Knowing the location and function of the major organs and structures within each body system will help you to more accurately assess a patient's condition and provide the best care.

### Body Systems

The human body performs many complex functions. Vital organs are organs whose functions are essential for life, such as the brain, heart, and lungs.

A body system is a group of organs and other structures that are especially adapted to perform specific body functions. They work together to carry out a function needed for life. For example, the heart, blood, and blood vessels make up the circulatory system, which keeps all parts of the body supplied with oxygen-rich blood.

For the body to work properly, all of the body's systems must work well together. When one or more of the body systems does not perform its function properly, illness can occur.

### Interrelationships of Body Systems

Each body system plays a vital role in survival. Body systems work together to help the body maintain a constant healthy state. When the environment changes, the body's systems adapt to the new conditions. For example, because the musculoskeletal system works harder during exercise, the respiratory and circulatory systems must also work harder to meet the body's increased oxygen demands. Your body systems also react to the stresses caused by illness or injury.

The impact of an injury or a disease is rarely restricted to one body system. For example, a broken bone may result in nerve damage that may impair movement and feeling. Injuries to the ribs can make breathing difficult. If the heart stops beating for any reason, breathing will also stop.



In any significant illness or injury, body systems may be seriously affected. This may result in a progressive failure of multiple body systems. Generally, the more body systems that are involved in an emergency, the more serious the emergency.

**BODY SYSTEMS**


<b>System</b>	<b>Major Structures</b>	<b>Primary Functions</b>	<b>Relationship with Other Body Systems</b>
<b>Respiratory System</b>	Airway and lungs	Supplies the body with oxygen through breathing	Works with the circulatory system to provide oxygen to cells; is under the control of the nervous system
<b>Circulatory System</b>	Heart, blood, and blood vessels	Transports nutrients and oxygen to body cells and removes waste products	Works with the respiratory system to provide oxygen to cells; works with the urinary and digestive systems to remove waste products; helps give skin its colour; is under the control of the nervous system
<b>Lymphatic System</b>	Lymph, lymph nodes	Removes excess fluid (lymph) from body tissues; absorbs fatty acids and transports fat to the circulatory system; transports white blood cells around the body and initiates the formation of antibodies	Works to return fluid to the circulatory system
<b>Nervous System</b>	Brain, spinal cord, and nerves	Transmits messages to and from the brain	Regulates all body systems through a network of nerves
<b>Musculo-skeletal System</b>	Bones, ligaments, muscles, and tendons	Provides the body's framework; protects internal organs and other underlying structures; allows movement; produces heat; manufactures the components of blood	Provides protection to organs and structures of other body systems; muscle action is controlled by the nervous system

**BODY SYSTEMS**


<b>System</b>	<b>Major Structures</b>	<b>Primary Functions</b>	<b>Relationship with Other Body Systems</b>
<b>Integumentary System</b>	Skin, hair, and nails	An important part of the body's communication network; helps prevent infection and dehydration; assists with temperature regulation; aids in production of certain vitamins	Helps protect the body from disease-producing organisms; together with the circulatory system, helps regulate body temperature; communicates sensation to the brain by way of the nerves
<b>Endocrine System</b>	Glands	Secretes hormones and other substances into blood and onto skin	Together with the nervous system, coordinates the activities of other systems
<b>Digestive System</b>	Mouth, esophagus, stomach, intestines	Breaks down food into a usable form to supply the body with nutrients and energy	Works with the circulatory system to transport nutrients to the body
<b>Genitourinary System</b>	Uterus and genitalia	Performs the processes of reproduction	Regulated through hormones produced by the endocrine system
	Kidneys, bladder	Removes wastes and regulates water balance	Receives waste from the circulatory system




## The Abdominal Quadrants




## Respiratory System




## Circulatory System




## Nervous System




## Musculoskeletal System Muscles




## Musculoskeletal System Skeleton




## Integumentary System



## Digestive System



## Urinary System



## Reproductive System

## A

**Abdomen:** The part of the body below the chest and above the pelvis. It contains the stomach, intestines, liver, spleen, and other organs.

**Abdominal thrusts:** A method to remove a foreign object from the airway.

**Airway:** The pathway that allows air to travel from the mouth and nose to the lungs.

**Airway obstruction:** A blockage in the airway that stops air from reaching the lungs.

**Allergic reaction:** The body's response to a substance to which it is particularly sensitive. The response can be mild or very severe.

**Anemia:** A condition caused by a lack of red blood cells.

**Arteries:** Large blood vessels that carry oxygen-rich blood from the heart to the rest of the body.

**Aspiration:** Inhaling blood, vomit, saliva, or foreign material into the lungs.

**Asthma:** A chronic illness in which certain substances or conditions (triggers) cause inflammation and swelling of the bronchioles, making breathing difficult.

**Automated external defibrillator (AED):** An electronic device that analyzes the heart's electrical rhythm and, if necessary, tells the user to deliver a shock to a person in cardiac arrest.

## B

**Baby:** A child under the age of 12 months.

**Bandage:** Material used to wrap or cover a part of the body or to hold a dressing or splint in place.

**Breathing emergency:** A situation in which breathing is so impaired that the person's life is in danger.

**Bronchioles:** Small tubes at the base of the lungs where oxygen and carbon dioxide are exchanged.

**Burn:** An injury caused by heat, chemicals, electricity, or radiation.

**Bystander:** Someone who is present at the scene of a situation or emergency (other than an ill or injured person).

## C

**Cardiac arrest:** A condition in which the heart has stopped beating or beats too irregularly or too weakly to pump blood effectively.

**Cardiopulmonary resuscitation (CPR):** A first aid technique that combines rescue breaths and chest compressions for someone who has stopped breathing and whose heart has stopped beating.

**Cardiovascular disease:** Any disease of the heart and blood vessels (also called heart disease).

**Caustic:** Capable of causing chemical burns

**Child:** Anyone between the ages of 1 and 8 years. (When giving CPR or using an AED, a child is considered to be anyone between 1 year old and the onset of puberty.)

**Choking:** A condition in which someone's airway is partly or completely blocked by a foreign object, swelling of the mouth or throat, or fluids such as vomit.

**Cholesterol:** A fatty substance that can cause build-up on artery walls, making the arteries narrower and restricting blood flow.

**Compression:** Rhythmic pressure that is put on the chest to dislodge something blocking the airway or to circulate blood when the heart isn't beating effectively.

**Concussion:** A temporary impairment of brain function, usually caused by a blow to the head.

**Contraction:** A squeezing action made by the muscles in the womb during labour.

**Crowning:** The point in labour when the baby's head is at the opening of the vagina.

## D

**Defibrillation:** An electric shock that is given to correct a life-threatening heart rhythm.

**Direct pressure:** Pressure that is put on a wound to control bleeding.

**Dislocation:** An injury in which a bone is moved out of its normal position at a joint.

**Dispatcher:** The emergency medical services worker who answers the EMS/9-1-1 telephone number and decides which EMS personnel to send to the scene and who may give advice about first aid until EMS personnel arrive.

**Dressing:** A pad placed directly over a wound to absorb blood and other body fluids and to prevent infection.

## E

**Emergency medical services (EMS) personnel:** Trained and equipped people, including police, firefighters, and ambulance personnel, who are dispatched through a local emergency number to give emergency care to ill or injured people.

**Epinephrine:** A drug that can be injected into the body to counteract a severe allergic reaction.

**External bleeding:** Bleeding from an open wound in the skin.

## F

**First Aider:** A person with training who gives immediate care to someone who is ill or injured until more advanced care can be obtained.

**Foreign object:** Any item that enters the body from outside.

**Fracture:** A break, crack, or chip in bone tissue.

## H

**Head-tilt/chin-lift:** A technique for opening the airway in an unresponsive adult, child, or baby.

**Heart attack:** A sudden illness in which an artery that feeds the heart becomes blocked, stopping part of the heart from getting the oxygen-rich blood it needs.

**Hypothermia:** A life-threatening condition that develops when the body's temperature drops too low, usually from being exposed to cold temperatures for too long.

## I

**Immobilize:** To use a splint or other method to keep an injured body part from moving.

**Internal bleeding:** Bleeding that occurs inside the body.

**J**

**Joint:** The point where two or more bones are held together by ligaments that allow movement.

**L**

**Ligament:** A fibrous band that holds bones together at a joint.

**M**

**Medical identification product:** A wallet card, bracelet, watch strap, anklet, or necklace with a tag indicating that the person wearing it has a particular medical condition.

**P**

**Pelvis:** The part of the body between the abdomen and the legs. It contains the intestines, bladder, and reproductive organs.

**Poison:** Any substance that causes injury, illness, or death if it is inhaled, swallowed (ingested), absorbed, or injected.

**Poison Control Centre:** A centre staffed by medical professionals who can give information about first aid in cases of poisoning.

**Primary assessment:** An examination of an ill or injured person to determine his or her initial condition and whether there are any life-threatening concerns.

**Pulse:** The beat felt in arteries near the skin with each contraction of the heart.

**R**

**Recovery position:** A position for an unresponsive person that helps to keep the airway open and allows any blood or vomit to drain from the person's mouth.

**Respiratory arrest:** A condition in which breathing has stopped or isn't effective.

**Respiratory distress:** A condition in which breathing is difficult.

**Responsiveness:** Awareness of the environment and ability to respond to stimuli.

**Risk factors:** Conditions or behaviours that increase the chance that a person will develop a particular disease or suffer a particular injury.

**S**

**Secondary assessment:** A verbal, visual, and/or physical check of an ill or injured person for conditions that were not identified in the primary assessment.

**Seizure:** An episode of abnormal electrical signals in the brain that results in disturbed brain function, shaking or contraction of limbs, and/or an altered level of responsiveness.

**Shock:** A serious condition caused when the circulatory system cannot get enough oxygen-rich blood to all parts of the body. Causes include severe blood loss, allergic reactions, and emotional trauma.

**Sign:** An indicator of injury or illness that a First Aider can see, feel, smell, or hear.

**Soft tissue:** Skin, fat, muscles, and other soft body structures.

**Splint:** A device used to stop body parts from moving.

**Sprain:** The stretching or tearing of ligaments and other soft tissues at a joint.

**Strain:** The stretching or tearing of muscles and tendons.

**Stroke:** A disruption of blood flow in the brain, causing weakness and/or speech problems.

**Symptom:** An indicator of injury or illness that is felt and expressed by the ill or injured person.

## T

**Tendon:** A fibrous band that attaches muscle to bone.

**Tissue:** A group of cells that work together to perform specific functions.

**Tuberculosis (TB):** A potentially fatal bacterial infection of the lungs that is spread through the air from one person to another. It can also affect the bones, brain, kidneys, and other organs. Treatment is complex and involves taking many different medications over an extended period of time.

## U

**Unresponsive:** A state in which a person is unaware of the environment and does not react to stimuli.

## V

**Veins:** Blood vessels that carry oxygen-poor blood from all parts of the body back to the heart.

**Vertebrae:** The 33 bones that make up the spine.

**Vital signs:** Three key characteristics of a person's condition (level of responsiveness, breathing, and skin).

## W

**Wound:** An injury to soft tissues.

Each time a program is revised, it is built on the great work completed in the previous revisions. The Canadian Red Cross would like to recognize everyone who worked on developing these programs in the past; their work set the foundation for our success.

We would like to thank our Training Partners, Master Instructor Trainers, Instructor Trainers, and Instructors who provided the feedback that helped guide this revision and shape our new programs.

This revision required the hard work and dedication of many teams who put in countless hours to contribute to its success. The Canadian Red Cross extends its thanks to the First Aid Technical Advisory Group (FATAG) for its overall leadership, dedication, and direction. FATAG members include:

Dr. Andrew MacPherson  
*Medical Director*

Jason Brinson

Richard Czech

Blair Doyle

Jason Durham

Domenic Filippelli

Dr. Gordon Giesbrecht

Dr. Morgan Hillier

Carolyn Hoekstra

Kevin Holder

Dr. Loriann Hynes

Lyle Karasiuk

Sioban Kennedy

Shelly Longmore

Jodie Marshall

Roger Mayo

Kyle Mohler

Michael Nemeth

Tyrone Power

Bob Reside

Kathy Sampson

Mike Skinner

Diane Story

Hugo Surprenant

Kristopher Tharris

Debbie Van't Kruis

Dave Wooldridge

Charna Young

*Illustrations by Allison MacAlister*

*Photos by Matt Bork, Tara Hillyer, and Lisa Petrole*

94 bruise; © Shutterstock.com/Stephane Bidouze

155, 158 child with medication; © Shutterstock.com/ Thomas M Perkins

161, 163 poison oak; © iStockphoto.com/Devin Allphin

161, 163 poison sumac; © iStockphoto.com/zorani

162, 163 Giant Hogweed ©Rob Paxton

166 raccoon; © bookguy/iStock/Thinkstock

167 bee; © ThinkstockPhotos.ca/Ale-ks<<http://www.thinkstockphotos.ca/search/2/image?artist=Ale-ks&family=creative>>

170, 171 rattlesnake; © iStockphoto.com/Dave Rodriguez

171 Massasauga snake; © Reptilia Zoo and Education Centre

171 Prairie Rattlesnake; © Reptilia Zoo and Education Centre

172 black widow spider; © iStockphoto.com/Mark Kostich

172 brown recluse spider; © iStockphoto.com/Clint Spencer

173 sting ray; © iStockphoto.com/Niko Guido

173 jellyfish; © The Vancouver Aquarium

188 taking temperature; © Shutterstock.com/SpeedKingz<<https://www.shutterstock.com/g/SpeedKingz>>

190 checking medication; © Shutterstock.com/ bikeriderlondon<<https://www.shutterstock.com/g/bikeriderlondon>>

# Stay Connected.

[myrc.redcross.ca](http://myrc.redcross.ca)

Log in for program standards, teaching resources and forms, FAQs, contact with technical experts, and much more.



[shop.redcross.ca](http://shop.redcross.ca)

Visit your online storefront for course materials, training equipment, and gifts or promotional items.



[redcross.ca/apps](http://redcross.ca/apps)

Download the free First Aid App and get instant access to videos, interactive quizzes, and simple, step-by-step advice.



[redcross.ca](http://redcross.ca)

Stay tuned for training and volunteer opportunities in the Red Cross community.

**Red Cross First Aid. The Experience to Make a Difference.**



[redcross.ca/firstaid](http://redcross.ca/firstaid) | 1-877-356-3226

Copyright © 2017 The Red Cross Society  
ISSN: 978-1-55104-709-6